

Eun Seok Kang

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/2569097/eun-seok-kang-publications-by-year.pdf>
Version: 2024-04-04

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

237 papers	5,751 citations	38 h-index	65 g-index
246 ext. papers	6,928 ext. citations	5.3 avg, IF	5.51 L-index

#	Paper	IF	Citations
237	Severe Hypoglycemia Increases Dementia Risk and Related Mortality: A Nationwide, Population-Based Cohort Study.. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2022 ,	5.6	2
236	DA-1241, a Novel GPR119 Agonist, Improves Hyperglycaemia by Inhibiting Hepatic Gluconeogenesis and Enhancing Insulin Secretion in Diabetic Mice.. <i>Diabetes and Metabolism Journal</i> , 2022 ,	5	1
235	Blood Pressure Levels and Risks of Dementia: a Nationwide Study of 4.5 Million People. <i>Hypertension</i> , 2022 , 79, 218-229	8.5	0
234	Analysis of Severe Hypoglycemia Among Adults With Type 2 Diabetes and Nonalcoholic Fatty Liver Disease.. <i>JAMA Network Open</i> , 2022 , 5, e220262	10.4	0
233	Ezetimibe combination therapy with statin for non-alcoholic fatty liver disease: an open-label randomized controlled trial (ESSENTIAL study).. <i>BMC Medicine</i> , 2022 , 20, 93	11.4	4
232	Short Term Isocaloric Ketogenic Diet Modulates NLRP3 Inflammasome B-hydroxybutyrate and Fibroblast Growth Factor 21.. <i>Frontiers in Immunology</i> , 2022 , 13, 843520	8.4	2
231	Sodium Glucose Cotransporter-2 Inhibitors as an Add-on Therapy to Metformin Plus Dipeptidyl Peptidase-4 Inhibitor in Patients with Type 2 Diabetes. <i>Yonsei Medical Journal</i> , 2022 , 63, 539	3	
230	Proteinuria as a significant predictive factor for the progression of carotid artery atherosclerosis in non-albuminuric type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2021 , 181, 109082	7.4	
229	Association between domain-specific physical activity and diabetes in Korean adults. <i>Scientific Reports</i> , 2021 , 11, 13066	4.9	1
228	Cardiovascular Risk Is Elevated in Lean Subjects with Nonalcoholic Fatty Liver Disease. <i>Gut and Liver</i> , 2021 ,	4.8	4
227	Safety and effectiveness of linagliptin in Korean patients with type 2 diabetes: A postmarketing surveillance study. <i>Diabetes, Obesity and Metabolism</i> , 2021 , 23, 1208-1212	6.7	0
226	Ipragliflozin, an SGLT2 Inhibitor, Ameliorates High-Fat Diet-Induced Metabolic Changes by Upregulating Energy Expenditure through Activation of the AMPK/ SIRT1 Pathway. <i>Diabetes and Metabolism Journal</i> , 2021 , 45, 921-932	5	7
225	Renal Tubular Damage Marker, Urinary N-acetyl-D-Glucosaminidase, as a Predictive Marker of Hepatic Fibrosis in Type 2 Diabetes Mellitus. <i>Diabetes and Metabolism Journal</i> , 2021 ,	5	1
224	Non-Laboratory-Based Simple Screening Model for Nonalcoholic Fatty Liver Disease in Patients with Type 2 Diabetes Developed Using Multi-Center Cohorts. <i>Endocrinology and Metabolism</i> , 2021 , 36, 823-834	3.5	1
223	Nonalcoholic fatty liver disease, diastolic dysfunction, and impaired myocardial glucose uptake in patients with type 2 diabetes. <i>Diabetes, Obesity and Metabolism</i> , 2021 , 23, 1041-1051	6.7	1
222	Uric Acid Variability as a Predictive Marker of Newly Developed Cardiovascular Events in Type 2 Diabetes.. <i>Frontiers in Cardiovascular Medicine</i> , 2021 , 8, 775753	5.4	0
221	Non-alcoholic steatohepatitis and progression of carotid atherosclerosis in patients with type 2 diabetes: a Korean cohort study. <i>Cardiovascular Diabetology</i> , 2020 , 19, 81	8.7	11

220	Current and emerging pharmacological options for the treatment of nonalcoholic steatohepatitis. <i>Metabolism: Clinical and Experimental</i> , 2020 , 111S, 154203	12.7	46
219	Comparison of Renal Effects of Ezetimibe-Statin Combination versus Statin Monotherapy: A Propensity-Score-Matched Analysis. <i>Journal of Clinical Medicine</i> , 2020 , 9,	5.1	2
218	Ipragliflozin Additively Ameliorates Non-Alcoholic Fatty Liver Disease in Patients with Type 2 Diabetes Controlled with Metformin and Pioglitazone: A 24-Week Randomized Controlled Trial. <i>Journal of Clinical Medicine</i> , 2020 , 9,	5.1	17
217	Proteinuria is Associated with Carotid Artery Atherosclerosis in Non-Albuminuric Type 2 Diabetes: A Cross-Sectional Study. <i>Journal of Clinical Medicine</i> , 2020 , 9,	5.1	3
216	SGLT2 inhibition modulates NLRP3 inflammasome activity via ketones and insulin in diabetes with cardiovascular disease. <i>Nature Communications</i> , 2020 , 11, 2127	17.4	96
215	Cohort Profile: Firefighter Research on the Enhancement of Safety and Health (FRESH), a Prospective Cohort Study on Korean Firefighters. <i>Yonsei Medical Journal</i> , 2020 , 61, 103-109	3	8
214	Association between Non-Alcoholic Steatohepatitis and Left Ventricular Diastolic Dysfunction in Type 2 Diabetes Mellitus. <i>Diabetes and Metabolism Journal</i> , 2020 , 44, 267-276	5	13
213	Effect of dapagliflozin, a sodium-glucose co-transporter-2 inhibitor, on gluconeogenesis in proximal renal tubules. <i>Diabetes, Obesity and Metabolism</i> , 2020 , 22, 373-382	6.7	12
212	Efficacy and safety of gemigliptin as add-on therapy to insulin, with or without metformin, in patients with type 2 diabetes mellitus (ZEUS II study). <i>Diabetes, Obesity and Metabolism</i> , 2020 , 22, 123-127	6.7	3
211	Efficacy and safety of insulin glargine 300 U/mL versus insulin glargine 100 U/mL in Asia Pacific insulin-naïve people with type 2 diabetes: The EDITION AP randomized controlled trial. <i>Diabetes, Obesity and Metabolism</i> , 2020 , 22, 612-621	6.7	6
210	Dipeptidyl peptidase-4 inhibitor protects against non-alcoholic steatohepatitis in mice by targeting TRAIL receptor-mediated lipoapoptosis via modulating hepatic dipeptidyl peptidase-4 expression. <i>Scientific Reports</i> , 2020 , 10, 19429	4.9	4
209	Effect of Ezetimibe on Glucose Metabolism and Inflammatory Markers in Adipose Tissue. <i>Biomedicines</i> , 2020 , 8,	4.8	3
208	Effect of Switching from Linagliptin to Tenueligliptin Dipeptidyl Peptidase-4 Inhibitors in Older Patients with Type 2 Diabetes Mellitus. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2020 , 13, 4113-4121	3.4	2
207	Association Between Serum Bilirubin and the Progression of Carotid Atherosclerosis in Type 2 Diabetes. <i>Journal of Lipid and Atherosclerosis</i> , 2020 , 9, 195-204	3	2
206	Commentary: Nonalcoholic or metabolic dysfunction-associated fatty liver disease? The epidemic of the 21st century in search of the most appropriate name. <i>Metabolism: Clinical and Experimental</i> , 2020 , 113, 154413	12.7	16
205	Hepatic fibrosis is associated with total proteinuria in Korean patients with type 2 diabetes. <i>Medicine (United States)</i> , 2020 , 99, e21038	1.8	2
204	Differential Diabetogenic Effect of Pitavastatin and Rosuvastatin, in vitro and in vivo. <i>Journal of Atherosclerosis and Thrombosis</i> , 2020 , 27, 429-440	4	3
203	Gamma glutamyltransferase and risk of dementia in prediabetes and diabetes. <i>Scientific Reports</i> , 2020 , 10, 6800	4.9	5

202	Nonalcoholic Fatty Liver Disease and Sarcopenia Are Independently Associated With Cardiovascular Risk. <i>American Journal of Gastroenterology</i> , 2020 , 115, 584-595	0.7	24
201	Risk of Incident Dementia According to Metabolic Health and Obesity Status in Late Life: A Population-Based Cohort Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019 , 104, 2942-2952	5.6	21
200	Incident Hepatocellular Carcinoma Risk in Patients Treated with a Sulfonylurea: A Nationwide, Nested, Case-Control Study. <i>Scientific Reports</i> , 2019 , 9, 8532	4.9	3
199	Predictors of the Therapeutic Efficacy and Consideration of the Best Combination Therapy of Sodium-Glucose Co-transporter 2 Inhibitors. <i>Diabetes and Metabolism Journal</i> , 2019 , 43, 158-173	5	10
198	Low-dose pioglitazone can ameliorate learning and memory impairment in a mouse model of dementia by increasing LRP1 expression in the hippocampus. <i>Scientific Reports</i> , 2019 , 9, 4414	4.9	33
197	Reduced expression of pyruvate kinase in kidney proximal tubule cells is a potential mechanism of pravastatin altered glucose metabolism. <i>Scientific Reports</i> , 2019 , 9, 5318	4.9	1
196	Prevention of Hepatocellular Carcinoma by Statins: Clinical Evidence and Plausible Mechanisms. <i>Seminars in Liver Disease</i> , 2019 , 39, 141-152	7.3	9
195	Statin Use is Associated with Decreased Hepatocellular Carcinoma Recurrence in Liver Transplant Patients. <i>Scientific Reports</i> , 2019 , 9, 1467	4.9	11
194	Response: Predictors of the Therapeutic Efficacy and Consideration of the Best Combination Therapy of Sodium-Glucose Co-transporter 2 Inhibitors (2019;43:158-73). <i>Diabetes and Metabolism Journal</i> , 2019 , 43, 379-380	5	1
193	Predictive factors for the development of diabetes in cancer patients treated with phosphatidylinositol 3-kinase inhibitors. <i>Cancer Chemotherapy and Pharmacology</i> , 2019 , 84, 405-414	3.5	5
192	Glucometabolic characteristics and higher vascular complication risk in Korean patients with type 2 diabetes with non-albumin proteinuria. <i>Journal of Diabetes and Its Complications</i> , 2019 , 33, 585-591	3.2	2
191	The Effectiveness of Intermittent Fasting to Reduce Body Mass Index and Glucose Metabolism: A Systematic Review and Meta-Analysis. <i>Journal of Clinical Medicine</i> , 2019 , 8,	5.1	48
190	Nationwide Trends in Pancreatitis and Pancreatic Cancer Risk Among Patients With Newly Diagnosed Type 2 Diabetes Receiving Dipeptidyl Peptidase 4 Inhibitors. <i>Diabetes Care</i> , 2019 , 42, 2057-2064	14.6	18
189	2019 Clinical Practice Guidelines for Type 2 Diabetes Mellitus in Korea. <i>Diabetes and Metabolism Journal</i> , 2019 , 43, 398-406	5	124
188	Spontaneous ketonuria and risk of incident diabetes: a 12-year prospective study. <i>Diabetologia</i> , 2019 , 62, 779-788	10.3	5
187	Current Management of Type 2 Diabetes Mellitus in Primary Care Clinics in Korea. <i>Endocrinology and Metabolism</i> , 2019 , 34, 282-290	3.5	9
186	Injectable Therapy for Diabetes Mellitus: Glucagon-Like Peptide-1 Receptor Agonist. <i>Journal of Korean Diabetes</i> , 2019 , 20, 149	0.1	
185	Efficacy and Safety of Gemigliptin in Post-Transplant Patients With Type 2 Diabetes Mellitus. <i>Transplantation Proceedings</i> , 2019 , 51, 3444-3448	1.1	1

184	Differential Effects of Thiazolidinediones and Dipeptidyl Peptidase-4 Inhibitors on Insulin Resistance and β -Cell Function in Type 2 Diabetes Mellitus: A Propensity Score-Matched Analysis. <i>Diabetes Therapy</i> , 2019 , 10, 149-158	3.6	4
183	Sodium-glucose cotransporter 2 inhibitors regulate ketone body metabolism via inter-organ crosstalk. <i>Diabetes, Obesity and Metabolism</i> , 2019 , 21, 801-811	6.7	20
182	Teneligliptin versus sitagliptin in Korean patients with type 2 diabetes inadequately controlled with metformin and glimepiride: A randomized, double-blind, non-inferiority trial. <i>Diabetes, Obesity and Metabolism</i> , 2019 , 21, 631-639	6.7	5
181	Comparison of hepatocellular carcinoma risk between patients treated with glimepiride and gliclazide. <i>Diabetes and Metabolism</i> , 2019 , 45, 83-85	5.4	4
180	The association between skin auto-fluorescence of palmoplantar sites and microvascular complications in Asian patients with type 2 diabetes mellitus. <i>Scientific Reports</i> , 2018 , 8, 6309	4.9	2
179	Statin use and the risk of hepatocellular carcinoma in patients at high risk: A nationwide nested case-control study. <i>Journal of Hepatology</i> , 2018 , 68, 476-484	13.4	63
178	Waist-to-calf circumference ratio is an independent predictor of hepatic steatosis and fibrosis in patients with type 2 diabetes. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2018 , 33, 1082-1091	4	11
177	Pharmacotherapy of type 2 diabetes: An update. <i>Metabolism: Clinical and Experimental</i> , 2018 , 78, 13-42	12.7	103
176	Combining SGLT2 Inhibition With a Thiazolidinedione Additively Attenuate the Very Early Phase of Diabetic Nephropathy Progression in Type 2 Diabetes Mellitus. <i>Frontiers in Endocrinology</i> , 2018 , 9, 412	5.7	17
175	Comparison of the Effects of Ezetimibe-Statin Combination Therapy on Major Adverse Cardiovascular Events in Patients with and without Diabetes: A Meta-Analysis. <i>Endocrinology and Metabolism</i> , 2018 , 33, 219-227	3.5	9
174	Differential association of ezetimibe-simvastatin combination with major adverse cardiovascular events in patients with or without diabetes: a retrospective propensity score-matched cohort study. <i>Scientific Reports</i> , 2018 , 8, 11925	4.9	5
173	Elevated N-acetyl- β -glucosaminidase, a urinary tubular damage marker, is a significant predictor of carotid artery atherosclerosis in type 1 diabetes, independent of albuminuria: A cross-sectional study. <i>Journal of Diabetes and Its Complications</i> , 2018 , 32, 777-783	3.2	4
172	Cardiovascular Outcome Trials of Sodium Glucose Cotransporter 2 Inhibitor and Its Possible Cardioprotective Mechanism. <i>Journal of Lipid and Atherosclerosis</i> , 2018 , 7, 21	3	
171	Association of non-alcoholic steatohepatitis with subclinical myocardial dysfunction in non-cirrhotic patients. <i>Journal of Hepatology</i> , 2018 , 68, 764-772	13.4	55
170	Relationship Between Circulating Netrin-1 Concentration, Impaired Fasting Glucose, and Newly Diagnosed Type 2 Diabetes. <i>Frontiers in Endocrinology</i> , 2018 , 9, 691	5.7	14
169	Diabetes Fact Sheet in Korea, 2016: An Appraisal of Current Status. <i>Diabetes and Metabolism Journal</i> , 2018 , 42, 415-424	5	64
168	Effects of Serum Albumin, Calcium Levels, Cancer Stage and Performance Status on Weight Loss in Parathyroid Hormone-Related Peptide Positive or Negative Patients with Cancer. <i>Endocrinology and Metabolism</i> , 2018 , 33, 97-104	3.5	0
167	Clinical parameters affecting dapagliflozin response in patients with type 2 diabetes. <i>Diabetes and Metabolism</i> , 2017 , 43, 191-194	5.4	9

166	High-dose sulphonylurea treatment in patients with renal impairment should be considered with caution. <i>Evidence-Based Medicine</i> , 2017 , 22, 72		
165	The renal tubular damage marker urinary N-acetyl- β -glucosaminidase may be more closely associated with early detection of atherosclerosis than the glomerular damage marker albuminuria in patients with type 2 diabetes. <i>Cardiovascular Diabetology</i> , 2017 , 16, 16	8.7	18
164	The Relationship between Increases in Morning Spot Urinary Glucose Excretion and Decreases in HbA in Patients with Type 2 Diabetes After Taking an SGLT2 Inhibitor: A Retrospective, Longitudinal Study. <i>Diabetes Therapy</i> , 2017 , 8, 601-609	3.6	3
163	Morning Spot Urine Glucose-to-Creatinine Ratios Predict Overnight Urinary Glucose Excretion in Patients With Type 2 Diabetes. <i>Annals of Laboratory Medicine</i> , 2017 , 37, 9-17	3.1	4
162	Association between Metformin Use and Risk of Lactic Acidosis or Elevated Lactate Concentration in Type 2 Diabetes. <i>Yonsei Medical Journal</i> , 2017 , 58, 312-318	3	11
161	Ezetimibe ameliorates steatohepatitis via AMP activated protein kinase-TFEB-mediated activation of autophagy and NLRP3 inflammasome inhibition. <i>Autophagy</i> , 2017 , 13, 1767-1781	10.2	97
160	Diabetes self-assessment score and the development of diabetes: A 10-year prospective study. <i>Medicine (United States)</i> , 2017 , 96, e7067	1.8	2
159	Pre-sarcopenia is associated with renal hyperfiltration independent of obesity or insulin resistance: Nationwide Surveys (KNHANES 2008-2011). <i>Medicine (United States)</i> , 2017 , 96, e7165	1.8	4
158	Anatomic fat depots and cardiovascular risk: a focus on the leg fat using nationwide surveys (KNHANES 2008-2011). <i>Cardiovascular Diabetology</i> , 2017 , 16, 54	8.7	15
157	Effect of statin on hepatocellular carcinoma in patients with type 2 diabetes: A nationwide nested case-control study. <i>International Journal of Cancer</i> , 2017 , 140, 798-806	7.5	42
156	Impact of diabetes mellitus and chronic liver disease on the incidence of dementia and all-cause mortality among patients with dementia. <i>Medicine (United States)</i> , 2017 , 96, e8753	1.8	8
155	Comparison between Atorvastatin and Rosuvastatin in Renal Function Decline among Patients with Diabetes. <i>Endocrinology and Metabolism</i> , 2017 , 32, 274-280	3.5	4
154	Exogenous administration of DLK1 ameliorates hepatic steatosis and regulates gluconeogenesis via activation of AMPK. <i>International Journal of Obesity</i> , 2016 , 40, 356-65	5.5	14
153	Paroxetine-induced Hypoglycemia in Type 2 Diabetic Patient. <i>The Ewha Medical Journal</i> , 2016 , 39, 14	0.1	0
152	Sarcopenia is associated with albuminuria independently of hypertension and diabetes: KNHANES 2008-2011. <i>Metabolism: Clinical and Experimental</i> , 2016 , 65, 1531-40	12.7	33
151	Association between dietary acid load and the risk of cardiovascular disease: nationwide surveys (KNHANES 2008-2011). <i>Cardiovascular Diabetology</i> , 2016 , 15, 122	8.7	38
150	Efficacy of different dipeptidyl peptidase-4 (DPP-4) inhibitors on metabolic parameters in patients with type 2 diabetes undergoing dialysis. <i>Medicine (United States)</i> , 2016 , 95, e4543	1.8	11
149	Urinary N-acetyl- β -glucosaminidase, an early marker of diabetic kidney disease, might reflect glucose excursion in patients with type 2 diabetes. <i>Medicine (United States)</i> , 2016 , 95, e4114	1.8	27

148	Sarcopenia is associated with significant liver fibrosis independently of obesity and insulin resistance in nonalcoholic fatty liver disease: Nationwide surveys (KNHANES 2008-2011). <i>Hepatology</i> , 2016 , 63, 776-86	11.2	175
147	Serum PTHrP Predicts Weight Loss in Cancer Patients Independent of Hypercalcemia, Inflammation, and Tumor Burden. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2016 , 101, 1207-14	5.6	21
146	Depot-Specific Changes in Fat Metabolism with Aging in a Type 2 Diabetic Animal Model. <i>PLoS ONE</i> , 2016 , 11, e0148141	3.7	8
145	Hypoglycemia and Health Costs. <i>Journal of Korean Diabetes</i> , 2016 , 17, 11	0.1	
144	A Case of Autoantibody-Positive Ketosis-Prone Diabetes Mellitus. <i>Journal of Korean Diabetes</i> , 2016 , 17, 60	0.1	1
143	Effects of Dipeptidyl Peptidase-4 Inhibitors on Hyperglycemia and Blood Cyclosporine Levels in Renal Transplant Patients with Diabetes: A Pilot Study. <i>Endocrinology and Metabolism</i> , 2016 , 31, 161-7	3.5	15
142	Characteristics Predictive for a Successful Switch from Insulin Analogue Therapy to Oral Hypoglycemic Agents in Patients with Type 2 Diabetes. <i>Yonsei Medical Journal</i> , 2016 , 57, 1395-403	3	
141	Metformin Restores Parkin-Mediated Mitophagy, Suppressed by Cytosolic p53. <i>International Journal of Molecular Sciences</i> , 2016 , 17,	6.3	52
140	Effects of Omega-3 Fatty Acid Supplementation on Diabetic Nephropathy Progression in Patients with Diabetes and Hypertriglyceridemia. <i>PLoS ONE</i> , 2016 , 11, e0154683	3.7	28
139	Economic Burden of Hypoglycemia in Patients with Type 2 Diabetes Mellitus from Korea. <i>PLoS ONE</i> , 2016 , 11, e0151282	3.7	10
138	Rosiglitazone Use and the Risk of Bladder Cancer in Patients With Type 2 Diabetes. <i>Medicine (United States)</i> , 2016 , 95, e2786	1.8	21
137	Comparison and Validation of 10 Equations Including a Novel Method for Estimation of LDL-cholesterol in a 168,212 Asian Population. <i>Medicine (United States)</i> , 2016 , 95, e3230	1.8	10
136	A genetic variant in GLP1R is associated with response to DPP-4 inhibitors in patients with type 2 diabetes. <i>Medicine (United States)</i> , 2016 , 95, e5155	1.8	22
135	Association between betatrophin/ANGPTL8 and non-alcoholic fatty liver disease: animal and human studies. <i>Scientific Reports</i> , 2016 , 6, 24013	4.9	54
134	A novel imaging platform for non-invasive screening of abnormal glucose tolerance. <i>Diabetes Research and Clinical Practice</i> , 2016 , 116, 83-5	7.4	6
133	Risk assessment and management of post-transplant diabetes mellitus. <i>Metabolism: Clinical and Experimental</i> , 2016 , 65, 1559-69	12.7	9
132	Chronic HMGCR/HMG-CoA reductase inhibitor treatment contributes to dysglycemia by upregulating hepatic gluconeogenesis through autophagy induction. <i>Autophagy</i> , 2015 , 11, 2089-2101	10.2	34
131	Undiagnosed diabetes is prevalent in younger adults and associated with a higher risk cardiometabolic profile compared to diagnosed diabetes. <i>American Heart Journal</i> , 2015 , 170, 760-769.e2	4.9	14

130	Increased expression of ATP-binding cassette transporter A1 (ABCA1) as a possible mechanism for the protective effect of cilostazol against hepatic steatosis. <i>Metabolism: Clinical and Experimental</i> , 2015 , 64, 1444-53	12.7	16
129	Sarcopaenia is associated with NAFLD independently of obesity and insulin resistance: Nationwide surveys (KNHANES 2008-2011). <i>Journal of Hepatology</i> , 2015 , 63, 486-93	13.4	186
128	Risk of diabetes in patients treated with HMG-CoA reductase inhibitors. <i>Metabolism: Clinical and Experimental</i> , 2015 , 64, 482-8	12.7	34
127	Association Between Heme Oxygenase-1 Promoter Polymorphisms and the Development of Albuminuria in Type 2 Diabetes: A Case-Control Study. <i>Medicine (United States)</i> , 2015 , 94, e1825	1.8	17
126	Visceral adiposity is associated with altered myocardial glucose uptake measured by (18)FDG-PET in 346 subjects with normal glucose tolerance, prediabetes, and type 2 diabetes. <i>Cardiovascular Diabetology</i> , 2015 , 14, 148	8.7	22
125	Use of a Diabetes Self-Assessment Score to Predict Nonalcoholic Fatty Liver Disease and Nonalcoholic Steatohepatitis. <i>Medicine (United States)</i> , 2015 , 94, e1103	1.8	6
124	Optimal candidates for the switch from glimepiride to sitagliptin to reduce hypoglycemia in patients with type 2 diabetes mellitus. <i>Endocrinology and Metabolism</i> , 2015 , 30, 84-91	3.5	11
123	1,5-anhydroglucitol as a useful marker for assessing short-term glycemic excursions in type 1 diabetes. <i>Diabetes and Metabolism Journal</i> , 2015 , 39, 164-70	5	19
122	Glycated Albumin Levels in Patients with Type 2 Diabetes Increase Relative to HbA1c with Time. <i>BioMed Research International</i> , 2015 , 2015, 576306	3	6
121	Glycated albumin and the risk of micro- and macrovascular complications in subjects with type 1 diabetes. <i>Cardiovascular Diabetology</i> , 2015 , 14, 53	8.7	13
120	Metformin alleviates hepatosteatosis by restoring SIRT1-mediated autophagy induction via an AMP-activated protein kinase-independent pathway. <i>Autophagy</i> , 2015 , 11, 46-59	10.2	179
119	Comparing kidney outcomes in type 2 diabetes treated with different sulphonylureas in real-life clinical practice. <i>Diabetes and Metabolism</i> , 2015 , 41, 208-15	5.4	12
118	Postprandial C-peptide to glucose ratio as a predictor of β -cell function and its usefulness for staged management of type 2 diabetes. <i>Journal of Diabetes Investigation</i> , 2014 , 5, 517-24	3.9	20
117	Tolerability, effectiveness and predictive parameters for the therapeutic usefulness of exenatide in obese, Korean patients with type 2 diabetes. <i>Journal of Diabetes Investigation</i> , 2014 , 5, 554-62	3.9	14
116	The hypolipidemic effect of cilostazol can be mediated by regulation of hepatic low-density lipoprotein receptor-related protein 1 (LRP1) expression. <i>Metabolism: Clinical and Experimental</i> , 2014 , 63, 112-9	12.7	15
115	HMG CoA reductase inhibitor treatment induces dysglycemia in renal allograft recipients. <i>Transplantation</i> , 2014 , 97, 419-25	1.8	9
114	Response: The Effect of DPP-4 Inhibitors on Metabolic Parameters in Patients with Type 2 Diabetes (Diabetes Metab J 2014;38:211-9). <i>Diabetes and Metabolism Journal</i> , 2014 , 38, 319-20	5	
113	The relationship between BMI and glycated albumin to glycated hemoglobin (GA/A1c) ratio according to glucose tolerance status. <i>PLoS ONE</i> , 2014 , 9, e89478	3.7	24

112	Non-laboratory-based self-assessment screening score for non-alcoholic fatty liver disease: development, validation and comparison with other scores. <i>PLoS ONE</i> , 2014 , 9, e107584	3.7	53
111	Inverse association between glycated albumin and insulin secretory function may explain higher levels of glycated albumin in subjects with longer duration of diabetes. <i>PLoS ONE</i> , 2014 , 9, e108772	3.7	8
110	The Effect of DPP-4 Inhibitors on Metabolic Parameters in Patients with Type 2 Diabetes. <i>Diabetes and Metabolism Journal</i> , 2014 , 38, 211-9	5	26
109	The glycated albumin to glycated hemoglobin ratio might not be associated with carotid atherosclerosis in patients with type 1 diabetes. <i>Diabetes and Metabolism Journal</i> , 2014 , 38, 456-63	5	10
108	Adipocytokines, metabolic syndrome, and exercise. <i>International Journal of Endocrinology</i> , 2014 , 2014, 597162	2.7	7
107	GPR30 mediates anorectic estrogen-induced STAT3 signaling in the hypothalamus. <i>Metabolism: Clinical and Experimental</i> , 2014 , 63, 1455-61	12.7	18
106	Statin therapy is associated with the development of new-onset diabetes after transplantation in liver recipients with high fasting plasma glucose levels. <i>Liver Transplantation</i> , 2014 , 20, 557-63	4.5	18
105	Screening for prediabetes using machine learning models. <i>Computational and Mathematical Methods in Medicine</i> , 2014 , 2014, 618976	2.8	37
104	Serum cholesterol concentration and prevalence, awareness, treatment, and control of high low-density lipoprotein cholesterol in the Korea National Health and Nutrition Examination Surveys 2008-2010: Beyond the Tip of the Iceberg. <i>Journal of the American Heart Association</i> , 2014 , 3, e000650	6	33
103	Anti-obesity effects of KR-66195, a synthetic DPP-IV inhibitor, in diet-induced obese mice and obese-diabetic ob/ob mice. <i>Metabolism: Clinical and Experimental</i> , 2014 , 63, 793-9	12.7	4
102	The effect of insulin resistance on postprandial triglycerides in Korean type 2 diabetic patients. <i>Acta Diabetologica</i> , 2014 , 51, 15-22	3.9	2
101	Effect of cilostazol on carotid intima-media thickness in type 2 diabetic patients without cardiovascular event. <i>Endocrine</i> , 2014 , 47, 138-45	4	7
100	Management of Diabetes in Organ Transplant Patients. <i>Journal of Korean Diabetes</i> , 2014 , 15, 134	0.1	1
99	Transcription factor Snail is a novel regulator of adipocyte differentiation via inhibiting the expression of peroxisome proliferator-activated receptor α <i>Cellular and Molecular Life Sciences</i> , 2013 , 70, 3959-71	10.3	37
98	Variants of the adiponectin gene and diabetic microvascular complications in patients with type 2 diabetes. <i>Metabolism: Clinical and Experimental</i> , 2013 , 62, 677-85	12.7	18
97	Gastrectomy for early gastric cancer is associated with decreased cardiovascular mortality in association with postsurgical metabolic changes. <i>Annals of Surgical Oncology</i> , 2013 , 20, 1250-7	3.1	20
96	Glycated albumin causes pancreatic β -cells dysfunction through autophagy dysfunction. <i>Endocrinology</i> , 2013 , 154, 2626-39	4.8	18
95	Rosiglitazone attenuates casein-induced hepatic endoplasmic reticulum stress in Sprague-Dawley rats: a novel model of endoplasmic reticulum stress. <i>Endocrine Journal</i> , 2013 , 60, 1231-40	2.9	7

94	A Case of Pheochromocytoma Associated with Diabetic Ketoacidosis and Infective Endocarditis. <i>Journal of Korean Diabetes</i> , 2013 , 14, 156	0.1	
93	Association between EPCs count and rate of coronary revascularization in asymptomatic type 2 diabetic patients. <i>Acta Diabetologica</i> , 2012 , 49, 413-20	3.9	5
92	Serum glycated albumin predicts the progression of carotid arterial atherosclerosis. <i>Atherosclerosis</i> , 2012 , 225, 450-5	3.1	48
91	Epicardial adipose tissue thickness is an indicator for coronary artery stenosis in asymptomatic type 2 diabetic patients: its assessment by cardiac magnetic resonance. <i>Cardiovascular Diabetology</i> , 2012 , 11, 83	8.7	40
90	The risk of bladder cancer in korean diabetic subjects treated with pioglitazone. <i>Diabetes and Metabolism Journal</i> , 2012 , 36, 371-8	5	31
89	Balsamic Vinegar Improves High Fat-Induced Beta Cell Dysfunction via Beta Cell ABCA1. <i>Diabetes and Metabolism Journal</i> , 2012 , 36, 275-9	5	10
88	Korean red ginseng (<i>Panax ginseng</i>) improves insulin sensitivity in high fat fed Sprague-Dawley rats. <i>Phytotherapy Research</i> , 2012 , 26, 142-7	6.7	39
87	Variants of the adiponectin and adiponectin receptor-1 genes and posttransplantation diabetes mellitus in renal allograft recipients. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2012 , 97, E129-35	5.6	18
86	Pentoxifylline attenuates methionine- and choline-deficient-diet-induced steatohepatitis by suppressing TNF- α expression and endoplasmic reticulum stress. <i>Experimental Diabetes Research</i> , 2012 , 2012, 762565		16
85	Upregulation of hepatic LRP1 by rosiglitazone: a possible novel mechanism of the beneficial effect of thiazolidinediones on atherogenic dyslipidemia. <i>Journal of Molecular Endocrinology</i> , 2012 , 49, 165-74	4.5	14
84	Higher morning to evening ratio in total dose of twice-daily biphasic insulin analog might be effective in achieving glucose control in patients with poorly controlled type 2 diabetes. <i>Diabetes Technology and Therapeutics</i> , 2012 , 14, 508-14	8.1	8
83	The effect of rosiglitazone on LRP1 expression and amyloid β uptake in human brain microvascular endothelial cells: a possible role of a low-dose thiazolidinedione for dementia treatment. <i>International Journal of Neuropsychopharmacology</i> , 2012 , 15, 135-42	5.8	28
82	A Case of Latent Autoimmune Diabetes in Adults Developed after Surgical Cure of Growth Hormone Secreting Pituitary Tumor. <i>Endocrinology and Metabolism</i> , 2012 , 27, 318	3.5	
81	Postprandial Triglyceride Is Associated with Fasting Triglyceride and HOMA-IR in Korean Subjects with Type 2 Diabetes. <i>Diabetes and Metabolism Journal</i> , 2011 , 35, 404-10	5	13
80	Associations between Fatness, Fitness, IGF and IMT among Obese Korean Male Adolescents. <i>Diabetes and Metabolism Journal</i> , 2011 , 35, 610-8	5	17
79	A Case of Gabapentin-induced Myoclonus in a Type 2 Diabetic Patient with End-stage Renal Disease. <i>Journal of Korean Diabetes</i> , 2011 , 12, 171	0.1	
78	Dietary oleate has beneficial effects on every step of non-alcoholic Fatty liver disease progression in a methionine- and choline-deficient diet-fed animal model. <i>Diabetes and Metabolism Journal</i> , 2011 , 35, 489-96	5	16
77	Lithospermic acid B protects β cells from cytokine-induced apoptosis by alleviating apoptotic pathways and activating anti-apoptotic pathways of Nrf2-HO-1 and Sirt1. <i>Toxicology and Applied Pharmacology</i> , 2011 , 252, 47-54	4.6	35

76	Fat redistribution preferentially reflects the anti-inflammatory benefits of pioglitazone treatment. <i>Metabolism: Clinical and Experimental</i> , 2011 , 60, 165-72	12.7	13
75	Up-regulation of hepatic low-density lipoprotein receptor-related protein 1: a possible novel mechanism of antiatherogenic activity of hydroxymethylglutaryl-coenzyme A reductase inhibitor Atorvastatin and hepatic LRP1 expression. <i>Metabolism: Clinical and Experimental</i> , 2011 , 60, 930-40	12.7	33
74	Leptin replacement improves postprandial glycemia and insulin sensitivity in human immunodeficiency virus-infected lipotrophic men treated with pioglitazone: a pilot study. <i>Metabolism: Clinical and Experimental</i> , 2011 , 60, 1045-9	12.7	32
73	Effects of leptin and adiponectin on pancreatic β cell function. <i>Metabolism: Clinical and Experimental</i> , 2011 , 60, 1664-72	12.7	98
72	Glycated albumin is a useful glycation index for monitoring fluctuating and poorly controlled type 2 diabetic patients. <i>Acta Diabetologica</i> , 2011 , 48, 167-72	3.9	64
71	Circulating vaspin and visfatin are not affected by acute or chronic energy deficiency or leptin administration in humans. <i>European Journal of Endocrinology</i> , 2011 , 164, 911-7	6.5	14
70	Effects of pioglitazone on subclinical atherosclerosis and insulin resistance in nondiabetic renal allograft recipients. <i>Nephrology Dialysis Transplantation</i> , 2010 , 25, 976-84	4.3	14
69	Protective effects of magnesium lithospermate B against diabetic atherosclerosis via Nrf2-ARE-NQO1 transcriptional pathway. <i>Atherosclerosis</i> , 2010 , 211, 69-76	3.1	49
68	Natural therapeutic magnesium lithospermate B potentially protects the endothelium from hyperglycaemia-induced dysfunction. <i>Cardiovascular Research</i> , 2010 , 87, 713-22	9.9	36
67	Dietary monounsaturated fatty acids but not saturated fatty acids preserve the insulin signaling pathway via IRS-1/PI3K in rat skeletal muscle. <i>Lipids</i> , 2010 , 45, 1109-16	1.6	31
66	Effect of 17-beta Estradiol on Adipocyte Lipin-1 Expression in OLETF Rat. <i>Endocrinology and Metabolism</i> , 2010 , 25, 199	3.5	1
65	The C609T variant of NQO1 is associated with carotid artery plaques in patients with type 2 diabetes. <i>Molecular Genetics and Metabolism</i> , 2009 , 97, 85-90	3.7	20
64	Korean red ginseng (<i>Panax ginseng</i>) improves insulin sensitivity and attenuates the development of diabetes in Otsuka Long-Evans Tokushima fatty rats. <i>Metabolism: Clinical and Experimental</i> , 2009 , 58, 1170-7	12.7	80
63	Retinol-binding protein-4 is associated with endothelial dysfunction in adults with newly diagnosed type 2 diabetes mellitus. <i>Atherosclerosis</i> , 2009 , 204, 23-5	3.1	37
62	Association of common type 2 diabetes risk gene variants and posttransplantation diabetes mellitus in renal allograft recipients in Korea. <i>Transplantation</i> , 2009 , 88, 693-8	1.8	32
61	A Case of Autoimmune Hypoglycemia in the Presence of Both Anti-insulin and Anti-insulin Receptor Antibodies. <i>Korean Clinical Diabetes</i> , 2009 , 10, 123		2
60	Staged diabetes management according to individual patient insulin resistance and beta-cell function ameliorates glycaemic control in type 2 diabetes mellitus. <i>Clinical Endocrinology</i> , 2008 , 69, 549-554	3.4	6
59	Lithospermic acid B ameliorates the development of diabetic nephropathy in OLETF rats. <i>European Journal of Pharmacology</i> , 2008 , 579, 418-25	5.3	26

58	Therapeutic effect of magnesium lithospermate B on neointimal formation after balloon-induced vascular injury. <i>European Journal of Pharmacology</i> , 2008 , 586, 226-33	5.3	36
57	Adiponectin gene polymorphism 45T>G is associated with carotid artery plaques in patients with type 2 diabetes mellitus. <i>Metabolism: Clinical and Experimental</i> , 2008 , 57, 274-9	12.7	19
56	LPIN1 genetic variation is associated with rosiglitazone response in type 2 diabetic patients. <i>Molecular Genetics and Metabolism</i> , 2008 , 95, 96-100	3.7	21
55	Rosiglitazone improves insulin sensitivity with increased serum leptin levels in patients with type 2 diabetes mellitus. <i>Diabetes Research and Clinical Practice</i> , 2008 , 81, 42-9	7.4	27
54	Plasma free fatty acid level patterns according to cardiovascular risk status in postmenopausal women. <i>Clinica Chimica Acta</i> , 2008 , 392, 11-6	6.2	20
53	A polymorphism in the zinc transporter gene SLC30A8 confers resistance against posttransplantation diabetes mellitus in renal allograft recipients. <i>Diabetes</i> , 2008 , 57, 1043-7	0.9	66
52	Rosiglitazone inhibits early stage of glucolipotoxicity-induced beta-cell apoptosis. <i>Hormone Research</i> , 2008 , 70, 165-73		18
51	A variant of the transcription factor 7-like 2 (TCF7L2) gene and the risk of posttransplantation diabetes mellitus in renal allograft recipients. <i>Diabetes Care</i> , 2008 , 31, 63-8	14.6	45
50	NFkappaB activation is associated with its O-GlcNAcylation state under hyperglycemic conditions. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008 , 105, 17345-50	11.5	177
49	Effects of Walking and Physical Activity on Glucose Regulation among Type 2 Diabetics. <i>Korean Diabetes Journal</i> , 2008 , 32, 60		5
48	Multiple endocrine neoplasia type 1 with multiple leiomyomas linked to a novel mutation in the MEN1 gene. <i>Yonsei Medical Journal</i> , 2008 , 49, 655-61	3	8
47	Visceral fat thickness predicts fatty liver in Koreans with type 2 diabetes mellitus. <i>Journal of Korean Medical Science</i> , 2008 , 23, 256-61	4.7	3
46	Echogenic carotid artery plaques are associated with vertebral fractures in postmenopausal women with low bone mass. <i>Calcified Tissue International</i> , 2008 , 82, 411-7	3.9	19
45	A new method of screening for diabetic neuropathy using laser Doppler and photoplethysmography. <i>Medical and Biological Engineering and Computing</i> , 2008 , 46, 61-7	3.1	17
44	Adenoviral vector-mediated glucagon-like peptide 1 gene therapy improves glucose homeostasis in Zucker diabetic fatty rats. <i>Journal of Gene Medicine</i> , 2008 , 10, 260-8	3.5	17
43	Association between polymorphisms in SLC30A8, HHEX, CDKN2A/B, IGF2BP2, FTO, WFS1, CDKAL1, KCNQ1 and type 2 diabetes in the Korean population. <i>Journal of Human Genetics</i> , 2008 , 53, 991-998	4.3	126
42	Protective Effects of Lithospermic Acid B on Diabetic Nephropathy in OLETF Rats Comparing with Amlodipine and Losartan. <i>Korean Diabetes Journal</i> , 2008 , 32, 10		
41	Rosiglitazone protects human neuroblastoma SH-SY5Y cells against MPP+ induced cytotoxicity via inhibition of mitochondrial dysfunction and ROS production. <i>Journal of the Neurological Sciences</i> , 2007 , 253, 53-60	3.2	91

40	Depot-specific regulation of perilipin by rosiglitazone in a diabetic animal model. <i>Metabolism: Clinical and Experimental</i> , 2007 , 56, 676-85	12.7	22
39	A Case of Turner's Syndrome with Transient Hypopituitarism. <i>Journal of Korean Endocrine Society</i> , 2007 , 22, 266		
38	In vivo Corneal Confocal Microscopy and Nerve Growth Factor in Diabetic Microvascular Complications. <i>The Journal of Korean Diabetes Association</i> , 2007 , 31, 351		
37	Design, synthesis, and discovery of stilbene derivatives based on lithospermic acid B as potent protein tyrosine phosphatase 1B inhibitors. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2007 , 17, 4481-6	2.9	21
36	The adaptation and relationship of FGF-23 to changes in mineral metabolism in Graves Disease. <i>Clinical Endocrinology</i> , 2007 , 66, 854-8	3.4	12
35	Risk factors associated with the onset and progression of posttransplantation diabetes in renal allograft recipients. <i>Diabetes Care</i> , 2007 , 30, 609-15	14.6	76
34	Riedel's thyroiditis in a patient with recurrent subacute thyroiditis: a case report and review of the literature. <i>Endocrine Journal</i> , 2007 , 54, 559-62	2.9	24
33	Effects of rosiglitazone and metformin on inflammatory markers and adipokines: decrease in interleukin-18 is an independent factor for the improvement of homeostasis model assessment-beta in type 2 diabetes mellitus. <i>Clinical Endocrinology</i> , 2007 , 66, 282-9	3.4	56
32	Activation of NF- κ B and AP-1 in Peripheral Blood Mononuclear Cells Isolated from Patients with Diabetic Nephropathy. <i>The Journal of Korean Diabetes Association</i> , 2007 , 31, 261		
31	Adiponectin protects human neuroblastoma SH-SY5Y cells against acetaldehyde-induced cytotoxicity. <i>Biochemical Pharmacology</i> , 2006 , 72, 616-23	6	27
30	Antidiabetic agents from medicinal plants. <i>Current Medicinal Chemistry</i> , 2006 , 13, 1203-18	4.3	329
29	Decrement of postprandial insulin secretion determines the progressive nature of type-2 diabetes. <i>European Journal of Endocrinology</i> , 2006 , 155, 615-22	6.5	23
28	Rosiglitazone relieves acute ethanol-induced hangover in Sprague-Dawley rats. <i>Alcohol and Alcoholism</i> , 2006 , 41, 231-5	3.5	9
27	The 11482G >A polymorphism in the perilipin gene is associated with weight gain with rosiglitazone treatment in type 2 diabetes. <i>Diabetes Care</i> , 2006 , 29, 1320-4	14.6	42
26	Effects of lifestyle modification on metabolic parameters and carotid intima-media thickness in patients with type 2 diabetes mellitus. <i>Metabolism: Clinical and Experimental</i> , 2006 , 55, 1053-9	12.7	88
25	Clinical and biochemical characteristics of nonobese type 2 diabetic patients with glutamic acid decarboxylase antibody in Korea. <i>Metabolism: Clinical and Experimental</i> , 2006 , 55, 1107-12	12.7	7
24	Rosiglitazone protects human neuroblastoma SH-SY5Y cells against acetaldehyde-induced cytotoxicity. <i>Biochemical and Biophysical Research Communications</i> , 2006 , 340, 221-7	3.4	37
23	Adiponectin protects human neuroblastoma SH-SY5Y cells against MPP ⁺ -induced cytotoxicity. <i>Biochemical and Biophysical Research Communications</i> , 2006 , 343, 564-70	3.4	38

22	Normative values of carotid artery intima-media thickness in healthy Korean adults and estimation of macrovascular diseases relative risk using this data in type 2 diabetes patients. <i>Diabetes Research and Clinical Practice</i> , 2006 , 72, 183-9	7.4	13
21	Haplotype combination of Calpain-10 gene polymorphism is associated with metabolic syndrome in type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2006 , 73, 268-75	7.4	15
20	The association of total and differential white blood cell count with metabolic syndrome in type 2 diabetic patients. <i>Diabetes Research and Clinical Practice</i> , 2006 , 73, 284-91	7.4	82
19	A Case of Autoimmune Hypoglycemia due to Insulin Antibody in Patient with End Stage Renal Disease. <i>Journal of Korean Endocrine Society</i> , 2006 , 21, 536		
18	Does birthweight predict adolescent adiponectin levels?. <i>Clinical Endocrinology</i> , 2006 , 64, 162-8	3.4	14
17	Effects of growth hormone on insulin resistance and atherosclerotic risk factors in obese type 2 diabetic patients with poor glycaemic control. <i>Clinical Endocrinology</i> , 2006 , 64, 444-9	3.4	17
16	The long-term effects of rosiglitazone on serum lipid concentrations and body weight. <i>Clinical Endocrinology</i> , 2006 , 65, 453-9	3.4	21
15	A novel 111/121 diplotype in the Calpain-10 gene is associated with type 2 diabetes. <i>Journal of Human Genetics</i> , 2006 , 51, 629-33	4.3	26
14	Contributing Factors to Different Natural Courses of Posttransplantation Diabetes Mellitus in Renal Allograft Recipients. <i>Journal of Korean Endocrine Society</i> , 2006 , 21, 373		2
13	The Relationship between the Leptin Concentration and the Small Dense Low Density Lipoprotein Cholesterol Concentration in Korean Type 2 Diabetic Patients. <i>Journal of Korean Endocrine Society</i> , 2006 , 21, 319		2
12	A Case of Patient with Opioid-Induced Adrenocortical Insufficiency and Hypogonadism. <i>Journal of Korean Endocrine Society</i> , 2006 , 21, 257		
11	A Case of Multiple Endocrine Neoplasia Type 1 with Papillary Thyroid Carcinoma. <i>Journal of Korean Endocrine Society</i> , 2006 , 21, 79		1
10	Limitation of the validity of the homeostasis model assessment as an index of insulin resistance in Korea. <i>Metabolism: Clinical and Experimental</i> , 2005 , 54, 206-11	12.7	101
9	Relationship of serum high sensitivity C-reactive protein to metabolic syndrome and microvascular complications in type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2005 , 69, 151-9	7.4	55
8	Effects of Pro12Ala polymorphism of peroxisome proliferator-activated receptor gamma2 gene on rosiglitazone response in type 2 diabetes. <i>Clinical Pharmacology and Therapeutics</i> , 2005 , 78, 202-8	6.1	94
7	Adiponectin Gene Polymorphism and Carotid Artery Intima-Media thickness in Type 2 Diabetes. <i>Journal of Korean Endocrine Society</i> , 2005 , 20, 29		
6	The influence of adiponectin gene polymorphism on the rosiglitazone response in patients with type 2 diabetes. <i>Diabetes Care</i> , 2005 , 28, 1139-44	14.6	88
5	A Case of Multiple Endocrine Neoplasia Type 1 with Mutation in MENIN Gene. <i>Journal of Korean Endocrine Society</i> , 2005 , 20, 71		6

4	Hypercalcemia in hepatic tuberculosis: a case report in Korea. <i>American Journal of Tropical Medicine and Hygiene</i> , 2005 , 72, 368-9	3.2	2
3	Serum high sensitivity C-reactive protein is associated with carotid intima-media thickness in type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2004 , 66 Suppl 1, S115-20	7.4	28
2	Hydrochloric Acid Induced Hyperkalemia. <i>Daehan Jeonhaejjil Daejsa Yeonjguhoeji</i> , 2004 , 2, 65		
1	Rhabdomyolysis and Disseminated Intravascular Coagulation Associated with Severe Exercise-induced Hypernatremic Dehydration. <i>Daehan Jeonhaejjil Daejsa Yeonjguhoeji</i> , 2004 , 2, 62		