## Yixun Yu

## List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/257573/publications.pdf Version: 2024-02-01



| #  | Article  | IF  | CITATIONS |
|----|--|-----|-----------|
| 1  | Mitochondrial citrate synthase plays important roles in anthocyanin synthesis in petunia. Plant<br>Science, 2021, 305, 110835.   | 3.6 | 18        |
| 2  | Phosphoproteome analysis reveals the involvement of protein dephosphorylation in ethylene-induced corolla senescence in petunia. BMC Plant Biology, 2021, 21, 512.   | 3.6 | 2         |
| 3  | Genomic identification and expression analysis of the BBX transcription factor gene family in Petunia<br>hybrida. Molecular Biology Reports, 2020, 47, 6027-6041.  | 2.3 | 12        |
| 4  | PaACL silencing accelerates flower senescence and changes the proteome to maintain metabolic homeostasis in Petunia hybrida. Journal of Experimental Botany, 2020, 71, 4858-4876.                            | 4.8 | 11        |
| 5  | The N <sup>1</sup> -Methyladenosine Methylome of Petunia mRNA. Plant Physiology, 2020, 183, 1710-1724.   | 4.8 | 31        |
| 6  | Genome-Wide Identification and Expression Profile Analysis of the NF-Y Transcription Factor Gene<br>Family in Petunia hybrida. Plants, 2020, 9, 336.   | 3.5 | 10        |
| 7  | Suppression of chorismate synthase, which is localized in chloroplasts and peroxisomes, results in abnormal flower development and anthocyanin reduction in petunia. Scientific Reports, 2020, 10, 10846.    | 3.3 | 10        |
| 8  | PhDHS Is Involved in Chloroplast Development in Petunia. Frontiers in Plant Science, 2019, 10, 284.  | 3.6 | 9         |
| 9  | The acyl-activating enzyme PhAAE13 is an alternative enzymatic source of precursors for anthocyanin biosynthesis in petunia flowers. Journal of Experimental Botany, 2017, 68, erw426.                       | 4.8 | 12        |
| 10 | PhCESA3 silencing inhibits elongation and stimulates radial expansion in petunia. Scientific Reports, 2017, 7, 41471.  | 3.3 | 13        |
| 11 | PhERF6, interacting with EOBI, negatively regulates fragrance biosynthesis in petunia flowers. New<br>Phytologist, 2017, 215, 1490-1502.   | 7.3 | 45        |
| 12 | Proteomes and Ubiquitylomes Analysis Reveals the Involvement of Ubiquitination in Protein<br>Degradation in Petunias. Plant Physiology, 2017, 173, 668-687.  | 4.8 | 80        |
| 13 | Molecular Characterization and Functional Analysis of Two Petunia PhEILs. Frontiers in Plant<br>Science, 2016, 7, 1606.  | 3.6 | 10        |
| 14 | Expression and functional analysis of PhEOL1 and PhEOL2 during flower senescence in petunia.<br>Functional Plant Biology, 2016, 43, 413.   | 2.1 | 6         |
| 15 | Functional characterization of PhGR and PhGRL1 during flower senescence in the petunia. Plant Cell Reports, 2015, 34, 1561-1568.   | 5.6 | 8         |
| 16 | PhGRL2 Protein, Interacting with PhACO1, Is Involved in Flower Senescence in the Petunia. Molecular Plant, 2014, 7, 1384-1387.   | 8.3 | 43        |
| 17 | The R2R3-MYB–Like Regulatory Factor EOBI, Acting Downstream of EOBII, Regulates Scent Production by Activating <i>ODO1</i> and Structural Scent-Related Genes in Petunia Â. Plant Cell, 2013, 24, 5089-5105. | 6.6 | 114       |
| 18 | Identification and expression analysis of ERF transcription factor genes in petunia during flower senescence and in response to hormone treatments. Journal of Experimental Botany, 2011, 62, 825-840.       | 4.8 | 85        |

Yixun Yu

| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 19 | Cloning and characterization of a DCEIN2 gene responsive to ethylene and sucrose in cut flower carnation. Plant Cell, Tissue and Organ Culture, 2011, 105, 447-455. | 2.3 | 16        |
| 20 | Relationship between Rh-RTH1 and ethylene receptor gene expression in response to ethylene in cut<br>rose. Plant Cell Reports, 2010, 29, 895-904.                   | 5.6 | 9         |