

Pierre J Richard

List of Publications by Citations

Source: <https://exaly.com/author-pdf/2583259/pierre-j-richard-publications-by-citations.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

110
papers

4,288
citations

39
h-index

60
g-index

111
ext. papers

4,760
ext. citations

2.9
avg, IF

5.15
L-index

#	Paper	IF	Citations
110	Isotopic Determination of Food Sources of <i>Crassostrea gigas</i> Along a Trophic Gradient in the Estuarine Bay of Marennes-Ol'fon. <i>Estuarine, Coastal and Shelf Science</i> , 1996 , 42, 347-360	2.9	230
109	Isotopic niches and trophic levels of myctophid fishes and their predators in the Southern Ocean. <i>Limnology and Oceanography</i> , 2010 , 55, 324-332	4.8	150
108	Stable isotopes reveal the trophic position and mesopelagic fish diet of female southern elephant seals breeding on the Kerguelen Islands. <i>Marine Ecology - Progress Series</i> , 2008 , 370, 239-247	2.6	145
107	Seabird satellite tracking validates the use of latitudinal isoscapes to depict predators' foraging areas in the Southern Ocean. <i>Rapid Communications in Mass Spectrometry</i> , 2010 , 24, 3456-60	2.2	141
106	Whisker isotopic signature depicts migration patterns and multi-year intra- and inter-individual foraging strategies in fur seals. <i>Biology Letters</i> , 2009 , 5, 830-2	3.6	125
105	Chemical and Isotopic Composition of the Organic Matter Sources in the Gulf of Gdansk (Southern Baltic Sea). <i>Estuarine, Coastal and Shelf Science</i> , 2000 , 51, 585-598	2.9	124
104	Resource partitioning within a tropical seabird community: new information from stable isotopes. <i>Marine Ecology - Progress Series</i> , 2008 , 366, 281-291	2.6	113
103	Determination of food sources for benthic invertebrates in a salt marsh (Aiguillon Bay, France) by carbon and nitrogen stable isotopes: importance of locally produced sources. <i>Marine Ecology - Progress Series</i> , 1999 , 187, 301-307	2.6	109
102	Food sources of an infaunal suspension-feeding bivalve <i>Cerastoderma edule</i> in a muddy sandflat of Marennes-Ol'fon Bay, as determined by analyses of carbon and nitrogen stable isotopes. <i>Marine Ecology - Progress Series</i> , 1999 , 187, 147-158	2.6	93
101	Quantifying the short-term temperature effect on light-saturated photosynthesis of intertidal microphytobenthos. <i>Marine Ecology - Progress Series</i> , 1996 , 134, 309-313	2.6	85
100	Effect of starvation on RNA, DNA and protein content of laboratory-reared larvae and juveniles of <i>Solea solea</i> . <i>Marine Ecology - Progress Series</i> , 1991 , 72, 69-77	2.6	79
99	Use of stable isotopes to quantify seasonal changes of trophic niche and levels of population and individual specialisation in seabirds. <i>Marine Ecology - Progress Series</i> , 2010 , 401, 269-277	2.6	78
98	Food source of intertidal nematodes in the Bay of Marennes-Ol'fon (France), as determined by dual stable isotope analysis. <i>Marine Ecology - Progress Series</i> , 1996 , 142, 303-309	2.6	78
97	Temporal variation of delta13C in particulate organic matter and oyster <i>Crassostrea gigas</i> in Marennes-Ol'fon Bay (France): effect of freshwater inflow. <i>Marine Ecology - Progress Series</i> , 1997 , 147, 105-115	2.6	76
96	The effect of geomorphological structures on potential biostabilisation by microphytobenthos on intertidal mudflats. <i>Continental Shelf Research</i> , 2000 , 20, 1243-1256	2.4	73
95	Dynamics of spatial patterns of microphytobenthic biomass: inferences from a geostatistical analysis of two comprehensive surveys in Marennes-Ol'fon Bay (France). <i>Marine Ecology - Progress Series</i> , 1998 , 166, 131-141	2.6	71
94	Long-term species, sexual and individual variations in foraging strategies of fur seals revealed by stable isotopes in whiskers. <i>PLoS ONE</i> , 2012 , 7, e32916	3.7	69

93	Trophic importance of diatoms in an intertidal <i>Zostera noltii</i> seagrass bed: Evidence from stable isotope and fatty acid analyses. <i>Estuarine, Coastal and Shelf Science</i> , 2011 , 92, 140-153	2.9	67
92	Stable isotopes document winter trophic ecology and maternal investment of adult female southern elephant seals (<i>Mirounga leonina</i>) breeding at the Kerguelen Islands. <i>Marine Biology</i> , 2008 , 155, 413-420	2.5	67
91	Short- and long-term consistency in the foraging niche of wandering albatrosses. <i>Marine Biology</i> , 2012 , 159, 1581-1591	2.5	66
90	SEASONAL EFFECT ON THE RELATIONSHIP BETWEEN THE PHOTOSYNTHETIC CAPACITY OF INTERTIDAL MICROPHYTOBENTHOS AND TEMPERATURE ¹ . <i>Journal of Phycology</i> , 1997 , 33, 723-728	3	64
89	Using carbon and nitrogen isotopic values of body feathers to infer inter- and intra-individual variations of seabird feeding ecology during moult. <i>Marine Biology</i> , 2009 , 156, 1233-1240	2.5	54
88	Microstructure of microphytobenthic biofilm and its spatio-temporal dynamics in an intertidal mudflat (Aiguillon Bay, France). <i>Marine Ecology - Progress Series</i> , 2004 , 282, 33-44	2.6	54
87	Tissue, ontogenic and sex-related differences in $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of the oceanic squid <i>Todarodes filippovae</i> (Cephalopoda: Ommastrephidae). <i>Marine Biology</i> , 2009 , 156, 699-708	2.5	52
86	Distribution and sources of organic matter in surface sediments of Bohai Sea near the Yellow River Estuary, China. <i>Estuarine, Coastal and Shelf Science</i> , 2015 , 165, 128-136	2.9	51
85	Ecological niche segregation within a community of sympatric dolphins around a tropical island. <i>Marine Ecology - Progress Series</i> , 2011 , 433, 273-288	2.6	51
84	Stable isotopes document the trophic structure of a deep-sea cephalopod assemblage including giant octopod and giant squid. <i>Biology Letters</i> , 2009 , 5, 364-7	3.6	50
83	$\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ in the Mondego estuary food web: seasonal variation in producers and consumers. <i>Marine Environmental Research</i> , 2009 , 67, 109-16	3.3	50
82	Spatio-temporal structure of the epipelagic diatom assemblage from an intertidal mudflat in Marennes-Olignon Bay, France. <i>Estuarine, Coastal and Shelf Science</i> , 2005 , 64, 385-394	2.9	49
81	Methodology of light response curves: application of chlorophyll fluorescence to microphytobenthic biofilms. <i>Marine Biology</i> , 2007 , 153, 91-101	2.5	48
80	Seabird year-round and historical feeding ecology: blood and feather $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values document foraging plasticity of small sympatric petrels. <i>Marine Ecology - Progress Series</i> , 2014 , 505, 267-280	2.6	48
79	Measurement of ingestion rate of <i>Hydrobia ulvae</i> (Pennant) on intertidal epipelagic microalgae: the effect of mud snail density. <i>Journal of Experimental Marine Biology and Ecology</i> , 2000 , 255, 247-260	2.1	46
78	Seasonal variation in consumption of benthic bacteria by meio- and macrofauna in an intertidal mudflat. <i>Limnology and Oceanography</i> , 2009 , 54, 1048-1059	4.8	45
77	Bacterivory in the common foraminifer <i>Ammonia tepida</i> : Isotope tracer experiment and the controlling factors. <i>Journal of Experimental Marine Biology and Ecology</i> , 2008 , 359, 55-61	2.1	45
76	Dynamics of particulate organic matter composition in coastal systems: A spatio-temporal study at multi-systems scale. <i>Progress in Oceanography</i> , 2017 , 156, 221-239	3.8	43

75	Eutrophication and trophic structure in response to the presence of the eelgrass <i>Zostera noltii</i> . <i>Marine Biology</i> , 2009 , 156, 2107-2120	2.5	43
74	Sequential isotopic signature along gladius highlights contrasted individual foraging strategies of jumbo squid (<i>Dosidicus gigas</i>). <i>PLoS ONE</i> , 2011 , 6, e22194	3.7	42
73	Food sources used by sediment meiofauna in an intertidal <i>Zostera noltii</i> seagrass bed: a seasonal stable isotope study. <i>Marine Biology</i> , 2012 , 159, 1537-1550	2.5	40
72	Effects of Lipid Extraction on $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ Values in Seabird Muscle, Liver and Feathers. <i>Waterbirds</i> , 2008 , 31, 169-178	0.5	40
71	Utilization of estuarine organic matter during growth and migration by juvenile brown shrimp <i>Penaeus aztecus</i> in a South Texas estuary. <i>Marine Ecology - Progress Series</i> , 2000 , 199, 205-216	2.6	37
70	A preliminary study of habitat and resource partitioning among co-occurring tropical dolphins around Mayotte, southwest Indian Ocean. <i>Estuarine, Coastal and Shelf Science</i> , 2009 , 84, 367-374	2.9	36
69	Archaeal Methane Cycling Communities Associated with Gassy Subsurface Sediments of Marennes-Olfon Bay (France). <i>Geomicrobiology Journal</i> , 2009 , 26, 31-43	2.5	35
68	Excretion of nitrogenous products by <i>Penaeus japonicus</i> bate in relation to environmental osmotic conditions. <i>Comparative Biochemistry and Physiology A, Comparative Physiology</i> , 1982 , 72, 673-678		35
67	Trophic ecology of mullets during their spring migration in a European saltmarsh: A stable isotope study. <i>Estuarine, Coastal and Shelf Science</i> , 2011 , 91, 502-510	2.9	34
66	Relative contribution of natural productivity and compound feed to tissue growth in blue shrimp (<i>Litopenaeus stylirostris</i>) reared in biofloc: Assessment by C and N stable isotope ratios and effect on key digestive enzymes. <i>Aquaculture</i> , 2015 , 448, 288-297	4.4	33
65	Ingestion rate of the deposit-feeder <i>Hydrobia ulvae</i> (Gastropoda) on epipellic diatoms: effect of cell size and algal biomass. <i>Journal of Experimental Marine Biology and Ecology</i> , 2005 , 317, 1-12	2.1	33
64	Stable isotope analysis of food source and trophic position of intertidal nematodes and copepods. <i>Marine Ecology - Progress Series</i> , 2008 , 359, 145-150	2.6	32
63	Are epiphytes a significant component of intertidal <i>Zostera noltii</i> beds?. <i>Aquatic Botany</i> , 2009 , 91, 82-90	1.8	31
62	Age, sex, and breeding status shape a complex foraging pattern in an extremely long-lived seabird. <i>Ecology</i> , 2014 , 95, 2324-33	4.6	30
61	A comprehensive isotopic investigation of habitat preferences in nonbreeding albatrosses from the Southern Ocean. <i>Ecography</i> , 2013 , 36, 277-286	6.5	30
60	Comparative foraging ecology and ecological niche of a superabundant tropical seabird: the sooty tern <i>Sterna fuscata</i> in the southwest Indian Ocean. <i>Marine Biology</i> , 2008 , 155, 505-520	2.5	30
59	Importance of Marine-Derived Nutrients Supplied by Planktivorous Seabirds to High Arctic Tundra Plant Communities. <i>PLoS ONE</i> , 2016 , 11, e0154950	3.7	30
58	Bacterivory of a mudflat nematode community under different environmental conditions. <i>Marine Biology</i> , 2008 , 154, 671-682	2.5	29

57	A geochemical record of environmental changes in sediments from Sishili Bay, northern Yellow Sea, China: anthropogenic influence on organic matter sources and composition over the last 100 years. <i>Marine Pollution Bulletin</i> , 2013 , 77, 227-36	6.7	28
56	The use of stable isotope analyses from skin biopsy samples to assess trophic relationships of sympatric delphinids off Moorea (French Polynesia). <i>Journal of Experimental Marine Biology and Ecology</i> , 2010 , 395, 48-54	2.1	28
55	Bacterivory by benthic organisms in sediment: Quantification using ^{15}N -enriched bacteria. <i>Journal of Experimental Marine Biology and Ecology</i> , 2008 , 355, 18-26	2.1	28
54	Identifying carbon sources and trophic position of coral reef fishes using diet and stable isotope (^{15}N and ^{13}C) analyses in two contrasted bays in Moorea, French Polynesia. <i>Coral Reefs</i> , 2013 , 32, 1091-1102	4.2	27
53	CHARACTERIZING AND QUANTIFYING PHOTOINHIBITION IN INTERTIDAL MICROPHYTOBENTHOS1. <i>Journal of Phycology</i> , 2004 , 40, 692-696	3	27
52	Trophic resource use by macrozoobenthic primary consumers within a semi-enclosed coastal ecosystem: Stable isotope and fatty acid assessment. <i>Journal of Sea Research</i> , 2014 , 88, 87-99	1.9	26
51	Influence of environment factors on bacterial ingestion rate of the deposit-feeder <i>Hydrobia ulvae</i> and comparison with meiofauna. <i>Journal of Sea Research</i> , 2008 , 60, 151-156	1.9	25
50	Benthic contribution to pelagic microalgal communities in two semi-enclosed, European-type littoral ecosystems (Marennes-Olfon Bay and Aiguillon Bay, France). <i>Journal of Sea Research</i> , 2004 , 52, 241-258	1.9	25
49	Trophic structure in the northern Humboldt Current system: new perspectives from stable isotope analysis. <i>Marine Biology</i> , 2017 , 164, 1	2.5	24
48	How does the resuspension of the biofilm alter the functioning of the benthos-pelagos coupled food web of a bare mudflat in Marennes-Olfon Bay (NE Atlantic)? <i>Journal of Sea Research</i> , 2014 , 92, 144-157	1.9	24
47	Key features of intertidal food webs that support migratory shorebirds. <i>PLoS ONE</i> , 2013 , 8, e76739	3.7	24
46	OSmother where wert thou? Maternal strategies in the southern elephant seal: a stable isotope investigation. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2012 , 279, 2681-90	4.4	24
45	Seasonal Variations in Suspended Particulate Matter in the Marennes-Olfon Bay, France, using Lipids as Biomarkers. <i>Estuarine, Coastal and Shelf Science</i> , 1996 , 43, 335-357	2.9	24
44	Incorporation of diet information derived from Bayesian stable isotope mixing models into mass-balanced marine ecosystem models: A case study from the Marennes-Olfon Estuary, France. <i>Ecological Modelling</i> , 2013 , 267, 127-137	3	23
43	Macroalgae ^{15}N values in well-mixed estuaries: Indicator of anthropogenic nitrogen input or macroalgae metabolism?. <i>Estuarine, Coastal and Shelf Science</i> , 2013 , 119, 126-138	2.9	23
42	Feeding ecology of striped dolphins, <i>Stenella coeruleoalba</i> , in the north-western Mediterranean Sea based on stable isotope analyses. <i>Journal of the Marine Biological Association of the United Kingdom</i> , 2012 , 92, 1677-1687	1.1	23
41	Is benthic food web structure related to diversity of marine macrobenthic communities?. <i>Estuarine, Coastal and Shelf Science</i> , 2012 , 108, 76-86	2.9	23
40	Spatial Variability of Stable Isotope Ratios in Oysters (<i>Crassostrea gigas</i>) and Primary Producers Along an Estuarine Gradient (Bay of Brest, France). <i>Estuaries and Coasts</i> , 2013 , 36, 808-819	2.8	21

39	Stable isotope patterns in micronekton from the Mozambique Channel. <i>Deep-Sea Research Part II: Topical Studies in Oceanography</i> , 2014 , 100, 153-163	2.3	21
38	Quantifying the trophic base for benthic secondary production in the Nakdong River estuary of Korea using stable C and N isotopes. <i>Journal of Experimental Marine Biology and Ecology</i> , 2009 , 382, 18-26 ^{2,1}	2.1	21
37	Photo-regulation in microphytobenthos from intertidal mudflats and non-tidal coastal shallows. <i>Estuarine, Coastal and Shelf Science</i> , 2015 , 152, 153-161	2.9	20
36	Influence de la temperature sur la croissance et la mue de Palaemon serratus en fonction de leur taille. <i>Aquaculture</i> , 1978 , 14, 13-22	4.4	20
35	Feeding strategies and resource partitioning ¶mitigate the effects of oligotrophy for marine cave mysids. <i>Marine Ecology - Progress Series</i> , 2011 , 440, 163-176	2.6	20
34	Hg Stable Isotope Time Trend in Ringed Seals Registers Decreasing Sea Ice Cover in the Alaskan Arctic. <i>Environmental Science & Technology</i> , 2015 , 49, 8977-85	10.3	19
33	Spatial variations in dietary organic matter sources modulate the size and condition of fish juveniles in temperate lagoon nursery sites. <i>Estuarine, Coastal and Shelf Science</i> , 2015 , 152, 78-90	2.9	19
32	Stable isotopes document inter- and intra-specific variation in feeding ecology of nine large southern Procellariiformes. <i>Marine Ecology - Progress Series</i> , 2013 , 490, 255-266	2.6	19
31	Organic matter exploitation in a highly turbid environment: Planktonic food web in the Charente estuary, France. <i>Estuarine, Coastal and Shelf Science</i> , 2012 , 98, 126-137	2.9	18
30	Shift in foraging grounds and diet broadening during ontogeny in southern elephant seals from Kerguelen Islands. <i>Marine Biology</i> , 2013 , 160, 977-986	2.5	18
29	NH ₄ ⁺ turnover in intertidal sediments of Marennes-Olŕon Bay (France): effect of sediment temperature. <i>Oceanologica Acta: European Journal of Oceanology - Revue Europeene De Oceanologie</i> , 2000 , 23, 575-584		18
28	Selection of effective macroalgal species and tracing nitrogen sources on the different part of Yantai coast, China indicated by macroalgal ¶(15)N values. <i>Science of the Total Environment</i> , 2016 , 542, 306-14	10.2	16
27	The influence of long emersion on biota, ammonium fluxes and nitrification in intertidal sediments of Marennes-Olŕon Bay, France. <i>Marine Environmental Research</i> , 2002 , 53, 381-402	3.3	15
26	Trophic structure of the macrobenthic community of Hornsund, Spitsbergen, based on the determination of stable carbon and nitrogen isotopic signatures. <i>Polar Biology</i> , 2014 , 37, 1247-1260	2	13
25	Differences in spatial structures between juveniles and adults of the gastropod <i>Hydrobia ulvae</i> on an intertidal mudflat (Marennes-Olŕon Bay, France) potentially affect estimates of local demographic processes. <i>Journal of Sea Research</i> , 2004 , 51, 63-68	1.9	13
24	Variations circadiennes des acides amin¶ libres du muscle de <i>Penaeus kerathurus</i> . <i>Biochemical Systematics and Ecology</i> , 1979 , 7, 65-67	1.4	13
23	Distribution of adsorbed ammonium pools in two intertidal sedimentary structures, Marennes-Olŕon Bay, France. <i>Marine Ecology - Progress Series</i> , 1999 , 182, 29-35	2.6	13
22	Carbon Stable Isotope Analysis of Methylmercury Toxin in Biological Materials by Gas Chromatography Isotope Ratio Mass Spectrometry. <i>Analytical Chemistry</i> , 2015 , 87, 11732-8	7.8	11

21	Tolérance aux températures extrêmes de <i>Palaemon serratus</i> (Pennant): Influence de la taille et de l'acclimatation. <i>Journal of Experimental Marine Biology and Ecology</i> , 1978 , 35, 137-146	2.1	11
20	Spatio-temporal differentiation in the population structure of <i>Hydrobia ulvae</i> on an intertidal mudflat (Marennes-Olifon Bay, France). <i>Journal of the Marine Biological Association of the United Kingdom</i> , 2002 , 82, 605-614	1.1	10
19	Transfer of ornithogenic influence through different trophic levels of the Arctic terrestrial ecosystem of Bjørnøya (Bear Island), Svalbard. <i>Soil Biology and Biochemistry</i> , 2017 , 115, 475-489	7.5	9
18	Does the size of the microphytobenthic biofilm on intertidal mudflats depend on the available photosynthetic biomass?. <i>Marine Ecology - Progress Series</i> , 2005 , 298, 95-100	2.6	9
17	Stable carbon isotope ratios differentiate autotrophs supporting animal diversity in Lake Baikal. <i>Comptes Rendus De L'Académie Des Sciences Série 3, Sciences De La Vie</i> , 1998 , 321, 509-516		8
16	Influence d'une microsporidiose sur les acides aminés libres de <i>Carcinus mediterraneus</i> Czerniavsky, 1884 soumis à diverses salinités et à des valeurs extrêmes de température. <i>Journal of Experimental Marine Biology and Ecology</i> , 1980 , 43, 207-220	2.1	8
15	Fluff deposition on intertidal sediments: effects on benthic biota, ammonium fluxes and nitrification rates. <i>Biogeochemistry</i> , 2002 , 61, 115-133	3.8	7
14	Cd transfer in the deposit-feeder Prosobranch <i>Hydrobia ulvae</i> (Pennant) from benthic diatoms: the kinetics of rapid Cd assimilation and efflux. <i>Journal of Experimental Marine Biology and Ecology</i> , 2005 , 317, 159-174	2.1	7
13	Anomalies de calcification chez la palourde japonaise <i>Ruditapes philippinarum</i> : Caractérisation et comparaison des compositions en acides aminés de différentes parties de la coquille analysés par HPLC. <i>Aquaculture</i> , 1989 , 81, 169-183	4.4	7
12	Long- and short-term photoacclimation in epipsammon from non-tidal coastal shallows compared to epipelon from intertidal mudflat. <i>Journal of Sea Research</i> , 2018 , 136, 1-9	1.9	6
11	Trophic shift in young-of-the-year Mugilidae during salt-marsh colonization. <i>Journal of Fish Biology</i> , 2013 , 82, 1297-307	1.9	6
10	Influence de la température sur les acides aminés libres de <i>Palaemon serratus</i> et <i>P. squilla</i> . <i>Biochemical Systematics and Ecology</i> , 1977 , 5, 297-300	1.4	6
9	Non-photochemical quenching in epipsammic and epipellic microalgal assemblages from two marine ecosystems. <i>Continental Shelf Research</i> , 2017 , 136, 74-82	2.4	4
8	Acides aminés de microsporidies parasites de crustacés décapodes marins. <i>Biochemical Systematics and Ecology</i> , 1980 , 8, 205-209	1.4	4
7	Le métabolisme amino-acide de <i>Palaemon serratus</i> : Variations des acides aminés libres du muscle et de l'hépatopancreas au cours du cycle de mue. <i>Comparative Biochemistry and Physiology A, Comparative Physiology</i> , 1980 , 67, 553-560		4
6	Evolution de quelques Enzymes Digestives chez le Jeune <i>Mugil capito</i> - Diverses Etapes de sa Croissance. <i>Biochemical Systematics and Ecology</i> , 1982 , 10, 185-189	1.4	4
5	The effects of temperature and salinity on phosphate levels in two euryhaline crustacean species. <i>Journal of Sea Research</i> , 1982 , 15, 284-292		4
4	Analyzing biases of nitrogen contents and δ ¹⁵ N values arising from acidified marine sediments with different CaCO ₃ concentrations. <i>Acta Oceanologica Sinica</i> , 2018 , 37, 1-5	1	3

3	Application of trichloroacetic acid (TCA) to extraction of soft body for the determination of tissue Cd, Cu, Pb and Zn in the prosobranch <i>Hydrobia ulvae</i> (Pennant). <i>Marine Pollution Bulletin</i> , 2003 , 46, 1326-33	6.7	3
2	Nitrification rates related to sedimentary structures in an Atlantic intertidal mudflat, Marennes-Olfon Bay, France. <i>Marine Ecology - Progress Series</i> , 1999 , 191, 33-41	2.6	3
1	Morphology and distribution of a little known but widespread diatom (Bacillariophyceae), <i>Navicula spartinetensis</i> Sullivan et Reimer. <i>Diatom Research</i> , 2012 , 27, 43-51	0.9	2