

Alison R Yung

List of Publications by Citations

Source: <https://exaly.com/author-pdf/2587055/alison-r-yung-publications-by-citations.pdf>

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

399
papers

30,840
citations

85
h-index

166
g-index

467
ext. papers

35,109
ext. citations

4.9
avg, IF

7.08
L-index

#	Paper	IF	Citations
399	Mapping the onset of psychosis: the Comprehensive Assessment of At-Risk Mental States. <i>Australian and New Zealand Journal of Psychiatry</i> , 2005 , 39, 964-71	2.6	1307
398	Neuroanatomical abnormalities before and after onset of psychosis: a cross-sectional and longitudinal MRI comparison. <i>Lancet, The</i> , 2003 , 361, 281-8	4.0	1044
397	Predicting psychosis: meta-analysis of transition outcomes in individuals at high clinical risk. <i>Archives of General Psychiatry</i> , 2012 , 69, 220-9		983
396	The psychosis high-risk state: a comprehensive state-of-the-art review. <i>JAMA Psychiatry</i> , 2013 , 70, 107-20	4.5	965
395	The prodromal phase of first-episode psychosis: past and current conceptualizations. <i>Schizophrenia Bulletin</i> , 1996 , 22, 353-70	1.3	833
394	Psychosis prediction: 12-month follow up of a high-risk ("prodromal") group. <i>Schizophrenia Research</i> , 2003 , 60, 21-32	3.6	805
393	Randomized controlled trial of interventions designed to reduce the risk of progression to first-episode psychosis in a clinical sample with subthreshold symptoms. <i>Archives of General Psychiatry</i> , 2002 , 59, 921-8		757
392	Risk factors for psychosis in an ultra high-risk group: psychopathology and clinical features. <i>Schizophrenia Research</i> , 2004 , 67, 131-42	3.6	597
391	Prediction of psychosis: A step towards indicated prevention of schizophrenia. <i>British Journal of Psychiatry</i> , 1998 , 172, 14-20	5.4	580
390	Monitoring and care of young people at incipient risk of psychosis. <i>Schizophrenia Bulletin</i> , 1996 , 22, 283-303	3.3	559
389	Clinical staging of psychiatric disorders: a heuristic framework for choosing earlier, safer and more effective interventions. <i>Australian and New Zealand Journal of Psychiatry</i> , 2006 , 40, 616-22	2.6	548
388	Hippocampal and amygdala volumes according to psychosis stage and diagnosis: a magnetic resonance imaging study of chronic schizophrenia, first-episode psychosis, and ultra-high-risk individuals. <i>Archives of General Psychiatry</i> , 2006 , 63, 139-49		496
387	Cognitive functioning in prodromal psychosis: a meta-analysis. <i>Archives of General Psychiatry</i> , 2012 , 69, 562-71		454
386	The Lancet Psychiatry Commission: a blueprint for protecting physical health in people with mental illness. <i>Lancet Psychiatry</i> , 2019 , 6, 675-712	23.3	411
385	Structural brain imaging evidence for multiple pathological processes at different stages of brain development in schizophrenia. <i>Schizophrenia Bulletin</i> , 2005 , 31, 672-96	1.3	409
384	Comorbid depressive and anxiety disorders in 509 individuals with an at-risk mental state: impact on psychopathology and transition to psychosis. <i>Schizophrenia Bulletin</i> , 2014 , 40, 120-31	1.3	402
383	The initial prodrome in psychosis: descriptive and qualitative aspects. <i>Australian and New Zealand Journal of Psychiatry</i> , 1996 , 30, 587-99	2.6	337

382	Declining transition rate in ultra high risk (prodromal) services: dilution or reduction of risk?. <i>Schizophrenia Bulletin</i> , 2007 , 33, 673-81	1.3	332
381	A systematic review and meta-analysis of exercise interventions in schizophrenia patients. <i>Psychological Medicine</i> , 2015 , 45, 1343-61	6.9	324
380	Long-term follow-up of a group at ultra high risk ("prodromal") for psychosis: the PACE 400 study. <i>JAMA Psychiatry</i> , 2013 , 70, 793-802	14.5	323
379	Early intervention in psychosis: concepts, evidence and future directions. <i>World Psychiatry</i> , 2008 , 7, 148-54	14.4	318
378	Memory impairments identified in people at ultra-high risk for psychosis who later develop first-episode psychosis. <i>American Journal of Psychiatry</i> , 2005 , 162, 71-8	11.9	317
377	Validation of "prodromal" criteria to detect individuals at ultra high risk of psychosis: 2 year follow-up. <i>Schizophrenia Research</i> , 2008 , 105, 10-7	3.6	287
376	Heterogeneity of Psychosis Risk Within Individuals at Clinical High Risk: A Meta-analytical Stratification. <i>JAMA Psychiatry</i> , 2016 , 73, 113-20	14.5	270
375	Preventing a first episode of psychosis: meta-analysis of randomized controlled prevention trials of 12 month and longer-term follow-ups. <i>Schizophrenia Research</i> , 2013 , 149, 56-62	3.6	270
374	Progressive gray matter reduction of the superior temporal gyrus during transition to psychosis. <i>Archives of General Psychiatry</i> , 2009 , 66, 366-76		266
373	Psychotic-like experiences in a community sample of adolescents: implications for the continuum model of psychosis and prediction of schizophrenia. <i>Australian and New Zealand Journal of Psychiatry</i> , 2009 , 43, 118-28	2.6	235
372	Progressive brain structural changes mapped as psychosis develops in 'at risk' individuals. <i>Schizophrenia Research</i> , 2009 , 108, 85-92	3.6	231
371	Motivating factors and barriers towards exercise in severe mental illness: a systematic review and meta-analysis. <i>Psychological Medicine</i> , 2016 , 46, 2869-2881	6.9	221
370	Intervention in individuals at ultra-high risk for psychosis: a review and future directions. <i>Journal of Clinical Psychiatry</i> , 2009 , 70, 1206-12	4.6	219
369	How much physical activity do people with schizophrenia engage in? A systematic review, comparative meta-analysis and meta-regression. <i>Schizophrenia Research</i> , 2016 , 176, 431-440	3.6	214
368	Mobile Phone Ownership and Endorsement of "mHealth" Among People With Psychosis: A Meta-analysis of Cross-sectional Studies. <i>Schizophrenia Bulletin</i> , 2016 , 42, 448-55	1.3	213
367	The "close-in" or ultra high-risk model: a safe and effective strategy for research and clinical intervention in prepsychotic mental disorder. <i>Schizophrenia Bulletin</i> , 2003 , 29, 771-90	1.3	212
366	Testing the Ultra High Risk (prodromal) criteria for the prediction of psychosis in a clinical sample of young people. <i>Schizophrenia Research</i> , 2006 , 84, 57-66	3.6	203
365	International clinical practice guidelines for early psychosis. <i>British Journal of Psychiatry</i> , 2005 , 48, s120-5	4.4	201

364	Basic self-disturbance predicts psychosis onset in the ultra high risk for psychosis "prodromal" population. <i>Schizophrenia Bulletin</i> , 2012 , 38, 1277-87	1.3	199
363	Generalized and specific cognitive performance in clinical high-risk cohorts: a review highlighting potential vulnerability markers for psychosis. <i>Schizophrenia Bulletin</i> , 2006 , 32, 538-55	1.3	199
362	Neuroanatomical abnormalities that predate the onset of psychosis: a multicenter study. <i>Archives of General Psychiatry</i> , 2011 , 68, 489-95		196
361	Aerobic Exercise Improves Cognitive Functioning in People With Schizophrenia: A Systematic Review and Meta-Analysis. <i>Schizophrenia Bulletin</i> , 2017 , 43, 546-556	1.3	189
360	Psychotic-like experiences in nonpsychotic help-seekers: associations with distress, depression, and disability. <i>Schizophrenia Bulletin</i> , 2006 , 32, 352-9	1.3	187
359	Outcomes of nontransitioned cases in a sample at ultra-high risk for psychosis. <i>American Journal of Psychiatry</i> , 2015 , 172, 249-58	11.9	184
358	Cognitive deficits in youth with familial and clinical high risk to psychosis: a systematic review and meta-analysis. <i>Acta Psychiatrica Scandinavica</i> , 2014 , 130, 1-15	6.5	174
357	Psychotic-like experiences and correlation with distress and depressive symptoms in a community sample of adolescents and young adults. <i>Schizophrenia Research</i> , 2010 , 119, 258-65	3.6	174
356	Pituitary volume predicts future transition to psychosis in individuals at ultra-high risk of developing psychosis. <i>Biological Psychiatry</i> , 2005 , 58, 417-23	7.9	174
355	At risk or not at risk? A meta-analysis of the prognostic accuracy of psychometric interviews for psychosis prediction. <i>World Psychiatry</i> , 2015 , 14, 322-32	14.4	172
354	Biomarkers and clinical staging in psychiatry. <i>World Psychiatry</i> , 2014 , 13, 211-23	14.4	170
353	Setting the stage: from prodrome to treatment resistance in bipolar disorder. <i>Bipolar Disorders</i> , 2007 , 9, 671-8	3.8	168
352	Neurocognitive predictors of functional outcome two to 13 years after identification as ultra-high risk for psychosis. <i>Schizophrenia Research</i> , 2011 , 132, 1-7	3.6	166
351	Effect of Ω 3 Polyunsaturated Fatty Acids in Young People at Ultrahigh Risk for Psychotic Disorders: The NEURAPRO Randomized Clinical Trial. <i>JAMA Psychiatry</i> , 2017 , 74, 19-27	14.5	162
350	Clinical staging: a heuristic and practical strategy for new research and better health and social outcomes for psychotic and related mood disorders. <i>Canadian Journal of Psychiatry</i> , 2010 , 55, 486-97	4.8	159
349	Impairment of olfactory identification ability in individuals at ultra-high risk for psychosis who later develop schizophrenia. <i>American Journal of Psychiatry</i> , 2003 , 160, 1790-4	11.9	157
348	Lessons learned from the psychosis high-risk state: towards a general staging model of prodromal intervention. <i>Psychological Medicine</i> , 2014 , 44, 17-24	6.9	149
347	Anatomic abnormalities of the anterior cingulate cortex before psychosis onset: an MRI study of ultra-high-risk individuals. <i>Biological Psychiatry</i> , 2008 , 64, 758-65	7.9	149

346	Pituitary volume in psychosis. <i>British Journal of Psychiatry</i> , 2004 , 185, 5-10	5.4	147
345	The Dark Side of the Moon: Meta-analytical Impact of Recruitment Strategies on Risk Enrichment in the Clinical High Risk State for Psychosis. <i>Schizophrenia Bulletin</i> , 2016 , 42, 732-43	1.3	144
344	At risk for schizophrenic or affective psychoses? A meta-analysis of DSM/ICD diagnostic outcomes in individuals at high clinical risk. <i>Schizophrenia Bulletin</i> , 2013 , 39, 923-32	1.3	143
343	Non-reduction in hippocampal volume is associated with higher risk of psychosis. <i>Schizophrenia Research</i> , 2002 , 58, 145-58	3.6	143
342	Road to full recovery: longitudinal relationship between symptomatic remission and psychosocial recovery in first-episode psychosis over 7.5 years. <i>Psychological Medicine</i> , 2012 , 42, 595-606	6.9	139
341	Mapping the Onset of Psychosis: The Comprehensive Assessment of At-Risk Mental States		136
340	Experience of trauma and conversion to psychosis in an ultra-high-risk (prodromal) group. <i>Acta Psychiatrica Scandinavica</i> , 2010 , 121, 377-84	6.5	134
339	Ethics and early intervention in psychosis: keeping up the pace and staying in step. <i>Schizophrenia Research</i> , 2001 , 51, 17-29	3.6	134
338	Clinical staging: a heuristic model for psychiatry and youth mental health. <i>Medical Journal of Australia</i> , 2007 , 187, S40-2	4	127
337	Insular cortex gray matter changes in individuals at ultra-high-risk of developing psychosis. <i>Schizophrenia Research</i> , 2009 , 111, 94-102	3.6	125
336	Early identification and intervention in depressive disorders: towards a clinical staging model. <i>Psychotherapy and Psychosomatics</i> , 2008 , 77, 263-70	9.4	121
335	A disturbed sense of self in the psychosis prodrome: linking phenomenology and neurobiology. <i>Neuroscience and Biobehavioral Reviews</i> , 2009 , 33, 807-17	9	116
334	Association between psychotic experiences and depression in a clinical sample over 6 months. <i>Schizophrenia Research</i> , 2007 , 91, 246-53	3.6	110
333	Medium term follow-up of a randomized controlled trial of interventions for young people at ultra high risk of psychosis. <i>Schizophrenia Research</i> , 2007 , 96, 25-33	3.6	110
332	Morphology of the anterior cingulate cortex in young men at ultra-high risk of developing a psychotic illness. <i>British Journal of Psychiatry</i> , 2003 , 182, 518-24	5.4	108
331	Stress and HPA-axis functioning in young people at ultra high risk for psychosis. <i>Journal of Psychiatric Research</i> , 2007 , 41, 561-9	5.2	106
330	Randomized controlled trial of interventions for young people at ultra high risk for psychosis: 6-month analysis. <i>Journal of Clinical Psychiatry</i> , 2011 , 72, 430-40	4.6	105
329	Randomized controlled trial of interventions for young people at ultra-high risk of psychosis: twelve-month outcome. <i>Journal of Clinical Psychiatry</i> , 2013 , 74, 349-56	4.6	105

328	Proton magnetic resonance spectroscopy in first episode psychosis and ultra high-risk individuals. <i>Schizophrenia Bulletin</i> , 2003 , 29, 831-43	1.3	101
327	Neuroimaging and emerging psychotic disorders: the Melbourne ultra-high risk studies. <i>International Review of Psychiatry</i> , 2007 , 19, 371-81	3.6	98
326	Early intervention in psychotic disorders: detection and treatment of the first episode and the critical early stages. <i>Medical Journal of Australia</i> , 2007 , 187, S8-10	4	97
325	The phenomenological critique and self-disturbance: implications for ultra-high risk ("prodrome") research. <i>Schizophrenia Bulletin</i> , 2008 , 34, 381-92	1.3	97
324	Hippocampal pathology in individuals at ultra-high risk for psychosis: a multi-modal magnetic resonance study. <i>NeuroImage</i> , 2010 , 52, 62-8	7.9	96
323	Recruitment and treatment practices for help-seeking "prodromal" patients. <i>Schizophrenia Bulletin</i> , 2007 , 33, 715-26	1.3	92
322	Social cognition in multiple sclerosis: A systematic review and meta-analysis. <i>Neurology</i> , 2016 , 87, 1727-1736	1.3	91
321	Sexual trauma increases the risk of developing psychosis in an ultra high-risk "prodromal" population. <i>Schizophrenia Bulletin</i> , 2014 , 40, 697-706	1.3	90
320	White matter volume changes in people who develop psychosis. <i>British Journal of Psychiatry</i> , 2008 , 193, 210-5	5.4	90
319	Cannabis use is not associated with the development of psychosis in an 'ultra' high-risk group. <i>Australian and New Zealand Journal of Psychiatry</i> , 2002 , 36, 800-6	2.6	89
318	Early intervention in psychosis: an overdue reform. <i>Australian and New Zealand Journal of Psychiatry</i> , 2003 , 37, 393-8	2.6	88
317	A preliminary evaluation of the validity of at-risk criteria for bipolar disorders in help-seeking adolescents and young adults. <i>Journal of Affective Disorders</i> , 2010 , 127, 316-20	6.6	86
316	Evidence-based guidelines for the pharmacological treatment of schizophrenia: Updated recommendations from the British Association for Psychopharmacology. <i>Journal of Psychopharmacology</i> , 2020 , 34, 3-78	4.6	85
315	Social cognition in clinical "at risk" for psychosis and first episode psychosis populations. <i>Schizophrenia Research</i> , 2012 , 141, 204-9	3.6	83
314	A comprehensive review and model of putative prodromal features of bipolar affective disorder. <i>Psychological Medicine</i> , 2011 , 41, 1567-77	6.9	83
313	Negative psychotic symptoms and impaired role functioning predict transition outcomes in the at-risk mental state: a latent class cluster analysis study. <i>Psychological Medicine</i> , 2013 , 43, 2311-25	6.9	82
312	Effectiveness of early intervention in psychosis. <i>Current Opinion in Psychiatry</i> , 2007 , 20, 121-5	4.9	82
311	What drives poor functioning in the at-risk mental state? A systematic review. <i>Schizophrenia Research</i> , 2014 , 159, 267-77	3.6	80

310	The predictive validity of bipolar at-risk (prodromal) criteria in help-seeking adolescents and young adults: a prospective study. <i>Bipolar Disorders</i> , 2014 , 16, 493-504	3.8	80
309	Identification of young people at risk of psychosis: validation of Personal Assessment and Crisis Evaluation Clinic intake criteria. <i>Australian and New Zealand Journal of Psychiatry</i> , 2000 , 34 Suppl, S164-9 ^{2.6}	2.6	80
308	33.1 ETHICAL ISSUES IN EARLY INTERVENTION. <i>Schizophrenia Bulletin</i> , 2019 , 45, S143-S143	1.3	78
307	Neuroimaging and treatment evidence for clinical staging in psychotic disorders: from the at-risk mental state to chronic schizophrenia. <i>Biological Psychiatry</i> , 2011 , 70, 619-25	7.9	77
306	A systematic review of the nature and correlates of sleep disturbance in early psychosis. <i>Sleep Medicine Reviews</i> , 2017 , 31, 25-38	10.2	75
305	Whither the attenuated psychosis syndrome?. <i>Schizophrenia Bulletin</i> , 2012 , 38, 1130-4	1.3	74
304	Volumetric abnormalities predating the onset of schizophrenia and affective psychoses: an MRI study in subjects at ultrahigh risk of psychosis. <i>Schizophrenia Bulletin</i> , 2012 , 38, 1083-91	1.3	73
303	Early-onset of symptoms predicts conversion to non-affective psychosis in ultra-high risk individuals. <i>Schizophrenia Research</i> , 2006 , 84, 67-76	3.6	73
302	Identification of young people at risk of psychosis: validation of Personal Assessment and Crisis Evaluation Clinic intake criteria. <i>Australian and New Zealand Journal of Psychiatry</i> , 2000 , 34, S164-S169	2.6	72
301	Hippocampal and anterior cingulate morphology in subjects at ultra-high-risk for psychosis: the role of family history of psychotic illness. <i>Schizophrenia Research</i> , 2005 , 75, 295-301	3.6	71
300	Sustained attention in young people at high risk of psychosis does not predict transition to psychosis. <i>Schizophrenia Research</i> , 2005 , 79, 127-36	3.6	71
299	Declining transition rates to psychotic disorder in "ultra-high risk" clients: Investigation of a dilution effect. <i>Schizophrenia Research</i> , 2016 , 170, 130-6	3.6	70
298	Predictive validity of clinical variables in the "at risk" for psychosis population: international comparison with results from the North American Prodrome Longitudinal Study. <i>Schizophrenia Research</i> , 2011 , 126, 51-7	3.6	69
297	Should a "Risk Syndrome for Psychosis" be included in the DSMV?. <i>Schizophrenia Research</i> , 2010 , 120, 7-15	3.6	69
296	PACE: a specialised service for young people at risk of psychotic disorders. <i>Medical Journal of Australia</i> , 2007 , 187, S43-6	4	69
295	Can we predict the onset of first-episode psychosis in a high-risk group?. <i>International Clinical Psychopharmacology</i> , 1998 , 13, S23-S30	2.2	69
294	Early intervention in psychosis: keeping faith with evidence-based health care. <i>Psychological Medicine</i> , 2010 , 40, 399-404	6.9	68
293	Clinical Staging of Psychiatric Disorders: A Heuristic Framework for Choosing Earlier, Safer and more Effective Interventions		68

292	Ultra high risk (UHR) for psychosis criteria: are there different levels of risk for transition to psychosis?. <i>Schizophrenia Research</i> , 2011 , 125, 62-8	3.6	67
291	Corpus callosum shape alterations in individuals prior to the onset of psychosis. <i>Schizophrenia Research</i> , 2008 , 103, 1-10	3.6	67
290	Exercise as an intervention for first-episode psychosis: a feasibility study. <i>Microbial Biotechnology</i> , 2018 , 12, 307-315	3.3	66
289	Validity of the 12-item General Health Questionnaire (GHQ-12) in detecting depressive and anxiety disorders among high school students. <i>Psychiatry Research</i> , 2011 , 187, 291-6	9.9	65
288	Rationale and first results of developing at-risk (prodromal) criteria for bipolar disorder. <i>Current Pharmaceutical Design</i> , 2012 , 18, 358-75	3.3	65
287	Back to the future: predicting and reshaping the course of psychotic disorder. <i>Archives of General Psychiatry</i> , 2008 , 65, 25-7		65
286	Preventative mental health care: accessing the target population. <i>Australian and New Zealand Journal of Psychiatry</i> , 1999 , 33, 912-7	2.6	65
285	The prevention of schizophrenia. <i>International Review of Psychiatry</i> , 2007 , 19, 633-46	3.6	63
284	Transdiagnostic clinical staging in youth mental health: a first international consensus statement. <i>World Psychiatry</i> , 2020 , 19, 233-242	14.4	61
283	Identification of Young People at Risk of Psychosis: Validation of Personal Assessment and Crisis Evaluation Clinic Intake Criteria. <i>Australian and New Zealand Journal of Psychiatry</i> , 2000 , 34, A164-A169	2.6	60
282	Social cognition deficits and the 'ultra high risk' for psychosis population: a review of literature. <i>Microbial Biotechnology</i> , 2011 , 5, 192-202	3.3	59
281	Prepsychotic phase of schizophrenia and related disorders: recent progress and future opportunities. <i>British Journal of Psychiatry</i> , 2005 , 48, s33-44	5.4	59
280	Treating Schizophrenia in the Prodromal Phase		59
279	Randomized controlled trial of interventions for young people at ultra-high risk of psychosis: study design and baseline characteristics. <i>Australian and New Zealand Journal of Psychiatry</i> , 2009 , 43, 818-29	2.6	58
278	Cardiometabolic risk factors in young people at ultra-high risk for psychosis: A systematic review and meta-analysis. <i>Schizophrenia Research</i> , 2016 , 170, 290-300	3.6	58
277	Diet as a hot topic in psychiatry: a population-scale study of nutritional intake and inflammatory potential in severe mental illness. <i>World Psychiatry</i> , 2018 , 17, 365-367	14.4	58
276	Prediction of a single psychotic episode: a 7.5-year, prospective study in first-episode psychosis. <i>Schizophrenia Research</i> , 2011 , 125, 236-46	3.6	55
275	The PACE Clinic: identification and management of young people at "ultra" high risk of psychosis. <i>Journal of Psychiatric Practice</i> , 2002 , 8, 255-69	1.3	54

274	Enhancing social functioning in young people at Ultra High Risk (UHR) for psychosis: A pilot study of a novel strengths and mindfulness-based online social therapy. <i>Schizophrenia Research</i> , 2018 , 202, 369-377	3.6	53
273	Superior temporal gyrus volume in antipsychotic-naive people at risk of psychosis. <i>British Journal of Psychiatry</i> , 2010 , 196, 206-11	5.4	53
272	Characterisation of the prodrome to a first episode of psychotic mania: results of a retrospective study. <i>Journal of Affective Disorders</i> , 2010 , 124, 341-5	6.6	53
271	The Validity and Value of Self-reported Physical Activity and Accelerometry in People With Schizophrenia: A Population-Scale Study of the UK Biobank. <i>Schizophrenia Bulletin</i> , 2018 , 44, 1293-1300	1.3	52
270	Antibodies to infectious agents in individuals at ultra-high risk for psychosis. <i>Biological Psychiatry</i> , 2007 , 61, 1215-7	7.9	52
269	Can an Internet-based intervention reduce suicidal ideation, depression and hopelessness among secondary school students: results from a pilot study. <i>Microbial Biotechnology</i> , 2016 , 10, 28-35	3.3	52
268	A cross-sectional exploration of the clinical characteristics of disengaged (NEET) young people in primary mental healthcare. <i>BMJ Open</i> , 2014 , 4, e006378	3	51
267	The association between early cannabis use and psychotic-like experiences in a community adolescent sample. <i>Schizophrenia Research</i> , 2009 , 112, 130-5	3.6	51
266	Is pre-psychotic intervention realistic in schizophrenia and related disorders?. <i>Australian and New Zealand Journal of Psychiatry</i> , 1997 , 31, 799-805	2.6	51
265	Imported malaria treated in Melbourne, Australia: epidemiology and clinical features in 246 patients. <i>Journal of Travel Medicine</i> , 2001 , 8, 76-81	12.9	51
264	Not all first-episode psychosis is the same: preliminary evidence of greater basic self-disturbance in schizophrenia spectrum cases. <i>Microbial Biotechnology</i> , 2013 , 7, 200-4	3.3	50
263	The ultra-high risk concept-a review. <i>Canadian Journal of Psychiatry</i> , 2013 , 58, 5-12	4.8	50
262	The longitudinal association of adolescent dating violence with psychiatric disorders and functioning. <i>Journal of Interpersonal Violence</i> , 2009 , 24, 1964-79	2.2	50
261	Clinical utility of the Mood and Anxiety Symptom Questionnaire (MASQ) in a sample of young help-seekers. <i>BMC Psychiatry</i> , 2007 , 7, 50	4.2	49
260	Cognitive decline following psychosis onset: data from the PACE clinic. <i>British Journal of Psychiatry</i> , 2007 , 51, s52-7	5.4	49
259	Commentary: The schizophrenia prodrome: a high-risk concept. <i>Schizophrenia Bulletin</i> , 2003 , 29, 859-65	1.3	49
258	Management of early psychosis in a generic adult mental health service. <i>Australian and New Zealand Journal of Psychiatry</i> , 2003 , 37, 429-36	2.6	49
257	Declining transition rates to psychosis: the contribution of potential changes in referral pathways to an ultra-high-risk service. <i>Microbial Biotechnology</i> , 2015 , 9, 200-6	3.3	48

256	Should attenuated psychosis syndrome be included in DSM-5?. <i>Lancet, The</i> , 2012 , 379, 591-2	4.0	48
255	The relationship between coping and subclinical psychotic experiences in adolescents from the general population--a longitudinal study. <i>Psychological Medicine</i> , 2011 , 41, 2535-46	6.9	48
254	Grip Strength Is Associated With Cognitive Performance in Schizophrenia and the General Population: A UK Biobank Study of 476559 Participants. <i>Schizophrenia Bulletin</i> , 2018 , 44, 728-736	1.3	47
253	Psychotic symptoms with sexual content in the "ultra high risk" for psychosis population: frequency and association with sexual trauma. <i>Psychiatry Research</i> , 2010 , 177, 84-91	9.9	47
252	Lithium suppression of tau induces brain iron accumulation and neurodegeneration. <i>Molecular Psychiatry</i> , 2017 , 22, 396-406	15.1	46
251	The Pace Clinic: Development of a Clinical Service for Young People at High Risk of Psychosis. <i>Australasian Psychiatry</i> , 1995 , 3, 345-349	1.7	45
250	Using clinical information to make individualized prognostic predictions in people at ultra high risk for psychosis. <i>Schizophrenia Research</i> , 2017 , 184, 32-38	3.6	44
249	The effects and determinants of exercise participation in first-episode psychosis: a qualitative study. <i>BMC Psychiatry</i> , 2016 , 16, 36	4.2	44
248	Prevalence of large cavum septi pellucidi in ultra high-risk individuals and patients with psychotic disorders. <i>Schizophrenia Research</i> , 2008 , 105, 236-44	3.6	44
247	The effects of vitamin and mineral supplementation on symptoms of schizophrenia: a systematic review and meta-analysis. <i>Psychological Medicine</i> , 2017 , 47, 1515-1527	6.9	43
246	Ecological momentary assessment and beyond: The rising interest in e-mental health research. <i>Journal of Psychiatric Research</i> , 2016 , 80, 3-4	5.2	43
245	Prediction and prevention of transition to psychosis in young people at incipient risk for schizophrenia. <i>American Journal of Medical Genetics Part A</i> , 2002 , 114, 929-37		43
244	Sources of clinical distress in young people at ultra high risk of psychosis. <i>Schizophrenia Research</i> , 2015 , 165, 15-21	3.6	42
243	Demographic and clinical characteristics of young people seeking help at youth mental health services: baseline findings of the Transitions Study. <i>Microbial Biotechnology</i> , 2015 , 9, 487-97	3.3	42
242	Omega-3 fatty acid supplementation in adolescents with borderline personality disorder and ultra-high risk criteria for psychosis: a post hoc subgroup analysis of a double-blind, randomized controlled trial. <i>Canadian Journal of Psychiatry</i> , 2013 , 58, 402-8	4.8	42
241	Neuroprotective effects of low-dose lithium in individuals at ultra-high risk for psychosis. A longitudinal MRI/MRS study. <i>Current Pharmaceutical Design</i> , 2012 , 18, 570-5	3.3	42
240	Neurobiological findings in early phase schizophrenia. <i>Brain Research Reviews</i> , 2000 , 31, 157-65		41
239	NEURAPRO-E study protocol: a multicentre randomized controlled trial of omega-3 fatty acids and cognitive-behavioural case management for patients at ultra high risk of schizophrenia and other psychotic disorders. <i>Microbial Biotechnology</i> , 2017 , 11, 418-428	3.3	40

238	The effectiveness of simple psychological and physical activity interventions for high prevalence mental health problems in young people: A factorial randomised controlled trial. <i>Journal of Affective Disorders</i> , 2016 , 196, 200-9	6.6	40
237	Does disturbance of self underlie social cognition deficits in schizophrenia and other psychotic disorders?. <i>Microbial Biotechnology</i> , 2009 , 3, 83-93	3.3	40
236	Neuroprotection in emerging psychotic disorders. <i>Microbial Biotechnology</i> , 2007 , 1, 114-127	3.3	40
235	Childhood maltreatment and transition to psychotic disorder independently predict long-term functioning in young people at ultra-high risk for psychosis. <i>Psychological Medicine</i> , 2015 , 45, 3453-65	6.9	39
234	Neurocognitive predictors of transition to psychosis: medium- to long-term findings from a sample at ultra-high risk for psychosis. <i>Psychological Medicine</i> , 2013 , 43, 2349-60	6.9	39
233	Discrete alterations of brain network structural covariance in individuals at ultra-high risk for psychosis. <i>Biological Psychiatry</i> , 2015 , 77, 989-96	7.9	38
232	Further examination of the reducing transition rate in ultra high risk for psychosis samples: The possible role of earlier intervention. <i>Schizophrenia Research</i> , 2016 , 174, 43-49	3.6	38
231	Reduced parahippocampal cortical thickness in subjects at ultra-high risk for psychosis. <i>Psychological Medicine</i> , 2014 , 44, 489-98	6.9	37
230	Young people at ultra high risk for psychosis: a research update. <i>Microbial Biotechnology</i> , 2011 , 5 Suppl 1, 52-7	3.3	37
229	Impact of co-occurring substance use on 6 month outcomes for young people seeking mental health treatment. <i>Australian and New Zealand Journal of Psychiatry</i> , 2007 , 41, 896-902	2.6	37
228	Amygdala and insula volumes prior to illness onset in bipolar disorder: a magnetic resonance imaging study. <i>Psychiatry Research - Neuroimaging</i> , 2012 , 201, 34-9	2.9	36
227	Neurocognitive functioning in the prodrome of mania--an exploratory study. <i>Journal of Affective Disorders</i> , 2013 , 147, 441-5	6.6	36
226	The safety and acceptability of delivering an online intervention to secondary students at risk of suicide: findings from a pilot study. <i>Microbial Biotechnology</i> , 2015 , 9, 498-506	3.3	36
225	The comprehensive assessment of at-risk mental states: from mapping the onset to mapping the structure. <i>Schizophrenia Research</i> , 2011 , 127, 107-14	3.6	36
224	A longitudinal study of obsessive-compulsive disorder in individuals at ultra-high risk for psychosis. <i>Journal of Psychiatric Research</i> , 2011 , 45, 1140-5	5.2	36
223	Antipsychotic drugs versus cognitive behavioural therapy versus a combination of both in people with psychosis: a randomised controlled pilot and feasibility study. <i>Lancet Psychiatry</i> , 2018 , 5, 411-423	23.3	35
222	Subclinical psychosis and depression: co-occurring phenomena that do not predict each other over time. <i>Schizophrenia Research</i> , 2011 , 130, 277-81	3.6	35
221	Randomized comparison of group cognitive behaviour therapy and group psychoeducation in acute patients with schizophrenia: effects on subjective quality of life. <i>Australian and New Zealand Journal of Psychiatry</i> , 2010 , 44, 144-50	2.6	35

220	Caudate nucleus volume in individuals at ultra-high risk of psychosis: a cross-sectional magnetic resonance imaging study. <i>Psychiatry Research - Neuroimaging</i> , 2010 , 182, 223-30	2.9	35
219	Childhood trauma and functional disability in psychosis, bipolar disorder and borderline personality disorder: a review of the literature. <i>Irish Journal of Psychological Medicine</i> , 2015 , 32, 21-30	3	34
218	Clinical trajectories in the ultra-high risk for psychosis population. <i>Schizophrenia Research</i> , 2018 , 197, 550-556	3.6	34
217	Sulcogyral pattern and sulcal count of the orbitofrontal cortex in individuals at ultra high risk for psychosis. <i>Schizophrenia Research</i> , 2014 , 154, 93-9	3.6	34
216	Can receipt of a regular postcard reduce suicide-related behaviour in young help seekers? A randomized controlled trial. <i>Microbial Biotechnology</i> , 2012 , 6, 145-52	3.3	34
215	Does screening high school students for psychological distress, deliberate self-harm, or suicidal ideation cause distress--and is it acceptable? An Australian-based study. <i>Crisis</i> , 2011 , 32, 254-63	2.8	34
214	The pro-cognitive mechanisms of physical exercise in people with schizophrenia. <i>British Journal of Pharmacology</i> , 2017 , 174, 3161-3172	8.6	33
213	Association Between Muscular Strength and Cognition in People With Major Depression or Bipolar Disorder and Healthy Controls. <i>JAMA Psychiatry</i> , 2018 , 75, 740-746	14.5	33
212	Measuring psychosocial outcome is good. <i>Current Opinion in Psychiatry</i> , 2013 , 26, 138-43	4.9	33
211	Cannabis use and symptom severity in individuals at ultra high risk for psychosis: a meta-analysis. <i>Acta Psychiatrica Scandinavica</i> , 2017 , 136, 5-15	6.5	32
210	Externalized attributional bias in the Ultra High Risk (UHR) for psychosis population. <i>Psychiatry Research</i> , 2013 , 206, 200-5	9.9	32
209	Substance use in youth at risk for psychosis. <i>Schizophrenia Research</i> , 2017 , 181, 23-29	3.6	32
208	Can we detect psychotic-like experiences in the general population?. <i>Current Pharmaceutical Design</i> , 2012 , 18, 376-85	3.3	32
207	Bipolar At-Risk Criteria: An Examination of Which Clinical Features Have Optimal Utility for Identifying Youth at Risk of Early Transition From Depression to Bipolar Disorders. <i>Schizophrenia Bulletin</i> , 2017 , 43, 737-744	1.3	31
206	Baseline grey matter volume of non-transitioned "ultra high risk" for psychosis individuals with and without attenuated psychotic symptoms at long-term follow-up. <i>Schizophrenia Research</i> , 2016 , 173, 152-158	3.6	31
205	Cost implications of specific and non-specific treatment for young persons at ultra high risk of developing a first episode of psychosis. <i>Microbial Biotechnology</i> , 2009 , 3, 28-34	3.3	31
204	Cannabis-induced attenuated psychotic symptoms: implications for prognosis in young people at ultra-high risk for psychosis. <i>Psychological Medicine</i> , 2017 , 47, 616-626	6.9	30
203	Metacognitive beliefs in the at-risk mental state: A systematic review and meta-analysis. <i>Behaviour Research and Therapy</i> , 2017 , 90, 25-31	5.2	30

202	History of trauma and the association with baseline symptoms in an Ultra-High Risk for psychosis cohort. <i>Psychiatry Research</i> , 2013 , 210, 75-81	9.9	30
201	Depressed mood as a risk factor for unprotected sex in young people. <i>Australasian Psychiatry</i> , 2006 , 14, 310-312	1.7	29
200	Transition to first episode psychosis in ultra high risk populations: does baseline functioning hold the key?. <i>Schizophrenia Research</i> , 2013 , 143, 132-7	3.6	28
199	Adhesio interthalamica in individuals at high-risk for developing psychosis and patients with psychotic disorders. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2008 , 32, 1708-14	5.5	28
198	Is basic self-disturbance in ultra-high risk for psychosis ('prodromal') patients associated with borderline personality pathology?. <i>Microbial Biotechnology</i> , 2013 , 7, 306-10	3.3	27
197	Follow-up factor structure of schizotypy and its clinical associations in a help-seeking sample meeting ultra-high risk for psychosis criteria at baseline. <i>Comprehensive Psychiatry</i> , 2013 , 54, 173-80	7.3	26
196	Psychotic-like experiences as overdetermined phenomena: when do they increase risk for psychotic disorder?. <i>Schizophrenia Research</i> , 2009 , 108, 303-4	3.6	26
195	Effects of NRG1 and DAOA genetic variation on transition to psychosis in individuals at ultra-high risk for psychosis. <i>Translational Psychiatry</i> , 2013 , 3, e251	8.6	25
194	Motives for substance use among young people seeking mental health treatment. <i>Microbial Biotechnology</i> , 2008 , 2, 188-94	3.3	25
193	Met and unmet need in youth mental health. <i>Journal of Mental Health</i> , 2008 , 17, 618-628	2.7	25
192	Neurocognition as a predictor of transition to psychotic disorder and functional outcomes in ultra-high risk participants: Findings from the NEURAPRO randomized clinical trial. <i>Schizophrenia Research</i> , 2019 , 206, 67-74	3.6	25
191	Transitions Study of predictors of illness progression in young people with mental ill health: study methodology. <i>Microbial Biotechnology</i> , 2015 , 9, 38-47	3.3	24
190	Can clinicians predict psychosis in an ultra high risk group?. <i>Australian and New Zealand Journal of Psychiatry</i> , 2010 , 44, 625-30	2.6	24
189	Early intervention in psychosis: evidence, evidence gaps, criticism, and confusion. <i>Australian and New Zealand Journal of Psychiatry</i> , 2012 , 46, 7-9	2.6	24
188	Persistent negative symptoms in individuals at Ultra High Risk for psychosis. <i>Schizophrenia Research</i> , 2019 , 206, 355-361	3.6	24
187	Antipsychotic medication versus psychological intervention versus a combination of both in adolescents with first-episode psychosis (MAPS): a multicentre, three-arm, randomised controlled pilot and feasibility study. <i>Lancet Psychiatry</i> , 2020 , 7, 788-800	23.3	23
186	Examining the association between social cognition and functioning in individuals at ultra-high risk for psychosis. <i>Australian and New Zealand Journal of Psychiatry</i> , 2017 , 51, 83-92	2.6	22
185	Psychotic experience subtypes, poor mental health status and help-seeking behaviour in a community sample of young adults. <i>Microbial Biotechnology</i> , 2012 , 6, 300-8	3.3	22

184	Psychotic experiences and their significance. <i>World Psychiatry</i> , 2016 , 15, 130-1	14.4	22
183	The NEURAPRO Biomarker Analysis: Long-Chain Omega-3 Fatty Acids Improve 6-Month and 12-Month Outcomes in Youths at Ultra-High Risk for Psychosis. <i>Biological Psychiatry</i> , 2020 , 87, 243-252	7.9	22
182	Social environmental risk factors for transition to psychosis in an Ultra-High Risk population. <i>Schizophrenia Research</i> , 2015 , 161, 150-5	3.6	21
181	Does specific psychopathology predict development of psychosis in ultra high-risk (UHR) patients?. <i>Australian and New Zealand Journal of Psychiatry</i> , 2013 , 47, 380-90	2.6	21
180	Managing deliberate self-harm in young people: an evaluation of a training program developed for school welfare staff using a longitudinal research design. <i>BMC Psychiatry</i> , 2008 , 8, 75	4.2	21
179	The Ultra-High-Risk for psychosis groups: Evidence to maintain the status quo. <i>Schizophrenia Research</i> , 2018 , 195, 543-548	3.6	20
178	Borderline personality features and development of psychosis in an 'ultra high risk' (UHR) population: a case control study. <i>Microbial Biotechnology</i> , 2012 , 6, 247-55	3.3	20
177	Should a risk syndrome for first episode psychosis be included in the DSM-5?. <i>Current Opinion in Psychiatry</i> , 2011 , 24, 128-33	4.9	20
176	Treatment of people at ultra-high risk for psychosis. <i>World Psychiatry</i> , 2017 , 16, 207-208	14.4	19
175	Long-term employment among people at ultra-high risk for psychosis. <i>Schizophrenia Research</i> , 2017 , 184, 26-31	3.6	19
174	Monitoring of physical health in services for young people at ultra-high risk of psychosis. <i>Microbial Biotechnology</i> , 2018 , 12, 153-159	3.3	19
173	Physical health promotion for young people at ultra-high risk for psychosis: An application of the COM-B model and behaviour-change wheel. <i>International Journal of Mental Health Nursing</i> , 2016 , 25, 536-545	3.8	19
172	Altered depth of the olfactory sulcus in ultra high-risk individuals and patients with psychotic disorders. <i>Schizophrenia Research</i> , 2014 , 153, 18-24	3.6	19
171	Dynamic association between interpersonal functioning and positive symptom dimensions of psychosis over time: a longitudinal study of healthy adolescents. <i>Schizophrenia Bulletin</i> , 2013 , 39, 179-85 ^{1.3}		19
170	The ultra high risk approach to define psychosis risk. <i>Current Pharmaceutical Design</i> , 2012 , 18, 346-50	3.3	19
169	Brain changes during the onset of schizophrenia: implications for neurodevelopmental theories. <i>Medical Journal of Australia</i> , 2009 , 190, S10-3	4	19
168	Examining the physical health and lifestyle of young people at ultra-high risk for psychosis: A qualitative study involving service users, parents and clinicians. <i>Psychiatry Research</i> , 2017 , 255, 87-93	9.9	18
167	Preferences and motivations for exercise in early psychosis. <i>Acta Psychiatrica Scandinavica</i> , 2016 , 134, 83-4	6.5	18

166	Olfactory identification deficits at identification as ultra-high risk for psychosis are associated with poor functional outcome. <i>Schizophrenia Research</i> , 2015 , 161, 156-62	3.6	18
165	Validation of a short adaptation of the Mood and Anxiety Symptoms Questionnaire (MASQ) in adolescents and young adults. <i>Psychiatry Research</i> , 2014 , 215, 778-83	9.9	18
164	Borderline personality pathology in young people at ultra high risk of developing a psychotic disorder. <i>Microbial Biotechnology</i> , 2017 , 11, 208-214	3.3	17
163	Opening the Black Box of Cognitive-Behavioural Case Management in Clients with Ultra-High Risk for Psychosis. <i>Psychotherapy and Psychosomatics</i> , 2017 , 86, 292-299	9.4	17
162	Do affective or dissociative symptoms mediate the association between childhood sexual trauma and transition to psychosis in an ultra-high risk cohort?. <i>Psychiatry Research</i> , 2016 , 236, 182-185	9.9	17
161	The development of a randomised controlled trial testing the effects of an online intervention among school students at risk of suicide. <i>BMC Psychiatry</i> , 2014 , 14, 155	4.2	17
160	Effects of drop-out on efficacy estimates in five Cochrane reviews of popular antipsychotics for schizophrenia. <i>Acta Psychiatrica Scandinavica</i> , 2012 , 126, 1-11	6.5	17
159	Structural abnormalities in the cuneus associated with Herpes Simplex Virus (type 1) infection in people at ultra high risk of developing psychosis. <i>Schizophrenia Research</i> , 2012 , 135, 175-80	3.6	17
158	Depression Education and Identification in Schools: An Australian-based Study. <i>School Mental Health</i> , 2010 , 2, 13-22	2.6	17
157	A factor analytic investigation of the Tripartite model of affect in a clinical sample of young Australians. <i>BMC Psychiatry</i> , 2008 , 8, 79	4.2	17
156	Improving collaboration between private psychiatrists, the public mental health sector and general practitioners: evaluation of the Partnership Project. <i>Australian and New Zealand Journal of Psychiatry</i> , 2004 , 38, 125-34	2.6	17
155	Depressed mood as a risk factor for unprotected sex in young people. <i>Australasian Psychiatry</i> , 2006 , 14, 310-2	1.7	17
154	Association of Structural Magnetic Resonance Imaging Measures With Psychosis Onset in Individuals at Clinical High Risk for Developing Psychosis: An ENIGMA Working Group Mega-analysis. <i>JAMA Psychiatry</i> , 2021 , 78, 753-766	14.5	17
153	The effectiveness of simple psychological and exercise interventions for high prevalence mental health problems in young people: a factorial randomised controlled trial. <i>Trials</i> , 2011 , 12, 76	2.8	16
152	When things are not as they seem: detecting first-episode psychosis upon referral to ultra high risk (prodromal) clinics. <i>Microbial Biotechnology</i> , 2007 , 1, 208-211	3.3	16
151	How age and gender predict illness course in a first-episode nonaffective psychosis cohort. <i>Journal of Clinical Psychiatry</i> , 2016 , 77, e283-9	4.6	16
150	Comorbidity rates of depression and anxiety in first episode psychosis: A systematic review and meta-analysis. <i>Schizophrenia Research</i> , 2020 , 216, 322-329	3.6	16
149	Long-term maintenance and effects of exercise in early psychosis. <i>Microbial Biotechnology</i> , 2018 , 12, 578-585	3.3	15

148	Markers of vulnerability to obsessive-compulsive disorder in an ultra-high risk sample of patients who developed psychosis. <i>Microbial Biotechnology</i> , 2012 , 6, 201-6	3.3	15
147	Handgrip Strength Is Associated With Hippocampal Volume and White Matter Hyperintensities in Major Depression and Healthy Controls: A UK Biobank Study. <i>Psychosomatic Medicine</i> , 2020 , 82, 39-46	3.7	15
146	Terminology and assessment tools of psychosis: A systematic narrative review. <i>Psychiatry and Clinical Neurosciences</i> , 2020 , 74, 226-246	6.2	15
145	Neuroanatomical Predictors of Functional Outcome in Individuals at Ultra-High Risk for Psychosis. <i>Schizophrenia Bulletin</i> , 2017 , 43, 449-458	1.3	15
144	Prolonged duration of untreated psychosis: a problem that needs addressing. <i>Microbial Biotechnology</i> , 2017 , 11, 263-268	3.3	14
143	Comparison of erythrocyte omega-3 index, fatty acids and molecular phospholipid species in people at ultra-high risk of developing psychosis and healthy people. <i>Schizophrenia Research</i> , 2020 , 226, 44-51	3.6	14
142	The reality of at risk mental state services: a response to recent criticisms. <i>Psychological Medicine</i> , 2021 , 51, 212-218	6.9	14
141	Youth depression alleviation: the Fish Oil Youth Depression Study (YoDA-F): A randomized, double-blind, placebo-controlled treatment trial. <i>Microbial Biotechnology</i> , 2016 , 10, 290-9	3.3	13
140	Dynamic prediction of transition to psychosis using joint modelling. <i>Schizophrenia Research</i> , 2018 , 202, 333-340	3.6	13
139	Relationship Between Polyunsaturated Fatty Acids and Psychopathology in the NEURAPRO Clinical Trial. <i>Frontiers in Psychiatry</i> , 2019 , 10, 393	5	13
138	Development and validation of a new measure of everyday adolescent functioning: the multidimensional adolescent functioning scale. <i>Journal of Adolescent Health</i> , 2013 , 52, 195-200	5.8	13
137	Ventricular volumes across stages of schizophrenia and other psychoses. <i>Australian and New Zealand Journal of Psychiatry</i> , 2017 , 51, 1041-1051	2.6	13
136	Neighbourhood characteristics and the rate of identification of young people at ultra-high risk for psychosis. <i>Schizophrenia Research</i> , 2015 , 169, 214-216	3.6	13
135	Identification and treatment of the prodromal phase of psychotic disorders: perspectives from the PACE Clinic. <i>Microbial Biotechnology</i> , 2007 , 1, 224-235	3.3	13
134	Obstetric complications and transition to psychosis in an "ultra" high risk sample. <i>Australian and New Zealand Journal of Psychiatry</i> , 2005 , 39, 460-6	2.6	13
133	Patterns of current comorbidity in young help-seekers: implications for service planning and delivery. <i>Australasian Psychiatry</i> , 2005 , 13, 379-83	1.7	13
132	Early Signs Monitoring to Prevent Relapse in Psychosis and Promote Well-Being, Engagement, and Recovery: Protocol for a Feasibility Cluster Randomized Controlled Trial Harnessing Mobile Phone Technology Blended With Peer Support. <i>JMIR Research Protocols</i> , 2020 , 9, e15058	2	13
131	Longitudinal Cognitive Performance in Individuals at Ultrahigh Risk for Psychosis: A 10-year Follow-up. <i>Schizophrenia Bulletin</i> , 2019 , 45, 1101-1111	1.3	13

130	Challenges in implementing an exercise intervention within residential psychiatric care: A mixed methods study. <i>Mental Health and Physical Activity</i> , 2017 , 12, 141-146	5	12
129	Prescribing in schizophrenia and psychosis: Increasing polypharmacy over time. <i>Human Psychopharmacology</i> , 2017 , 32, e2579	2.3	12
128	Self reported rates of criminal offending and victimization in young people at-risk for psychosis. <i>Schizophrenia Research</i> , 2015 , 166, 55-9	3.6	12
127	Identifying the ultra-high risk (prodromal) population: evaluation of training workshops with mental health services. <i>Australian and New Zealand Journal of Psychiatry</i> , 2008 , 42, 236-43	2.6	12
126	A clinical trials agenda for testing interventions in earlier stages of psychotic disorders. <i>Medical Journal of Australia</i> , 2009 , 190, S33-6	4	12
125	Distinguishing between anxiety and depression using the Mood and Anxiety Symptoms Questionnaire (MASQ). <i>British Journal of Clinical Psychology</i> , 2007 , 46, 235-9	3.6	12
124	Outcome of suicidal ideation and behavior in a young, help-seeking population over a 2-year period. <i>Crisis</i> , 2007 , 28, 4-10	2.8	12
123	Multiple lifestyle factors and depressed mood: a cross-sectional and longitudinal analysis of the UK Biobank (N = 84,860). <i>BMC Medicine</i> , 2020 , 18, 354	11.4	12
122	Distress in relation to attenuated psychotic symptoms in the ultra-high-risk population is not associated with increased risk of psychotic disorder. <i>Microbial Biotechnology</i> , 2016 , 10, 258-62	3.3	12
121	Prevention and treatment of long-term social disability amongst young people with emerging severe mental illness with social recovery therapy (The PRODIGY Trial): study protocol for a randomised controlled trial. <i>Trials</i> , 2017 , 18, 315	2.8	11
120	Adjunctive nutrients in first-episode psychosis: A systematic review of efficacy, tolerability and neurobiological mechanisms. <i>Microbial Biotechnology</i> , 2018 , 12, 774-783	3.3	11
119	Young people at ultra high risk for psychosis: research from the PACE clinic. <i>Revista Brasileira De Psiquiatria</i> , 2011 , 33 Suppl 2, s143-60	2.6	11
118	Public and Private Psychiatry: Can They Work Together and is it Worth the Effort?. <i>Australian and New Zealand Journal of Psychiatry</i> , 2005 , 39, 67-73	2.6	11
117	Perceived stress and psychosis: The effect of perceived stress on psychotic-like experiences in a community sample of adolescents. <i>Microbial Biotechnology</i> , 2019 , 13, 1465-1469	3.3	11
116	Changes in metabolic parameters in patients with severe mental illness over a 10-year period: A retrospective cohort study. <i>Australian and New Zealand Journal of Psychiatry</i> , 2017 , 51, 75-82	2.6	10
115	Are UHR patients who present with hallucinations alone at lower risk of transition to psychosis?. <i>Psychiatry Research</i> , 2016 , 235, 177-96	9.9	10
114	Evaluating complex, collaborative programmes: the Partnership Project as a case study. <i>Australian and New Zealand Journal of Psychiatry</i> , 2001 , 35, 639-46	2.6	10
113	Adulthood revictimization: looking beyond childhood trauma. <i>Acta Psychiatrica Scandinavica</i> , 2016 , 134, 368	6.5	10

112	A randomised, double-blind, placebo-controlled trial of minocycline and/or omega-3 fatty acids added to treatment as usual for at-risk mental states (NAYAB): study protocol. <i>Trials</i> , 2017 , 18, 524	2.8	9
111	Care management service and falls prevention: a case-control study in a Chinese population. <i>Journal of Aging and Health</i> , 2010 , 22, 348-61	2.6	9
110	Study protocol: the development of a randomised controlled trial testing a postcard intervention designed to reduce suicide risk among young help-seekers. <i>BMC Psychiatry</i> , 2009 , 9, 59	4.2	9
109	Study protocol: The development of a pilot study employing a randomised controlled design to investigate the feasibility and effects of a peer support program following discharge from a specialist first-episode psychosis treatment centre. <i>BMC Psychiatry</i> , 2010 , 10, 37	4.2	9
108	Recognition and Treatment of the Pre-psychotic Phase of Psychotic Disorders 2001 , 101-122		9
107	Preventive interventions for individuals at ultra high risk for psychosis: An updated and extended meta-analysis. <i>Clinical Psychology Review</i> , 2021 , 86, 102005	10.8	9
106	The role of coping in the association between subclinical psychotic experiences and functioning: A within study replication in two independent adolescent samples. <i>Schizophrenia Research</i> , 2018 , 201, 91-97	3.6	8
105	Cognitive precursors of severe mental disorders. <i>Cognitive Neuropsychiatry</i> , 2013 , 18, 1-8	2	8
104	Does cognitive behavioural therapy have a role in improving problem solving and coping in adolescents with suicidal ideation?. <i>The Cognitive Behaviour Therapist</i> , 2014 , 7,	2.9	8
103	Phenylthiocarbamide (PTC) perception in ultra-high risk for psychosis participants who develop schizophrenia: testing the evidence for an endophenotypic marker. <i>Psychiatry Research</i> , 2012 , 199, 8-11	9.9	8
102	Cognitive and olfactory deficits: Course from high risk to first-episode psychosis. <i>Schizophrenia Research</i> , 1998 , 29, 54-55	3.6	8
101	Evidence for preventive treatments in young patients at clinical high risk of psychosis: the need for context. <i>Lancet Psychiatry</i> , 2020 , 7, 378-380	23.3	8
100	Basic symptoms in young people at ultra-high risk of psychosis: Association with clinical characteristics and outcomes. <i>Schizophrenia Research</i> , 2020 , 216, 255-261	3.6	8
99	Distress related to psychotic symptoms in individuals at high risk of psychosis. <i>Schizophrenia Research</i> , 2020 , 215, 66-73	3.6	8
98	Importance of Variable Selection in Multimodal Prediction Models in Patients at Clinical High Risk for Psychosis and Recent-Onset Depression. <i>JAMA Psychiatry</i> , 2019 , 76, 339	14.5	7
97	Implementing an innovative intervention to increase research capacity for enhancing early psychosis care in Indonesia. <i>Journal of Psychiatric and Mental Health Nursing</i> , 2017 , 24, 671-680	2.4	7
96	The Singapore flagship programme in translational and clinical research in psychosis. <i>Microbial Biotechnology</i> , 2011 , 5, 290-300	3.3	7
95	The 'youth model' in mental health services. <i>Australasian Psychiatry</i> , 2008 , 16, 136-7	1.7	7

94	Ventricle volumes in emerging psychosis. A cross-sectional and longitudinal MRI study. <i>European Psychiatry</i> , 2007 , 22, S30-S31	6	7
93	Trajectories of symptom severity and functioning over a three-year period in a psychosis high-risk sample: A secondary analysis of the Neurapro trial. <i>Behaviour Research and Therapy</i> , 2020 , 124, 103527	5.2	7
92	The relationship between personality traits and psychotic like experiences in a large non-clinical adolescent sample. <i>Personality and Individual Differences</i> , 2015 , 73, 92-97	3.3	6
91	Study protocol for a randomised controlled trial of CBT vs antipsychotics vs both in 14-18-year-olds: Managing Adolescent first episode Psychosis: a Feasibility study (MAPS). <i>Trials</i> , 2019 , 20, 395	2.8	6
90	Transcending false dichotomies and diagnostic silos to reduce disease burden in mental disorders. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2020 , 55, 1095-1103	4.5	6
89	Can youth at high risk of illness progression be identified by measures of rumination and sleep-wake disturbance. <i>Microbial Biotechnology</i> , 2019 , 13, 1214-1219	3.3	6
88	A qualitative study of the prodrome to first-episode major depressive disorder in adolescents. <i>Psychopathology</i> , 2015 , 48, 153-61	3.4	5
87	Prognostic factors for progression to psychosis in high risk youth. <i>Evidence-Based Mental Health</i> , 2008 , 11, 72	11.1	5
86	0462 PSYCHIATRIC OUTPATIENT CLINIC FOR EMERGING PSYCHOLOGICAL DISORDERS (PREPP PROGRAM): VALIDATION OF THE FRENCH VERSION OF CAARMS. <i>Schizophrenia Research</i> , 2006 , 86, S80	3.6	5
85	Depression, Substance Use and Suicidality in Help-Seeking Adolescents: A Survey of Prevalence. <i>Australian Journal of Guidance and Counselling</i> , 2004 , 14, 162-175		5
84	Keeping an open mind: Investigating options for treatment of the pre-psychotic phase. <i>Journal of Mental Health</i> , 2003 , 12, 341-343	2.7	5
83	Developing Partnerships Between Public and Private Psychiatry. <i>Australasian Psychiatry</i> , 2000 , 8, 332-335.	7	5
82	Duration of Untreated Psychosis: Getting Both the Timing and the Sample Right. <i>American Journal of Psychiatry</i> , 2020 , 177, 1183	11.9	5
81	Exploring functional impairment in young people at ultra-high risk for psychosis: A qualitative study. <i>Microbial Biotechnology</i> , 2019 , 13, 789-797	3.3	5
80	Combined individual and family therapy in comparison to treatment as usual for people at-risk of psychosis: A feasibility study (IF CBT): Trial rationale, methodology and baseline characteristics. <i>Microbial Biotechnology</i> , 2021 , 15, 140-148	3.3	5
79	Investigating the Agreement Between Cardiovascular Disease Risk Calculators Among People Diagnosed With Schizophrenia. <i>Frontiers in Psychiatry</i> , 2018 , 9, 685	5	5
78	Longitudinal determinants of client treatment satisfaction in an intensive first-episode psychosis treatment programme. <i>Microbial Biotechnology</i> , 2017 , 11, 354-362	3.3	4
77	The relationship between childhood trauma and clinical characteristics in ultra-high risk for psychosis youth. <i>Psychosis</i> , 2019 , 11, 28-41	1.2	4

76	The value of early intervention in creating the new mental health system: Response to Allison et al. <i>Australian and New Zealand Journal of Psychiatry</i> , 2020 , 54, 667-669	2.6	4
75	Cognitive functioning in ultra-high risk for psychosis individuals with and without depression: Secondary analysis of findings from the NEURAPRO randomized clinical trial. <i>Schizophrenia Research</i> , 2020 , 218, 48-54	3.6	4
74	Moving beyond the weight-loss paradigm of exercise interventions for mental illness. <i>Psychiatry Research</i> , 2016 , 246, 392-393	9.9	4
73	Film review: Mental. <i>Microbial Biotechnology</i> , 2010 , 4, 6-6	3.3	4
72	People at risk of schizophrenia and other psychoses: comments on the Edinburgh High-Risk Study. <i>British Journal of Psychiatry</i> , 1999 , 175, 586-7	5.4	4
71	Cannabidiol for at risk for psychosis youth: A randomized controlled trial. <i>Microbial Biotechnology</i> , 2021 ,	3.3	4
70	Perceptual abnormalities in an ultra-high risk for psychosis population relationship to trauma and co-morbid disorder. <i>Microbial Biotechnology</i> , 2019 , 13, 231-240	3.3	4
69	The prognostic significance of attenuated psychotic symptoms in help-seeking youth. <i>Schizophrenia Research</i> , 2020 , 215, 277-283	3.6	4
68	Prediction of clinical outcomes beyond psychosis in the ultra-high risk for psychosis population. <i>Microbial Biotechnology</i> , 2021 , 15, 642-651	3.3	4
67	A pilot randomised controlled trial comparing antipsychotic medication, to cognitive behavioural therapy to a combination of both in people with psychosis: rationale, study design and baseline data of the COMPARE trial. <i>Psychosis</i> , 2017 , 9, 193-204	1.2	3
66	At-risk mental states 2020 , 47-57		3
65	Omega-3 fatty acids and neurocognitive ability in young people at ultra-high risk for psychosis. <i>Microbial Biotechnology</i> , 2021 , 15, 874-881	3.3	3
64	Intervention strategies for ultra-high risk for psychosis: Progress in delaying the onset and reducing the impact of first-episode psychosis. <i>Schizophrenia Research</i> , 2021 , 228, 344-356	3.6	3
63	Do schizotypal or borderline personality disorders predict onset of psychotic disorder or persistent attenuated psychotic symptoms in patients at high clinical risk?. <i>Schizophrenia Research</i> , 2020 , 220, 275-277	3.6	2
62	Commentary: Preventive Treatments for Psychosis: Umbrella Review (Just the Evidence). <i>Frontiers in Psychiatry</i> , 2020 , 11, 488	5	2
61	Impaired olfactory ability associated with larger left hippocampus and rectus volumes at earliest stages of schizophrenia: A sign of neuroinflammation?. <i>Psychiatry Research</i> , 2020 , 289, 112909	9.9	2
60	LONG TERM FOLLOW UP OF AN ULTRA HIGH RISK ("PRODROMAL") GROUP. <i>Schizophrenia Research</i> , 2010 , 117, 179	3.6	2
59	TRAUMA IN THE ULTRA HIGH RISK (PRODROMAL) POPULATION. <i>Schizophrenia Research</i> , 2008 , 102, 156	3.6	2

58	Amygdala Volume Status Might Reflect Dominant Mode of Emotional Information Processing Reply. <i>Archives of General Psychiatry</i> , 2007 , 64, 252		2
57	Precursors of schizophrenia. <i>Current Psychosis & Therapeutics Reports</i> , 2004 , 2, 67-72		2
56	Public and private psychiatry: can they work together and is it worth the effort?. <i>Australian and New Zealand Journal of Psychiatry</i> , 2005 , 39, 67-73	2.6	2
55	Evaluating Complex, Collaborative Programmes: The Partnership Project as a Case Study		2
54	Early persistence of psychotic-like experiences in a community sample of adolescents. <i>Microbial Biotechnology</i> , 2021 , 15, 1575-1583	3.3	2
53	Does cortical brain morphology act as a mediator between childhood trauma and transition to psychosis in young individuals at ultra-high risk?. <i>Schizophrenia Research</i> , 2020 , 224, 116-125	3.6	2
52	Examining healthcare professionals' beliefs and actions regarding the physical health of people with schizophrenia. <i>BMC Health Services Research</i> , 2020 , 20, 771	2.9	2
51	Individualised prediction of psychosis in individuals meeting at-risk mental state (ARMS) criteria: protocol for a systematic review of clinical prediction models. <i>Diagnostic and Prognostic Research</i> , 2019 , 3, 21	5.5	2
50	Ultra-high risk phase: A missed opportunity for physical health care. <i>Microbial Biotechnology</i> , 2018 , 12, 267-268	3.3	2
49	Pre-onset risk characteristics for mania among young people at clinical high risk for psychosis. <i>Schizophrenia Research</i> , 2018 , 192, 345-350	3.6	2
48	The EMPOWER blended digital intervention for relapse prevention in schizophrenia: a feasibility cluster randomised controlled trial in Scotland and Australia.. <i>Lancet Psychiatry</i> , 2022 , 9, 477-486	23.3	2
47	The structure of the ultra high risk mental state for psychosis. A latent class cluster analysis study. <i>European Psychiatry</i> , 2011 , 26, 2087-2087	6	1
46	Letter to the editor. The need for drug-naive research in first-episode psychosis: a response to Moncrieff & Leo (2010). <i>Psychological Medicine</i> , 2011 , 41, 1117-8	6.9	1
45	Selective bias in criticism of early intervention. <i>Australian and New Zealand Journal of Psychiatry</i> , 2012 , 46, 904-5	2.6	1
44	S.14.02 Transition to psychosis □what happens in the brain?. <i>European Neuropsychopharmacology</i> , 2008 , 18, S176	1.2	1
43	Obstetric complications and transition to psychosis in an 'ultra' high risk sample. <i>Australian and New Zealand Journal of Psychiatry</i> , 2005 , 39, 460-466	2.6	1
42	Proportion and characteristics of young people in a first-episode psychosis clinic who first attended an at-risk mental state service or other specialist youth mental health service.. <i>Schizophrenia Research</i> , 2022 , 241, 94-101	3.6	1
41	Improving Collaboration Between Private Psychiatrists, the Public Mental Health Sector and General Practitioners: Evaluation of the Partnership Project		1

40	First episode research. <i>Current Opinion in Psychiatry</i> , 1998 , 11, 27-32	4.9	1
39	The Impact of Early Intervention in Schizophrenia 2010 , 299-316		1
38	Improving Mood with Physical ACTivity (IMPACT) trial: a cluster randomised controlled trial to determine the effectiveness of a brief physical activity behaviour change intervention on depressive symptoms in young people, compared with psychoeducation, in addition to routine clinical care within youth mental health services-a protocol study. <i>BMJ Open</i> , 2019 , 9, e034002	3	1
37	More frequent lithium testing in UK Primary Care associates with a lower hospital admission rate for bipolar disorder-What this can tell us about GP practice engagement with this group. <i>International Journal of Clinical Practice</i> , 2019 , 73, e13277	2.9	1
36	Greater preference for eveningness is associated with negative symptoms in an ultra-high risk for psychosis sample. <i>Microbial Biotechnology</i> , 2021 , 15, 1793-1798	3.3	1
35	Prevalence of Major Cardiovascular Disease Events Among People Diagnosed With Schizophrenia Who Have Sleep Disturbance, Sedentary Behavior, or Muscular Weakness.. <i>Schizophrenia Bulletin Open</i> , 2021 , 2, sgaa069	2.2	1
34	Clinical and cost-effectiveness of social recovery therapy for the prevention and treatment of long-term social disability among young people with emerging severe mental illness (PRODIGY): randomised controlled trial.. <i>British Journal of Psychiatry</i> , 2022 , 1-9	5.4	0
33	Outcomes for first-episode psychosis after entry via an at-risk mental state clinic compared to direct entry to a first episode of psychosis service: A systematic review and meta-analysis.. <i>Schizophrenia Research</i> , 2022 , 240, 214-219	3.6	0
32	Intertwined associations between oxidative and nitrosative stress and endocannabinoid system pathways: Relevance for neuropsychiatric disorders. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2021 , 110481	5.5	0
31	The association of plasma inflammatory markers with omega-3 fatty acids and their mediating role in psychotic symptoms and functioning: An analysis of the NEURAPRO clinical trial. <i>Brain, Behavior, and Immunity</i> , 2022 , 99, 147-156	16.6	0
30	Validation of the Dutch version of the Multidimensional Adolescent Functioning Scale (MAFS). <i>Health and Quality of Life Outcomes</i> , 2020 , 18, 306	3	0
29	Supplementation with the omega-3 long chain polyunsaturated fatty acids: Changes in the concentrations of omega-3 index, fatty acids and molecular phospholipids of people at ultra high risk of developing psychosis. <i>Schizophrenia Research</i> , 2020 , 226, 52-60	3.6	0
28	Examining the feasibility, acceptability, validity and reliability of physical activity, sedentary behaviour and sleep measures in people with schizophrenia. <i>Mental Health and Physical Activity</i> , 2021 , 21, 100415	5	0
27	Patients', carers' and clinicians' attitudes towards alternative terms to describe the at-risk for psychosis state. <i>Schizophrenia Research</i> , 2021 , 237, 69-75	3.6	0
26	The association between migrant status and transition in an ultra-high risk for psychosis population. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2021 , 56, 943-952	4.5	0
25	A review of economic evaluations of health care for people at risk of psychosis and for first-episode psychosis.. <i>BMC Psychiatry</i> , 2022 , 22, 126	4.2	0
24	After the acute crisis - engaging people with psychosis in rehabilitation-oriented care.. <i>World Psychiatry</i> , 2022 , 21, 246-247	14.4	0
23	Digital smartphone intervention to recognise and manage early warning signs in schizophrenia to prevent relapse: the EMPOWER feasibility cluster RCT. <i>Health Technology Assessment</i> , 2022 , 26, 1-174	4.4	0

22	Making Early Intervention more equitable. <i>Australian and New Zealand Journal of Psychiatry</i> , 2017 , 51, 534-535	2.6
21	Lifestyle factors may be linked to symptoms of metabolic syndrome in people at risk for psychosis. <i>Schizophrenia Research</i> , 2017 , 183, 47-48	3.6
20	Gym rats: exercise reverses cognitive impairment in the phencyclidine rat model of schizophrenia. <i>European Neuropsychopharmacology</i> , 2017 , 27, S72	1.2
19	Stand-alone cognitive behavioural therapy is not in clinical equipoise with antipsychotic treatment - Author's reply. <i>Lancet Psychiatry</i> , 2018 , 5, 540-541	23.3
18	Women with schizophrenia are at increased risk of breast cancer. <i>Evidence-Based Mental Health</i> , 2018 , 21, e13	11.1
17	Comment on Lee et al. <i>Schizophrenia Research</i> , 2014 , 159, 253	3.6
16	Response to the letters of Dr Amos and Dr Preti and colleagues. <i>Schizophrenia Research</i> , 2014 , 153, 237-9.6	3.6
15	Is it appropriate to treat people at high-risk of psychosis before first onset - yes. <i>Medical Journal of Australia</i> , 2012 , 196, 556	4
14	The EPOS prediction model improves ability to predict transition to first episode psychosis in individuals at high risk. <i>Evidence-Based Mental Health</i> , 2010 , 13, 77	11.1
13	EVIDENCE FOR ALTERED ASYMMETRY OF FRONTAL CORTEX T2 RELAXATION TIME IN PATIENTS AT CLINICAL HIGH-RISK FOR PSYCHOSIS. <i>Schizophrenia Research</i> , 2010 , 117, 534	3.6
12	Diagnostic markers for schizophrenia: do we actually know what we're looking for?. <i>World Psychiatry</i> , 2011 , 10, 33-4	14.4
11	Evidencia de neuroimagen y de tratamiento respecto a los estadios clínicos en los trastornos psicóticos: del estado mental de riesgo a la esquizofrenia crónica. <i>Psiquiatria Biologica</i> , 2011 , 18, 117-123	0.2
10	Commentary on "Should the diagnostic boundaries of schizophrenia be expanded?". <i>Cognitive Neuropsychiatry</i> , 2011 , 16, 107-12	2
9	At-risk mental state and prediction83-106	
8	P.1.e.016 Anatomical abnormalities of the anterior cingulate cortex before the onset of psychosis. <i>European Neuropsychopharmacology</i> , 2008 , 18, S268-S269	1.2
7	S.22.01 Brain changes during the development of psychosis. <i>European Neuropsychopharmacology</i> , 2007 , 17, S209	1.2
6	The Management of Early Psychosis 2005 , 51-109	
5	Role of Diet in Depression in Young People: Recommendations for Implementation in Practice. <i>Psychiatric Annals</i> , 2022 , 52, 56-61	0.5

4	Physical health promotion in people with schizophrenia: why we should consider the ultra high-risk state. <i>Acta Psychiatrica Scandinavica</i> , 2016 , 133, 166-167	6.5
3	Investigation of structural brain correlates of neurological soft signs in individuals at ultra-high risk for psychosis. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2021 , 271, 1475-1485	5.1
2	Staff perspectives on the barriers and facilitators to exercise implementation in inpatient mental health services: A qualitative study. <i>Mental Health and Physical Activity</i> , 2022 , 22, 100452	5
1	Pineal morphology of the clinical high-risk state for psychosis and different psychotic disorders.. <i>Schizophrenia Research</i> , 2022 , 244, 1-7	3.6