

Yi-Feng Xu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2587585/publications.pdf>

Version: 2024-02-01

165
papers

9,198
citations

172207

29
h-index

48187

88
g-index

173
all docs

173
docs citations

173
times ranked

14519
citing authors

#	ARTICLE	IF	CITATIONS
1	A nationwide survey of psychological distress among Chinese people in the COVID-19 epidemic: implications and policy recommendations. <i>Annals of General Psychiatry</i> , 2020, 33, e100213.	1.1	2,951
2	Prevalence of mental disorders in China: a cross-sectional epidemiological study. <i>Lancet Psychiatry</i> , 2019, 6, 211-224.	3.7	1,273
3	Patients with mental health disorders in the COVID-19 epidemic. <i>Lancet Psychiatry</i> , 2020, 7, e21.	3.7	1,053
4	Genome-wide association analysis identifies 30 new susceptibility loci for schizophrenia. <i>Nature Genetics</i> , 2017, 49, 1576-1583.	9.4	395
5	Prevalence of depressive disorders and treatment in China: a cross-sectional epidemiological study. <i>Lancet Psychiatry</i> , 2021, 8, 981-990.	3.7	264
6	Mental health system in China: history, recent service reform and future challenges. <i>World Psychiatry</i> , 2011, 10, 210-216.	4.8	232
7	Common variants on 8p12 and 1q24.2 confer risk of schizophrenia. <i>Nature Genetics</i> , 2011, 43, 1224-1227.	9.4	224
8	The MATRICS Consensus Cognitive Battery (MCCB): Co-norming and standardization in China. <i>Schizophrenia Research</i> , 2015, 169, 109-115.	1.1	176
9	Cerebral metabolism in major depressive disorder: a voxel-based meta-analysis of positron emission tomography studies. <i>BMC Psychiatry</i> , 2014, 14, 321.	1.1	170
10	Rethinking online mental health services in China during the COVID-19 epidemic. <i>Asian Journal of Psychiatry</i> , 2020, 50, 102015.	0.9	151
11	Positive association between SIAT8B and schizophrenia in the Chinese Han population. <i>Schizophrenia Research</i> , 2007, 90, 108-114.	1.1	87
12	MicroRNA-223 protects neonatal rat cardiomyocytes and H9c2 cells from hypoxia-induced apoptosis and excessive autophagy via the Akt/mTOR pathway by targeting PARP-1. <i>Journal of Molecular and Cellular Cardiology</i> , 2018, 118, 133-146.	0.9	82
13	A case-control study of the relationship between the metabotropic glutamate receptor 3 gene and schizophrenia in the Chinese population. <i>Schizophrenia Research</i> , 2005, 73, 21-26.	1.1	80
14	Abnormal white matter microstructure in drug-naive first episode schizophrenia patients before and after eight weeks of antipsychotic treatment. <i>Schizophrenia Research</i> , 2016, 172, 1-8.	1.1	75
15	Dopamine D4 receptor polymorphism modulates cue-elicited heroin craving in Chinese. <i>Psychopharmacology</i> , 2006, 186, 185-190.	1.5	70
16	A study of N-methyl-D-aspartate receptor gene (GRIN2B) variants as predictors of treatment-resistant major depression. <i>Psychopharmacology</i> , 2014, 231, 685-693.	1.5	65
17	Genome-wide Analysis of the Role of Copy Number Variation in Schizophrenia Risk in Chinese. <i>Biological Psychiatry</i> , 2016, 80, 331-337.	0.7	55
18	Comparisons of the Efficacy and Tolerability of Extended-Release Venlafaxine, Mirtazapine, and Paroxetine in Treatment-Resistant Depression. <i>Journal of Clinical Psychopharmacology</i> , 2010, 30, 357-364.	0.7	48

#	ARTICLE	IF	CITATIONS
19	A Pilot Study of the Efficacy and Safety of Paroxetine Augmented With Risperidone, Valproate, Buspirone, Trazodone, or Thyroid Hormone in Adult Chinese Patients With Treatment-Resistant Major Depression. <i>Journal of Clinical Psychopharmacology</i> , 2011, 31, 638-642.	0.7	47
20	Pharmacogenetic effects of dopamine transporter gene polymorphisms on response to chlorpromazine and clozapine and on extrapyramidal syndrome in schizophrenia. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2010, 34, 1026-1032.	2.5	46
21	Family-Based Association Study of Synapsin II and Schizophrenia. <i>American Journal of Human Genetics</i> , 2004, 75, 873-877.	2.6	44
22	Increased Cognition Connectivity Network in Major Depression Disorder: A fMRI Study. <i>Psychiatry Investigation</i> , 2015, 12, 227.	0.7	40
23	The China Mental Health Survey (CMHS): I. background, aims and measures. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2016, 51, 1559-1569.	1.6	40
24	No association between the promoter variants of tumor necrosis factor alpha (TNF- α) and schizophrenia in Chinese Han population. <i>Neuroscience Letters</i> , 2004, 366, 139-143.	1.0	39
25	Regional Abnormality of Grey Matter in Schizophrenia: Effect from the Illness or Treatment?. <i>PLoS ONE</i> , 2016, 11, e0147204.	1.1	37
26	Genetic Structure Adds Power to Detect Schizophrenia Susceptibility at SLIT3 in the Chinese Han Population. <i>Genome Research</i> , 2004, 14, 1345-1349.	2.4	36
27	The GSK3B gene confers risk for both major depressive disorder and schizophrenia in the Han Chinese population. <i>Journal of Affective Disorders</i> , 2015, 185, 149-155.	2.0	34
28	The China Mental Health Survey: II. Design and field procedures. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2016, 51, 1547-1557.	1.6	34
29	Glucose and Insulin-Related Traits, Type 2 Diabetes and Risk of Schizophrenia: A Mendelian Randomization Study. <i>EBioMedicine</i> , 2018, 34, 182-188.	2.7	34
30	Striatal dysfunction in patients with schizophrenia and their unaffected first-degree relatives. <i>Schizophrenia Research</i> , 2018, 195, 215-221.	1.1	33
31	Positive association between OLIG2 and schizophrenia in the Chinese Han population. <i>Human Genetics</i> , 2008, 122, 659-660.	1.8	30
32	A family-based association study of the MOC gene with schizophrenia in the Chinese population. <i>Schizophrenia Research</i> , 2005, 73, 275-280.	1.1	29
33	RGS4 polymorphisms and risk of schizophrenia: An association study in Han Chinese plus meta-analysis. <i>Neuroscience Letters</i> , 2006, 406, 122-127.	1.0	29
34	Chronic mild restraint stress rats decreased CMKLR1 expression in distinct brain region. <i>Neuroscience Letters</i> , 2012, 524, 25-29.	1.0	29
35	<p>Repetitive transcranial magnetic stimulation as an adjunctive treatment for negative symptoms and cognitive impairment in patients with schizophrenia: a randomized, double-blind, sham-controlled trial</p>. <i>Neuropsychiatric Disease and Treatment</i> , 2019, Volume 15, 1141-1150.	1.0	27
36	Macrophage migration inhibitory factor mediates metabolic dysfunction induced by atypical antipsychotic therapy. <i>Journal of Clinical Investigation</i> , 2018, 128, 4997-5007.	3.9	27

#	ARTICLE	IF	CITATIONS
37	Hyperactivity within an extensive cortical distribution associated with excessive sensitivity in error processing in unmedicated depression: A combined event-related potential and sLORETA study. <i>International Journal of Psychophysiology</i> , 2013, 90, 282-289.	0.5	26
38	Positive association between ALDH1A2 and schizophrenia in the Chinese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2009, 33, 1491-1495.	2.5	25
39	The Ser9Gly polymorphism of the dopamine D3 receptor gene and risk of schizophrenia: An association study and a large meta-analysis. <i>Schizophrenia Research</i> , 2008, 101, 26-35.	1.1	22
40	Association between a COMT polymorphism and clinical response to risperidone treatment. <i>Psychiatric Genetics</i> , 2012, 22, 298-299.	0.6	22
41	Prospective memory performance in patients with drug-naïve, first-episode psychosis. <i>Schizophrenia Research</i> , 2013, 143, 285-290.	1.1	22
42	What is the optimal neuropsychological test battery for schizophrenia in China?. <i>Schizophrenia Research</i> , 2019, 208, 317-323.	1.1	22
43	Prevalence and dynamic features of psychological issues among Chinese healthcare workers during the COVID-19 pandemic: a systematic review and cumulative meta-analysis. <i>Annals of General Psychiatry</i> , 2021, 34, e100344.	1.1	22
44	A pharmacogenetic study of risperidone on histamine H3 receptor gene (<i>HRH3</i>) in Chinese Han schizophrenia patients. <i>Journal of Psychopharmacology</i> , 2012, 26, 813-818.	2.0	19
45	Histamine H4 Receptor Polymorphism. <i>Journal of Clinical Psychopharmacology</i> , 2013, 33, 221-225.	0.7	19
46	Association between SREBF2 gene polymorphisms and metabolic syndrome in clozapine-treated patients with schizophrenia. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2015, 56, 136-141.	2.5	19
47	Network pharmacology-based identification for therapeutic mechanism of Ling-Gui-Zhu-Gan decoction in the metabolic syndrome induced by antipsychotic drugs. <i>Computers in Biology and Medicine</i> , 2019, 110, 1-7.	3.9	19
48	Challenges and opportunities in mental health services during the COVID-19 pandemic. <i>Annals of General Psychiatry</i> , 2020, 33, e100275.	1.1	19
49	Quality of life in outpatients with depression in China. <i>Journal of Affective Disorders</i> , 2013, 150, 513-521.	2.0	18
50	–Individual Perceived Stress Mediates Psychological Distress in Medical Workers During COVID-19 Epidemic Outbreak in Wuhan–. <i>Neuropsychiatric Disease and Treatment</i> , 2020, Volume 16, 2529-2537.	1.0	18
51	Family-based association study between brain-derived neurotrophic factor gene and major depressive disorder of Chinese descent. <i>Psychiatry Research</i> , 2009, 169, 169-172.	1.7	17
52	Polymorphisms of XRCC4 are involved in reduced colorectal cancer risk in Chinese schizophrenia patients. <i>BMC Cancer</i> , 2010, 10, 523.	1.1	17
53	The NVL gene confers risk for both major depressive disorder and schizophrenia in the Han Chinese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2015, 62, 7-13.	2.5	17
54	Dysregulated 14-3-3 Family in Peripheral Blood Leukocytes of Patients with Schizophrenia. <i>Scientific Reports</i> , 2016, 6, 23791.	1.6	17

#	ARTICLE	IF	CITATIONS
55	Effect of Electroconvulsive Therapy on Medial Prefrontal \hat{I}^3 -Aminobutyric Acid Among Schizophrenia Patients. <i>Journal of ECT</i> , 2018, 34, 227-232.	0.3	17
56	The comorbidity of mental and physical disorders with self-reported chronic back or neck pain: Results from the China Mental Health Survey. <i>Journal of Affective Disorders</i> , 2020, 260, 334-341.	2.0	17
57	A family-based association study of kinesin heavy chain member 2 gene (KIF2) and schizophrenia. <i>Neuroscience Letters</i> , 2006, 407, 151-155.	1.0	16
58	A new method for identifying causal genes of schizophrenia and anti-tuberculosis drug-induced hepatotoxicity. <i>Scientific Reports</i> , 2016, 6, 32571.	1.6	16
59	Meta-Analysis-Based Preliminary Exploration of the Connection between ATDILI and Schizophrenia by GSTM1/T1 Gene Polymorphisms. <i>PLoS ONE</i> , 2015, 10, e0128643.	1.1	16
60	No significant association between the genetic polymorphisms in the GSK-3 \hat{I}^2 gene and schizophrenia in the Chinese population. <i>Journal of Psychiatric Research</i> , 2008, 42, 365-370.	1.5	15
61	Haplotype analysis confirms association of the serotonin transporter (5-HTT) gene with schizophrenia in the Han Chinese population. <i>Neuroscience Letters</i> , 2009, 453, 210-213.	1.0	15
62	<i>HTR2C</i> promoter polymorphisms are associated with risperidone efficacy in Chinese female patients. <i>Pharmacogenomics</i> , 2010, 11, 685-692.	0.6	15
63	Comparison of Chinese and international psychiatrists' views on classification of mental disorders. <i>Asia-Pacific Psychiatry</i> , 2014, 6, 267-273.	1.2	15
64	Neurochemical and brain functional changes in the ventromedial prefrontal cortex of first-episode psychosis patients: A combined functional magnetic resonance imaging \hat{I}^2 proton magnetic resonance spectroscopy study. <i>Australian and New Zealand Journal of Psychiatry</i> , 2020, 54, 519-527.	1.3	15
65	Wnt Signaling Pathway in Schizophrenia. <i>CNS and Neurological Disorders - Drug Targets</i> , 2014, 13, 755-764.	0.8	15
66	An association study between PPP1R1B gene and schizophrenia in the Chinese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2007, 31, 1303-1306.	2.5	14
67	No association between EGR gene family polymorphisms and schizophrenia in the Chinese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2010, 34, 506-509.	2.5	14
68	A study of negative life events driven depressive symptoms and academic engagement in Chinese college students. <i>Scientific Reports</i> , 2021, 11, 17160.	1.6	14
69	Polymorphisms and Haplotypes in theYWHAEGene Increase Susceptibility to Bipolar Disorder in Chinese Han Population. <i>Journal of Clinical Psychiatry</i> , 2012, 73, e1276-e1282.	1.1	14
70	Multicenter randomized controlled trial of bifrontal, bitemporal, and right unilateral electroconvulsive therapy in major depressive disorder. <i>Psychiatry and Clinical Neurosciences</i> , 2019, 73, 636-641.	1.0	13
71	Mitigating mental health consequences during the <i>COVID</i> \hat{I}^9 outbreak: Lessons from China. <i>Psychiatry and Clinical Neurosciences</i> , 2020, 74, 407-408.	1.0	13
72	The Prevalence of Psychological Status During the COVID-19 Epidemic in China: A Systemic Review and Meta-Analysis. <i>Frontiers in Psychology</i> , 2021, 12, 614964.	1.1	13

#	ARTICLE	IF	CITATIONS
73	Comparison of the density of gamma-aminobutyric acid in the ventromedial prefrontal cortex of patients with first-episode psychosis and healthy controls. <i>Shanghai Archives of Psychiatry</i> , 2015, 27, 341-7.	0.7	13
74	No association between the genetic polymorphisms within RTN4 and schizophrenia in the Chinese population. <i>Neuroscience Letters</i> , 2004, 365, 23-27.	1.0	12
75	Association between <i>SCAP</i> and <i>SREBF1</i> gene polymorphisms and metabolic syndrome in schizophrenia patients treated with atypical antipsychotics. <i>World Journal of Biological Psychiatry</i> , 2016, 17, 467-474.	1.3	12
76	Impaired cue identification and intention retrieval underlie prospective memory deficits in patients with first-episode schizophrenia. <i>Australian and New Zealand Journal of Psychiatry</i> , 2017, 51, 270-277.	1.3	12
77	Influence and interaction of genetic, cognitive, neuroendocrine and personalistic markers to antidepressant response in Chinese patients with major depression. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2021, 104, 110036.	2.5	12
78	Association study of 5-HT1A, 5-HT2A polymorphisms with schizophrenia and major depressive disorder in the Han Chinese population. <i>Neuroscience Letters</i> , 2016, 635, 39-43.	1.0	11
79	A case-control association study between the CYP3A4 and CYP3A5 genes and schizophrenia in the Chinese Han population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2009, 33, 1200-1204.	2.5	10
80	Identification of the <i>N</i> -acylsphingosine amidohydrolase 1 gene (<i>ASAH1</i>) for susceptibility to schizophrenia in a Han Chinese population. <i>World Journal of Biological Psychiatry</i> , 2012, 13, 106-113.	1.3	10
81	Effect of SOX10 gene polymorphism on early onset schizophrenia in Chinese Han population. <i>Neuroscience Letters</i> , 2012, 521, 93-97.	1.0	10
82	Common Variants in the TPH2 Promoter Confer Susceptibility to Paranoid Schizophrenia. <i>Journal of Molecular Neuroscience</i> , 2012, 47, 465-469.	1.1	10
83	A new risk locus in the ZEB2 gene for schizophrenia in the Han Chinese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2016, 66, 97-103.	2.5	10
84	Genome-wide significant, replicated and functional risk variants for Alzheimer's disease. <i>Journal of Neural Transmission</i> , 2017, 124, 1455-1471.	1.4	10
85	SNX29, a new susceptibility gene shared with major mental disorders in Han Chinese population. <i>World Journal of Biological Psychiatry</i> , 2021, 22, 526-534.	1.3	10
86	Role of rs454214 in Personality mediated Depression and Subjective Well-being. <i>Scientific Reports</i> , 2020, 10, 5702.	1.6	10
87	No genetic association between NCAM1 gene polymorphisms and schizophrenia in the Chinese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2008, 32, 1633-1636.	2.5	9
88	Genetic polymorphisms in the SCN8A gene are associated with suicidal behavior in psychiatric disorders in the Chinese population. <i>World Journal of Biological Psychiatry</i> , 2010, 11, 956-963.	1.3	9
89	SOX10 rs139883 Polymorphism Is Associated with the Age of Onset in Schizophrenia. <i>Journal of Molecular Neuroscience</i> , 2013, 50, 333-338.	1.1	9
90	The use of the SDQ with Chinese adolescents in the clinical context. <i>Psychiatry Research</i> , 2016, 246, 520-526.	1.7	9

#	ARTICLE	IF	CITATIONS
91	A study of single nucleotide polymorphisms of GRIN2B in schizophrenia from Chinese Han population. <i>Neuroscience Letters</i> , 2016, 630, 132-135.	1.0	9
92	Association between the variability of the <i>ABCA13</i> gene and the risk of major depressive disorder and schizophrenia in the Han Chinese population. <i>World Journal of Biological Psychiatry</i> , 2017, 18, 550-556.	1.3	9
93	Prediction of adolescent subjective well-being: A machine learning approach. <i>Annals of General Psychiatry</i> , 2019, 32, e100096.	1.1	9
94	An independent, replicable, functional and significant risk variant block at intron 3 of <i>CACNA1C</i> for schizophrenia. <i>Australian and New Zealand Journal of Psychiatry</i> , 2022, 56, 385-397.	1.3	9
95	Network pharmacology-based exploration of therapeutic mechanism of Liu-Yu-Tang in atypical antipsychotic drug-induced metabolic syndrome. <i>Computers in Biology and Medicine</i> , 2021, 134, 104452.	3.9	9
96	No genetic association between polymorphisms in the kainate-type glutamate receptor gene, <i>GRIK4</i> , and schizophrenia in the Chinese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2008, 32, 876-880.	2.5	8
97	Association study of <i>GRM7</i> polymorphisms and schizophrenia in the Chinese Han population. <i>Neuroscience Letters</i> , 2015, 604, 109-112.	1.0	8
98	Association study of dopamine receptor genes polymorphisms with the risk of schizophrenia in the Han Chinese population. <i>Psychiatry Research</i> , 2016, 245, 361-364.	1.7	8
99	No association found between the promoter variants of <i>ADRA1A</i> and schizophrenia in the Chinese population. <i>Journal of Psychiatric Research</i> , 2008, 42, 384-388.	1.5	7
100	Deficient inhibition of return in chronic but not first-episode patients with schizophrenia. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2010, 34, 961-967.	2.5	7
101	Ethics, patient rights and staff attitudes in Shanghai's psychiatric hospitals. <i>BMC Medical Ethics</i> , 2012, 13, 8.	1.0	7
102	The rights of psychiatric patients in China: A survey of medical staff and consumers' attitudes toward patient participation in clinical trials. <i>Social Science and Medicine</i> , 2012, 75, 823-827.	1.8	7
103	Dysfunction of Cognition Patterns Measured by MATRICS Consensus Cognitive Battery (MCCB) among First Episode Schizophrenia Patients and Their Biological Parents. <i>Shanghai Archives of Psychiatry</i> , 2017, 29, 154-160.	0.7	7
104	Does practice make perfect? Results from a Chinese feasibility study of cognitive remediation in schizophrenia. <i>Neuropsychological Rehabilitation</i> , 2013, 23, 580-596.	1.0	6
105	Down-regulation of <i>PRKCB1</i> expression in Han Chinese patients with subsyndromal symptomatic depression. <i>Journal of Psychiatric Research</i> , 2015, 69, 1-6.	1.5	6
106	Common variants in <i>SLC6A2</i> , <i>SLC6A3</i> , <i>DRD2</i> , and major depressive disorder. <i>Psychiatric Genetics</i> , 2017, 27, 103-104.	0.6	6
107	<i>CYP1A2</i> Genetic Polymorphism Is Associated With Treatment Remission to Antidepressant Venlafaxine in Han Chinese Population. <i>Clinical Neuropharmacology</i> , 2019, 42, 32-36.	0.2	6
108	Association study between <i>LEPR</i> , <i>MC4R</i> polymorphisms and overweight/obesity in Chinese Han adolescents. <i>Gene</i> , 2019, 692, 54-59.	1.0	6

#	ARTICLE	IF	CITATIONS
109	Cigarette smoking and schizophrenia: Mendelian randomisation study. <i>British Journal of Psychiatry</i> , 2021, 218, 98-103.	1.7	6
110	Sp1 Targeted PARP1 Inhibition Protects Cardiomyocytes From Myocardial Ischemiaâ€“Reperfusion Injury via Downregulation of Autophagy. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 621906.	1.8	6
111	An Evaluation of the Shanghai Mental Health Service Schizophrenia Rehabilitation Program. <i>American Journal of Psychiatric Rehabilitation</i> , 2011, 14, 287-306.	0.7	5
112	Ziprasidone, haloperidol and clonazepam intramuscular administration in the treatment of agitation symptoms in Chinese patients with schizophrenia: A network meta-analysis. <i>Annals of General Psychiatry</i> , 2018, 31, e000016.	1.1	5
113	Mental Health Service Challenges during the Early Stage of the COVID-19 Pandemic: Experience and Best Practices from China. <i>Canadian Journal of Psychiatry</i> , 2020, 66, 070674372097225.	0.9	5
114	Genetic risk of clozapine-induced leukopenia and neutropenia: a genome-wide association study. <i>Translational Psychiatry</i> , 2021, 11, 343.	2.4	5
115	Urotensin II Induces Cardiac Fibrosis through the TGF-Î²2/Smad Signaling Pathway during the Development of Cardiac Hypertrophy. <i>International Heart Journal</i> , 2021, 62, 1135-1144.	0.5	5
116	The Association of Depressive Symptoms with Disability among Adults in China. <i>Journal of Affective Disorders</i> , 2021, 296, 189-197.	2.0	5
117	Assessment of a six-week computer-based remediation program for social cognition in chronic schizophrenia. <i>Shanghai Archives of Psychiatry</i> , 2015, 27, 296-306.	0.7	5
118	No association found between the promoter variations of QKI and schizophrenia in the Chinese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2009, 33, 33-36.	2.5	4
119	No association between the KCNH1, KCNJ10 and KCNN3 genes and schizophrenia in the Han Chinese population. <i>Neuroscience Letters</i> , 2011, 487, 61-65.	1.0	4
120	Evaluations of treatment efficacy of depression from perspective of both patientsâ€™ symptoms and general sense of mental health and wellbeing: A large scale, multi-centered, longitudinal study in China. <i>Psychiatry Research</i> , 2016, 241, 55-60.	1.7	4
121	Analysis of association between common variants in the <i>SLCO6A1</i> gene with schizophrenia, bipolar disorder and major depressive disorder in the Han Chinese population. <i>World Journal of Biological Psychiatry</i> , 2016, 17, 140-146.	1.3	4
122	Association study of GRM7 polymorphisms with major depressive disorder in the Chinese Han population. <i>Psychiatric Genetics</i> , 2017, 27, 78-79.	0.6	4
123	Common variants in GRIK4 and major depressive disorder: An association study in the Chinese Han population. <i>Neuroscience Letters</i> , 2017, 653, 239-243.	1.0	4
124	HTR1A and HTR2A variants may not predict venlafaxine treatment response in China Han population with major depressive disorder. <i>Psychiatry Research</i> , 2018, 270, 1179-1180.	1.7	4
125	Genetic association between CELF4 rs1557341 polymorphism and neuroticism in Chinese Han population. <i>Psychiatry Research</i> , 2019, 279, 138-139.	1.7	4
126	Barriers and facilitators to implementing measurement-based care for depression in Shanghai, China: a situational analysis. <i>BMC Psychiatry</i> , 2021, 21, 430.	1.1	4

#	ARTICLE	IF	CITATIONS
127	Efficacy of Atypical Antipsychotics in the Management of Acute Agitation and Aggression in Hospitalized Patients with Schizophrenia or Bipolar Disorder: Results from a Systematic Review. Shanghai Archives of Psychiatry, 2016, 28, 241-252.	0.7	4
128	Efficacy of Artificial Intelligence-Assisted Psychotherapy in Patients With Anxiety Disorders: A Prospective, National Multicenter Randomized Controlled Trial Protocol. Frontiers in Psychiatry, 2021, 12, 799917.	1.3	4
129	Teachers of Psychiatry meeting in Shanghai: A leadership training course. Asia-Pacific Psychiatry, 2012, 4, 87-89.	1.2	3
130	Association study of APC polymorphisms with colorectal cancer in Han Chinese. Clinical Biochemistry, 2012, 45, 1669-1672.	0.8	3
131	Efficacy of HuaDan Anshen Mistura for treating insomnia: a randomized, double-blind, placebo-controlled, multi-center clinical trial. Journal of Traditional Chinese Medicine = Chung I Tsa Chih Ying Wen Pan / Sponsored By All-China Association of Traditional Chinese Medicine, Academy of Traditional Chinese Medicine. 2013, 33, 423-427.	0.4	3
132	Association study on the DLG4 gene and schizophrenia in the Chinese Han population. Psychiatric Genetics, 2013, 23, 247-250.	0.6	3
133	Improvement in social and cognitive functioning associated with paliperidone extended-release treatment in patients with schizophrenia: a 24-week, single arm, open-label study. Neuropsychiatric Disease and Treatment, 2016, Volume 12, 2095-2104.	1.0	3
134	Common variants in <i>QPCT</i> gene confer risk of schizophrenia in the Han Chinese population. American Journal of Medical Genetics Part B: Neuropsychiatric Genetics, 2016, 171, 237-242.	1.1	3
135	Role played by the SP4 gene in schizophrenia and major depressive disorder in the Han Chinese population. British Journal of Psychiatry, 2016, 208, 441-445.	1.7	3
136	No association of GRIK4 polymorphisms with schizophrenia in the Chinese Han population. Psychiatric Genetics, 2017, 27, 159-160.	0.6	3
137	Should repetitive Transcranial Magnetic Stimulation (rTMS) be considered an effective adjunctive treatment for auditory hallucinations in patients with schizophrenia?. Shanghai Archives of Psychiatry, 2013, 25, 254-5.	0.7	3
138	Impact of OXTR Polymorphisms on Subjective Well-Being: The Intermediary Role of Attributional Style. Frontiers in Genetics, 2021, 12, 763628.	1.1	3
139	Association between the volume of subregions of the amygdala and major depression with suicidal thoughts and anxiety in a Chinese cohort. Journal of Affective Disorders, 2022, 312, 39-45.	2.0	3
140	Association study of NOS1 gene polymorphisms with the risk of schizophrenia in Chinese Han origin. Psychiatry Research, 2016, 246, 844-845.	1.7	2
141	No association between SLC6A2, SLC6A3, DRD2 polymorphisms and schizophrenia in the Han Chinese population. Psychiatry Research, 2017, 253, 398-400.	1.7	2
142	Association study of the GLRX5 rs1007814 polymorphism with schizophrenia in the Han Chinese population. Psychiatric Genetics, 2017, 27, 76-77.	0.6	2
143	Sequential Multiple-Assignment Randomized Trials to Compare Antipsychotic Treatments (SMART-CAT) in first-episode schizophrenia patients: Rationale and trial design. Schizophrenia Research, 2021, 230, 87-94.	1.1	2
144	Interaction of CEND1 gene and life events in susceptibility to depressive symptoms in Chinese Han college students. Journal of Affective Disorders, 2021, 278, 570-575.	2.0	2

#	ARTICLE	IF	CITATIONS
145	Research progress in China on the assessment of cognitive function in schizophrenia. Shanghai Archives of Psychiatry, 2013, 25, 266-75.	0.7	2
146	Schizophrenia is not associated with the ERBB3 gene in a Han Chinese population sample: results from case-control and family-based studies. Genetics and Molecular Biology, 2009, 32, 729-730.	0.6	1
147	Social support and clinical depression in China. Asia-Pacific Psychiatry, 2012, 4, 195-200.	1.2	1
148	Sleep problems among clinically depressed adults in China. Journal of Mental Health, 2015, 24, 43-47.	1.0	1
149	No association of GRIA1 polymorphisms with schizophrenia in the Chinese Han population. Psychiatric Genetics, 2016, 26, 97-98.	0.6	1
150	New journey, new mission: bolstering international communication about mental health. Annals of General Psychiatry, 2018, 31, e100002.	1.1	1
151	Improvement of the Diagnosis of Left Ventricular Noncompaction Cardiomyopathy After Analyzing Both Advantages and Disadvantages of Echocardiography and CMRI. Progress in Cardiovascular Diseases, 2018, 61, 491-493.	1.6	1
152	Comparison of response to Chinese and Western videos of mental-health-related emotions in a representative Chinese sample. PeerJ, 2021, 9, e10440.	0.9	1
153	Middle temporal corpus callosum impairment as a predictor of eight-week treatment outcome of drug-naïve first-episode psychosis patients: A pilot longitudinal study. Schizophrenia Research, 2021, 232, 95-97.	1.1	1
154	MicroRNA Microarray Analysis Combined with Interaction Network Analysis to Investigate the Influence of Clozapine to Metabolic Syndrome. International Journal of Pharmacology, 2013, 9, 366-372.	0.1	1
155	Neural Correlates of the Preserved Inhibition of Return in Schizophrenia. PLoS ONE, 2015, 10, e0119521.	1.1	1
156	Research Progress in Biological Studies of Schizophrenia in China in 2017. Shanghai Archives of Psychiatry, 2018, 30, 147-153.	0.7	1
157	NAPG may not be a common risk gene shared by bipolar disorder and schizophrenia in Chinese population. Psychiatric Genetics, 2012, 22, 107-108.	0.6	0
158	The promoter polymorphisms in HTR2A gene associated with schizophrenia in Chinese of Han ethnicity. Psychiatry Research, 2018, 262, 636-637.	1.7	0
159	A case-control study of GRIN2B polymorphisms and major depressive disorder in the Chinese Han population. Psychiatry Research, 2018, 262, 626-627.	1.7	0
160	No Association Between SLC6A4 Gene Polymorphisms With Treatment Remission to Venlafaxine in Han Chinese Depressive Patients. Clinical Neuropharmacology, 2021, 44, 53-56.	0.2	0
161	The Epidemiology of Depressive Disorders and Service Use in China: Findings from the China Mental Health Survey. SSRN Electronic Journal, 0, , .	0.4	0
162	Case report of Folie À Trois. Shanghai Archives of Psychiatry, 2013, 25, 52-4.	0.7	0

#	ARTICLE	IF	CITATIONS
163	Advances in neuroimaging research of schizophrenia in China. Shanghai Archives of Psychiatry, 2014, 26, 181-93.	0.7	0
164	Symptom severity is more closely associated with social functioning status in inpatients with schizophrenia than cognitive deficits. Shanghai Archives of Psychiatry, 2012, 24, 83-90.	0.7	0
165	A novel heterozygous missense variant of the ARID4A gene identified in Han Chinese families with schizophrenia-diagnosed siblings that interferes with DNA-binding activity. Molecular Psychiatry, 2022, , .	4.1	0