

# Yi-Feng Xu

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2587585/publications.pdf>

Version: 2024-02-01

165  
papers

9,198  
citations

172207

29  
h-index

48187

88  
g-index

173  
all docs

173  
docs citations

173  
times ranked

14519  
citing authors

#	ARTICLE	IF	CITATIONS
1	An independent, replicable, functional and significant risk variant block at intron 3 of <i>CACNA1C</i> for schizophrenia. Australian and New Zealand Journal of Psychiatry, 2022, 56, 385-397.	1.3	9
2	A novel heterozygous missense variant of the ARID4A gene identified in Han Chinese families with schizophrenia-diagnosed siblings that interferes with DNA-binding activity. Molecular Psychiatry, 2022, , .	4.1	0
3	Association between the volume of subregions of the amygdala and major depression with suicidal thoughts and anxiety in a Chinese cohort. Journal of Affective Disorders, 2022, 312, 39-45.	2.0	3
4	Sequential Multiple-Assignment Randomized Trials to Compare Antipsychotic Treatments (SMART-CAT) in first-episode schizophrenia patients: Rationale and trial design. Schizophrenia Research, 2021, 230, 87-94.	1.1	2
5	SNX29, a new susceptibility gene shared with major mental disorders in Han Chinese population. World Journal of Biological Psychiatry, 2021, 22, 526-534.	1.3	10
6	Cigarette smoking and schizophrenia: Mendelian randomisation study. British Journal of Psychiatry, 2021, 218, 98-103.	1.7	6
7	Influence and interaction of genetic, cognitive, neuroendocrine and personalistic markers to antidepressant response in Chinese patients with major depression. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2021, 104, 110036.	2.5	12
8	Interaction of CEND1 gene and life events in susceptibility to depressive symptoms in Chinese Han college students. Journal of Affective Disorders, 2021, 278, 570-575.	2.0	2
9	No Association Between SLC6A4 Gene Polymorphisms With Treatment Remission to Venlafaxine in Han Chinese Depressive Patients. Clinical Neuropharmacology, 2021, 44, 53-56.	0.2	0
10	Comparison of response to Chinese and Western videos of mental-health-related emotions in a representative Chinese sample. PeerJ, 2021, 9, e10440.	0.9	1
11	The Prevalence of Psychological Status During the COVID-19 Epidemic in China: A Systemic Review and Meta-Analysis. Frontiers in Psychology, 2021, 12, 614964.	1.1	13
12	Prevalence and dynamic features of psychological issues among Chinese healthcare workers during the COVID-19 pandemic: a systematic review and cumulative meta-analysis. Annals of General Psychiatry, 2021, 34, e100344.	1.1	22
13	Sp1 Targeted PARP1 Inhibition Protects Cardiomyocytes From Myocardial Ischemia-Induced Reperfusion Injury via Downregulation of Autophagy. Frontiers in Cell and Developmental Biology, 2021, 9, 621906.	1.8	6
14	Middle temporal corpus callosum impairment as a predictor of eight-week treatment outcome of drug-naïve first-episode psychosis patients: A pilot longitudinal study. Schizophrenia Research, 2021, 232, 95-97.	1.1	1
15	Genetic risk of clozapine-induced leukopenia and neutropenia: a genome-wide association study. Translational Psychiatry, 2021, 11, 343.	2.4	5
16	Network pharmacology-based exploration of therapeutic mechanism of Liu-Yu-Tang in atypical antipsychotic drug-induced metabolic syndrome. Computers in Biology and Medicine, 2021, 134, 104452.	3.9	9
17	A study of negative life events driven depressive symptoms and academic engagement in Chinese college students. Scientific Reports, 2021, 11, 17160.	1.6	14
18	Prevalence of depressive disorders and treatment in China: a cross-sectional epidemiological study. Lancet Psychiatry, the, 2021, 8, 981-990.	3.7	264

#	ARTICLE	IF	CITATIONS
19	Urotensin II Induces Cardiac Fibrosis through the TGF- $\beta$ 2/Smad Signaling Pathway during the Development of Cardiac Hypertrophy. <i>International Heart Journal</i> , 2021, 62, 1135-1144.	0.5	5
20	The Association of Depressive Symptoms with Disability among Adults in China. <i>Journal of Affective Disorders</i> , 2021, 296, 189-197.	2.0	5
21	Barriers and facilitators to implementing measurement-based care for depression in Shanghai, China: a situational analysis. <i>BMC Psychiatry</i> , 2021, 21, 430.	1.1	4
22	Efficacy of Artificial Intelligence-Assisted Psychotherapy in Patients With Anxiety Disorders: A Prospective, National Multicenter Randomized Controlled Trial Protocol. <i>Frontiers in Psychiatry</i> , 2021, 12, 799917.	1.3	4
23	Impact of OXTR Polymorphisms on Subjective Well-Being: The Intermediary Role of Attributional Style. <i>Frontiers in Genetics</i> , 2021, 12, 763628.	1.1	3
24	The comorbidity of mental and physical disorders with self-reported chronic back or neck pain: Results from the China Mental Health Survey. <i>Journal of Affective Disorders</i> , 2020, 260, 334-341.	2.0	17
25	Mental Health Service Challenges during the Early Stage of the COVID-19 Pandemic: Experience and Best Practices from China. <i>Canadian Journal of Psychiatry</i> , 2020, 66, 070674372097225.	0.9	5
26	Challenges and opportunities in mental health services during the COVID-19 pandemic. <i>Annals of General Psychiatry</i> , 2020, 33, e100275.	1.1	19
27	Individual Perceived Stress Mediates Psychological Distress in Medical Workers During COVID-19 Epidemic Outbreak in Wuhan. <i>Neuropsychiatric Disease and Treatment</i> , 2020, Volume 16, 2529-2537.	1.0	18
28	Mitigating mental health consequences during the COVID-19 outbreak: Lessons from China. <i>Psychiatry and Clinical Neurosciences</i> , 2020, 74, 407-408.	1.0	13
29	Patients with mental health disorders in the COVID-19 epidemic. <i>Lancet Psychiatry</i> , 2020, 7, e21.	3.7	1,053
30	A nationwide survey of psychological distress among Chinese people in the COVID-19 epidemic: implications and policy recommendations. <i>Annals of General Psychiatry</i> , 2020, 33, e100213.	1.1	2,951
31	Neurochemical and brain functional changes in the ventromedial prefrontal cortex of first-episode psychosis patients: A combined functional magnetic resonance imaging-proton magnetic resonance spectroscopy study. <i>Australian and New Zealand Journal of Psychiatry</i> , 2020, 54, 519-527.	1.3	15
32	Rethinking online mental health services in China during the COVID-19 epidemic. <i>Asian Journal of Psychiatry</i> , 2020, 50, 102015.	0.9	151
33	Role of rs454214 in Personality mediated Depression and Subjective Well-being. <i>Scientific Reports</i> , 2020, 10, 5702.	1.6	10
34	Multicenter randomized controlled trial of bifrontal, bitemporal, and right unilateral electroconvulsive therapy in major depressive disorder. <i>Psychiatry and Clinical Neurosciences</i> , 2019, 73, 636-641.	1.0	13
35	Genetic association between CELF4 rs1557341 polymorphism and neuroticism in Chinese Han population. <i>Psychiatry Research</i> , 2019, 279, 138-139.	1.7	4
36	Network pharmacology-based identification for therapeutic mechanism of Ling-Gui-Zhu-Gan decoction in the metabolic syndrome induced by antipsychotic drugs. <i>Computers in Biology and Medicine</i> , 2019, 110, 1-7.	3.9	19

#	ARTICLE	IF	CITATIONS
37	&lt;p&gt;Repetitive transcranial magnetic stimulation as an adjunctive treatment for negative symptoms and cognitive impairment in patients with schizophrenia: a randomized, double-blind, sham-controlled trial&lt;p&gt;. Neuropsychiatric Disease and Treatment, 2019, Volume 15, 1141-1150.	1.0	27
38	Prevalence of mental disorders in China: a cross-sectional epidemiological study. Lancet Psychiatry, the, 2019, 6, 211-224.	3.7	1,273
39	What is the optimal neuropsychological test battery for schizophrenia in China?. Schizophrenia Research, 2019, 208, 317-323.	1.1	22
40	CYP1A2 Genetic Polymorphism Is Associated With Treatment Remission to Antidepressant Venlafaxine in Han Chinese Population. Clinical Neuropharmacology, 2019, 42, 32-36.	0.2	6
41	Prediction of adolescent subjective well-being: A machine learning approach. Annals of General Psychiatry, 2019, 32, e100096.	1.1	9
42	Association study between LEPR, MC4R polymorphisms and overweight/obesity in Chinese Han adolescents. Gene, 2019, 692, 54-59.	1.0	6
43	The promoter polymorphisms in HTR2A gene associated with schizophrenia in Chinese of Han ethnicity. Psychiatry Research, 2018, 262, 636-637.	1.7	0
44	MicroRNA-223 protects neonatal rat cardiomyocytes and H9c2 cells from hypoxia-induced apoptosis and excessive autophagy via the Akt/mTOR pathway by targeting PARP-1. Journal of Molecular and Cellular Cardiology, 2018, 118, 133-146.	0.9	82
45	Striatal dysfunction in patients with schizophrenia and their unaffected first-degree relatives. Schizophrenia Research, 2018, 195, 215-221.	1.1	33
46	A case-control study of GRIN2B polymorphisms and major depressive disorder in the Chinese Han population. Psychiatry Research, 2018, 262, 626-627.	1.7	0
47	New journey, new mission: bolstering international communication about mental health. Annals of General Psychiatry, 2018, 31, e100002.	1.1	1
48	Ziprasidone, haloperidol and clonazepam intramuscular administration in the treatment of agitation symptoms in Chinese patients with schizophrenia: A network meta-analysis. Annals of General Psychiatry, 2018, 31, e000016.	1.1	5
49	HTR1A and HTR2A variants may not predict venlafaxine treatment response in China Han population with major depressive disorder. Psychiatry Research, 2018, 270, 1179-1180.	1.7	4
50	Glucose and Insulin-Related Traits, Type 2 Diabetes and Risk of Schizophrenia: A Mendelian Randomization Study. EBioMedicine, 2018, 34, 182-188.	2.7	34
51	Improvement of the Diagnosis of Left Ventricular Noncompaction Cardiomyopathy After Analyzing Both Advantages and Disadvantages of Echocardiography and CMRI. Progress in Cardiovascular Diseases, 2018, 61, 491-493.	1.6	1
52	Effect of Electroconvulsive Therapy on Medial Prefrontal Î³-Aminobutyric Acid Among Schizophrenia Patients. Journal of ECT, 2018, 34, 227-232.	0.3	17
53	Macrophage migration inhibitory factor mediates metabolic dysfunction induced by atypical antipsychotic therapy. Journal of Clinical Investigation, 2018, 128, 4997-5007.	3.9	27
54	Research Progress in Biological Studies of Schizophrenia in China in 2017. Shanghai Archives of Psychiatry, 2018, 30, 147-153.	0.7	1

#	ARTICLE	IF	CITATIONS
55	Impaired cue identification and intention retrieval underlie prospective memory deficits in patients with first-episode schizophrenia. Australian and New Zealand Journal of Psychiatry, 2017, 51, 270-277.	1.3	12
56	No association between SLC6A2, SLC6A3, DRD2 polymorphisms and schizophrenia in the Han Chinese population. Psychiatry Research, 2017, 253, 398-400.	1.7	2
57	Association study of the GLRX5 rs1007814 polymorphism with schizophrenia in the Han Chinese population. Psychiatric Genetics, 2017, 27, 76-77.	0.6	2
58	Association study of GRM7 polymorphisms with major depressive disorder in the Chinese Han population. Psychiatric Genetics, 2017, 27, 78-79.	0.6	4
59	Common variants in SLC6A2, SLC6A3, DRD2, and major depressive disorder. Psychiatric Genetics, 2017, 27, 103-104.	0.6	6
60	Genome-wide association analysis identifies 30 new susceptibility loci for schizophrenia. Nature Genetics, 2017, 49, 1576-1583.	9.4	395
61	Genome-wide significant, replicated and functional risk variants for Alzheimer's disease. Journal of Neural Transmission, 2017, 124, 1455-1471.	1.4	10
62	Common variants in GRIK4 and major depressive disorder: An association study in the Chinese Han population. Neuroscience Letters, 2017, 653, 239-243.	1.0	4
63	No association of GRIK4 polymorphisms with schizophrenia in the Chinese Han population. Psychiatric Genetics, 2017, 27, 159-160.	0.6	3
64	Association between the variability of the ABCA13 gene and the risk of major depressive disorder and schizophrenia in the Han Chinese population. World Journal of Biological Psychiatry, 2017, 18, 550-556.	1.3	9
65	Dysfunction of Cognition Patterns Measured by MATRICS Consensus Cognitive Battery (MCCB) among First Episode Schizophrenia Patients and Their Biological Parents. Shanghai Archives of Psychiatry, 2017, 29, 154-160.	0.7	7
66	Improvement in social and cognitive functioning associated with paliperidone extended-release treatment in patients with schizophrenia: a 24-week, single arm, open-label study. Neuropsychiatric Disease and Treatment, 2016, Volume 12, 2095-2104.	1.0	3
67	Common variants in QPCT gene confer risk of schizophrenia in the Han Chinese population. American Journal of Medical Genetics Part B: Neuropsychiatric Genetics, 2016, 171, 237-242.	1.1	3
68	No association of GRIA1 polymorphisms with schizophrenia in the Chinese Han population. Psychiatric Genetics, 2016, 26, 97-98.	0.6	1
69	Association study of NOS1 gene polymorphisms with the risk of schizophrenia in Chinese Han origin. Psychiatry Research, 2016, 246, 844-845.	1.7	2
70	The use of the SDQ with Chinese adolescents in the clinical context. Psychiatry Research, 2016, 246, 520-526.	1.7	9
71	Association between SCAP and SREBF1 gene polymorphisms and metabolic syndrome in schizophrenia patients treated with atypical antipsychotics. World Journal of Biological Psychiatry, 2016, 17, 467-474.	1.3	12
72	Evaluations of treatment efficacy of depression from perspective of both patients' symptoms and general sense of mental health and wellbeing: A large scale, multi-centered, longitudinal study in China. Psychiatry Research, 2016, 241, 55-60.	1.7	4

#	ARTICLE	IF	CITATIONS
73	A study of single nucleotide polymorphisms of GRIN2B in schizophrenia from Chinese Han population. <i>Neuroscience Letters</i> , 2016, 630, 132-135.	1.0	9
74	Association study of dopamine receptor genes polymorphisms with the risk of schizophrenia in the Han Chinese population. <i>Psychiatry Research</i> , 2016, 245, 361-364.	1.7	8
75	A new method for identifying causal genes of schizophrenia and anti-tuberculosis drug-induced hepatotoxicity. <i>Scientific Reports</i> , 2016, 6, 32571.	1.6	16
76	Association study of 5-HT1A, 5-HT2A polymorphisms with schizophrenia and major depressive disorder in the Han Chinese population. <i>Neuroscience Letters</i> , 2016, 635, 39-43.	1.0	11
77	The China Mental Health Survey (CMHS): I. background, aims and measures. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2016, 51, 1559-1569.	1.6	40
78	The China Mental Health Survey: II. Design and field procedures. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2016, 51, 1547-1557.	1.6	34
79	Dysregulated 14-3-3 Family in Peripheral Blood Leukocytes of Patients with Schizophrenia. <i>Scientific Reports</i> , 2016, 6, 23791.	1.6	17
80	Role played by the SP4 gene in schizophrenia and major depressive disorder in the Han Chinese population. <i>British Journal of Psychiatry</i> , 2016, 208, 441-445.	1.7	3
81	A new risk locus in the ZEB2 gene for schizophrenia in the Han Chinese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2016, 66, 97-103.	2.5	10
82	Genome-wide Analysis of the Role of Copy Number Variation in Schizophrenia Risk in Chinese. <i>Biological Psychiatry</i> , 2016, 80, 331-337.	0.7	55
83	Analysis of association between common variants in the <i>SLCO6A1</i> gene with schizophrenia, bipolar disorder and major depressive disorder in the Han Chinese population. <i>World Journal of Biological Psychiatry</i> , 2016, 17, 140-146.	1.3	4
84	Abnormal white matter microstructure in drug-naive first episode schizophrenia patients before and after eight weeks of antipsychotic treatment. <i>Schizophrenia Research</i> , 2016, 172, 1-8.	1.1	75
85	Regional Abnormality of Grey Matter in Schizophrenia: Effect from the Illness or Treatment?. <i>PLoS ONE</i> , 2016, 11, e0147204.	1.1	37
86	Efficacy of Atypical Antipsychotics in the Management of Acute Agitation and Aggression in Hospitalized Patients with Schizophrenia or Bipolar Disorder: Results from a Systematic Review. <i>Shanghai Archives of Psychiatry</i> , 2016, 28, 241-252.	0.7	4
87	Increased Cognition Connectivity Network in Major Depression Disorder: A fMRI Study. <i>Psychiatry Investigation</i> , 2015, 12, 227.	0.7	40
88	The GSK3B gene confers risk for both major depressive disorder and schizophrenia in the Han Chinese population. <i>Journal of Affective Disorders</i> , 2015, 185, 149-155.	2.0	34
89	The NVL gene confers risk for both major depressive disorder and schizophrenia in the Han Chinese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2015, 62, 7-13.	2.5	17
90	Down-regulation of PRKCB1 expression in Han Chinese patients with subsyndromal symptomatic depression. <i>Journal of Psychiatric Research</i> , 2015, 69, 1-6.	1.5	6

#	ARTICLE	IF	CITATIONS
91	The MATRICS Consensus Cognitive Battery (MCCB): Co-norming and standardization in China. Schizophrenia Research, 2015, 169, 109-115.	1.1	176
92	Association study of GRM7 polymorphisms and schizophrenia in the Chinese Han population. Neuroscience Letters, 2015, 604, 109-112.	1.0	8
93	Sleep problems among clinically depressed adults in China. Journal of Mental Health, 2015, 24, 43-47.	1.0	1
94	Association between SREBF2 gene polymorphisms and metabolic syndrome in clozapine-treated patients with schizophrenia. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2015, 56, 136-141.	2.5	19
95	Assessment of a six-week computer-based remediation program for social cognition in chronic schizophrenia. Shanghai Archives of Psychiatry, 2015, 27, 296-306.	0.7	5
96	Meta-Analysis-Based Preliminary Exploration of the Connection between ATDILI and Schizophrenia by GSTM1/T1 Gene Polymorphisms. PLoS ONE, 2015, 10, e0128643.	1.1	16
97	Neural Correlates of the Preserved Inhibition of Return in Schizophrenia. PLoS ONE, 2015, 10, e0119521.	1.1	1
98	Comparison of the density of gamma-aminobutyric acid in the ventromedial prefrontal cortex of patients with first-episode psychosis and healthy controls. Shanghai Archives of Psychiatry, 2015, 27, 341-7.	0.7	13
99	Comparison of Chinese and international psychiatrists' views on classification of mental disorders. Asia-Pacific Psychiatry, 2014, 6, 267-273.	1.2	15
100	Cerebral metabolism in major depressive disorder: a voxel-based meta-analysis of positron emission tomography studies. BMC Psychiatry, 2014, 14, 321.	1.1	170
101	A study of N-methyl-D-aspartate receptor gene (GRIN2B) variants as predictors of treatment-resistant major depression. Psychopharmacology, 2014, 231, 685-693.	1.5	65
102	Wnt Signaling Pathway in Schizophrenia. CNS and Neurological Disorders - Drug Targets, 2014, 13, 755-764.	0.8	15
103	Advances in neuroimaging research of schizophrenia in China. Shanghai Archives of Psychiatry, 2014, 26, 181-93.	0.7	0
104	SOX10 rs139883 Polymorphism Is Associated with the Age of Onset in Schizophrenia. Journal of Molecular Neuroscience, 2013, 50, 333-338.	1.1	9
105	Prospective memory performance in patients with drug-naïve, first-episode psychosis. Schizophrenia Research, 2013, 143, 285-290.	1.1	22
106	Quality of life in outpatients with depression in China. Journal of Affective Disorders, 2013, 150, 513-521.	2.0	18
107	Hyperactivity within an extensive cortical distribution associated with excessive sensitivity in error processing in unmedicated depression: A combined event-related potential and sLORETA study. International Journal of Psychophysiology, 2013, 90, 282-289.	0.5	26
108	Efficacy of HuaDan Anshen Mistura for treating insomnia: a randomized, double-blind, placebo-controlled, multi-center clinical trial. Journal of Traditional Chinese Medicine = Chung I Tsa Chih Ying Wen Pan / Sponsored By All-China Association of Traditional Chinese Medicine, Academy of Traditional Chinese Medicine, 2013, 33, 423-427.	0.4	3

#	ARTICLE	IF	CITATIONS
109	Does practice make perfect? Results from a Chinese feasibility study of cognitive remediation in schizophrenia. <i>Neuropsychological Rehabilitation</i> , 2013, 23, 580-596.	1.0	6
110	Histamine H4 Receptor Polymorphism. <i>Journal of Clinical Psychopharmacology</i> , 2013, 33, 221-225.	0.7	19
111	Association study on the DLG4 gene and schizophrenia in the Chinese Han population. <i>Psychiatric Genetics</i> , 2013, 23, 247-250.	0.6	3
112	MicroRNA Microarray Analysis Combined with Interaction Network Analysis to Investigate the Influence of Clozapine to Metabolic Syndrome. <i>International Journal of Pharmacology</i> , 2013, 9, 366-372.	0.1	1
113	Case report of Folie À Trois. <i>Shanghai Archives of Psychiatry</i> , 2013, 25, 52-4.	0.7	0
114	Should repetitive Transcranial Magnetic Stimulation (rTMS) be considered an effective adjunctive treatment for auditory hallucinations in patients with schizophrenia?. <i>Shanghai Archives of Psychiatry</i> , 2013, 25, 254-5.	0.7	3
115	Research progress in China on the assessment of cognitive function in schizophrenia. <i>Shanghai Archives of Psychiatry</i> , 2013, 25, 266-75.	0.7	2
116	NAPG may not be a common risk gene shared by bipolar disorder and schizophrenia in Chinese population. <i>Psychiatric Genetics</i> , 2012, 22, 107-108.	0.6	0
117	Association between a COMT polymorphism and clinical response to risperidone treatment. <i>Psychiatric Genetics</i> , 2012, 22, 298-299.	0.6	22
118	A pharmacogenetic study of risperidone on histamine H3 receptor gene ( <i>HRH3</i> ) in Chinese Han schizophrenia patients. <i>Journal of Psychopharmacology</i> , 2012, 26, 813-818.	2.0	19
119	Teachers of Psychiatry meeting in Shanghai: A leadership training course. <i>Asia-Pacific Psychiatry</i> , 2012, 4, 87-89.	1.2	3
120	Association study of APC polymorphisms with colorectal cancer in Han Chinese. <i>Clinical Biochemistry</i> , 2012, 45, 1669-1672.	0.8	3
121	Identification of the <i>N-acylsphingosine amidohydrolase 1 gene (ASAH1)</i> for susceptibility to schizophrenia in a Han Chinese population. <i>World Journal of Biological Psychiatry</i> , 2012, 13, 106-113.	1.3	10
122	Effect of SOX10 gene polymorphism on early onset schizophrenia in Chinese Han population. <i>Neuroscience Letters</i> , 2012, 521, 93-97.	1.0	10
123	Chronic mild restraint stress rats decreased CMKLR1 expression in distinct brain region. <i>Neuroscience Letters</i> , 2012, 524, 25-29.	1.0	29
124	Ethics, patient rights and staff attitudes in Shanghai's psychiatric hospitals. <i>BMC Medical Ethics</i> , 2012, 13, 8.	1.0	7
125	Common Variants in the TPH2 Promoter Confer Susceptibility to Paranoid Schizophrenia. <i>Journal of Molecular Neuroscience</i> , 2012, 47, 465-469.	1.1	10
126	Social support and clinical depression in China. <i>Asia-Pacific Psychiatry</i> , 2012, 4, 195-200.	1.2	1



#	ARTICLE	IF	CITATIONS
127	The rights of psychiatric patients in China: A survey of medical staff and consumers' attitudes toward patient participation in clinical trials. <i>Social Science and Medicine</i> , 2012, 75, 823-827.	1.8	7
128	Polymorphisms and Haplotypes in the YWHAEGene Increase Susceptibility to Bipolar Disorder in Chinese Han Population. <i>Journal of Clinical Psychiatry</i> , 2012, 73, e1276-e1282.	1.1	14
129	Symptom severity is more closely associated with social functioning status in inpatients with schizophrenia than cognitive deficits. <i>Shanghai Archives of Psychiatry</i> , 2012, 24, 83-90.	0.7	0
130	Mental health system in China: history, recent service reform and future challenges. <i>World Psychiatry</i> , 2011, 10, 210-216.	4.8	232
131	Common variants on 8p12 and 1q24.2 confer risk of schizophrenia. <i>Nature Genetics</i> , 2011, 43, 1224-1227.	9.4	224
132	No association between the KCNH1, KCNJ10 and KCNN3 genes and schizophrenia in the Han Chinese population. <i>Neuroscience Letters</i> , 2011, 487, 61-65.	1.0	4
133	A Pilot Study of the Efficacy and Safety of Paroxetine Augmented With Risperidone, Valproate, Buspirone, Trazodone, or Thyroid Hormone in Adult Chinese Patients With Treatment-Resistant Major Depression. <i>Journal of Clinical Psychopharmacology</i> , 2011, 31, 638-642.	0.7	47
134	An Evaluation of the Shanghai Mental Health Service Schizophrenia Rehabilitation Program. <i>American Journal of Psychiatric Rehabilitation</i> , 2011, 14, 287-306.	0.7	5
135	Comparisons of the Efficacy and Tolerability of Extended-Release Venlafaxine, Mirtazapine, and Paroxetine in Treatment-Resistant Depression. <i>Journal of Clinical Psychopharmacology</i> , 2010, 30, 357-364.	0.7	48
136	Polymorphisms of XRCC4 are involved in reduced colorectal cancer risk in Chinese schizophrenia patients. <i>BMC Cancer</i> , 2010, 10, 523.	1.1	17
137	<i>HTR2C</i> promoter polymorphisms are associated with risperidone efficacy in Chinese female patients. <i>Pharmacogenomics</i> , 2010, 11, 685-692.	0.6	15
138	No association between EGR gene family polymorphisms and schizophrenia in the Chinese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2010, 34, 506-509.	2.5	14
139	Deficient inhibition of return in chronic but not first-episode patients with schizophrenia. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2010, 34, 961-967.	2.5	7
140	Pharmacogenetic effects of dopamine transporter gene polymorphisms on response to chlorpromazine and clozapine and on extrapyramidal syndrome in schizophrenia. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2010, 34, 1026-1032.	2.5	46
141	Genetic polymorphisms in the SCN8A gene are associated with suicidal behavior in psychiatric disorders in the Chinese population. <i>World Journal of Biological Psychiatry</i> , 2010, 11, 956-963.	1.3	9
142	Schizophrenia is not associated with the ERBB3 gene in a Han Chinese population sample: results from case-control and family-based studies. <i>Genetics and Molecular Biology</i> , 2009, 32, 729-730.	0.6	1
143	Family-based association study between brain-derived neurotrophic factor gene and major depressive disorder of Chinese descent. <i>Psychiatry Research</i> , 2009, 169, 169-172.	1.7	17
144	No association found between the promoter variations of QKI and schizophrenia in the Chinese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2009, 33, 33-36.	2.5	4

#	ARTICLE	IF	CITATIONS
145	A case-control association study between the CYP3A4 and CYP3A5 genes and schizophrenia in the Chinese Han population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2009, 33, 1200-1204.	2.5	10
146	Positive association between ALDH1A2 and schizophrenia in the Chinese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2009, 33, 1491-1495.	2.5	25
147	Haplotype analysis confirms association of the serotonin transporter (5-HTT) gene with schizophrenia in the Han Chinese population. <i>Neuroscience Letters</i> , 2009, 453, 210-213.	1.0	15
148	Positive association between OLIG2 and schizophrenia in the Chinese Han population. <i>Human Genetics</i> , 2008, 122, 659-660.	1.8	30
149	No significant association between the genetic polymorphisms in the GSK-3 $\beta$ gene and schizophrenia in the Chinese population. <i>Journal of Psychiatric Research</i> , 2008, 42, 365-370.	1.5	15
150	No association found between the promoter variants of ADRA1A and schizophrenia in the Chinese population. <i>Journal of Psychiatric Research</i> , 2008, 42, 384-388.	1.5	7
151	The Ser9Gly polymorphism of the dopamine D3 receptor gene and risk of schizophrenia: An association study and a large meta-analysis. <i>Schizophrenia Research</i> , 2008, 101, 26-35.	1.1	22
152	No genetic association between polymorphisms in the kainate-type glutamate receptor gene, GRIK4, and schizophrenia in the Chinese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2008, 32, 876-880.	2.5	8
153	No genetic association between NCAM1 gene polymorphisms and schizophrenia in the Chinese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2008, 32, 1633-1636.	2.5	9
154	An association study between PPP1R1B gene and schizophrenia in the Chinese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2007, 31, 1303-1306.	2.5	14
155	Positive association between SIAT8B and schizophrenia in the Chinese Han population. <i>Schizophrenia Research</i> , 2007, 90, 108-114.	1.1	87
156	RGS4 polymorphisms and risk of schizophrenia: An association study in Han Chinese plus meta-analysis. <i>Neuroscience Letters</i> , 2006, 406, 122-127.	1.0	29
157	A family-based association study of kinesin heavy chain member 2 gene (KIF2) and schizophrenia. <i>Neuroscience Letters</i> , 2006, 407, 151-155.	1.0	16
158	Dopamine D4 receptor polymorphism modulates cue-elicited heroin craving in Chinese. <i>Psychopharmacology</i> , 2006, 186, 185-190.	1.5	70
159	A case-control study of the relationship between the metabotropic glutamate receptor 3 gene and schizophrenia in the Chinese population. <i>Schizophrenia Research</i> , 2005, 73, 21-26.	1.1	80
160	A family-based association study of the MOG gene with schizophrenia in the Chinese population. <i>Schizophrenia Research</i> , 2005, 73, 275-280.	1.1	29
161	Genetic Structure Adds Power to Detect Schizophrenia Susceptibility at SLIT3 in the Chinese Han Population. <i>Genome Research</i> , 2004, 14, 1345-1349.	2.4	36
162	No association between the genetic polymorphisms within RTN4 and schizophrenia in the Chinese population. <i>Neuroscience Letters</i> , 2004, 365, 23-27.	1.0	12

#	ARTICLE	IF	CITATIONS
163	No association between the promoter variants of tumor necrosis factor alpha (TNF- $\hat{\pm}$ ) and schizophrenia in Chinese Han population. <i>Neuroscience Letters</i> , 2004, 366, 139-143.	1.0	39
164	Family-Based Association Study of Synapsin II and Schizophrenia. <i>American Journal of Human Genetics</i> , 2004, 75, 873-877.	2.6	44
165	The Epidemiology of Depressive Disorders and Service Use in China: Findings from the China Mental Health Survey. <i>SSRN Electronic Journal</i> , 0, , .	0.4	0