Michael Blackburn

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/258863/publications.pdf

Version: 2024-02-01

1684188 1372567 11 139 5 10 citations g-index h-index papers 11 11 11 244 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Transcriptome of the Lymantria dispar (Gypsy Moth) Larval Midgut in Response to Infection by Bacillus thuringiensis. PLoS ONE, 2013, 8, e61190.	2.5	46
2	Enteric bacteria of field-collected Colorado potato beetle larvae inhibit growth of the entomopathogens Photorhabdus temperata and Beauveria bassiana. Biological Control, 2008, 46, 434-441.	3.0	31
3	The Occurrence of Photorhabdus-Like Toxin Complexes in Bacillus thuringiensis. PLoS ONE, 2011, 6, e18122.	2.5	21
4	Phylogenetic Distribution of Phenotypic Traits in Bacillus thuringiensis Determined by Multilocus Sequence Analysis. PLoS ONE, 2013, 8, e66061.	2.5	12
5	Reproductive failure of Heterorhabditis marelatus in the Colorado potato beetle: Evidence of stress on the nematode symbiont Photorhabdus temperata and potential interference from the enteric bacteria of the beetle. Biological Control, 2007, 42, 207-215.	3.0	11
6	Insecticidal Activity of <i>Chromobacterium vaccinii</i> ¹ . Journal of Entomological Science, 2018, 53, 339-346.	0.3	5
7	Cabbage looper (Trichoplusia ni Hübner) labial glands contain unique bacterial flora in contrast with their alimentary canal, mandibular glands, and Malpighian tubules. MicrobiologyOpen, 2020, 9, e994.	3.0	5
8	The genome of the insecticidal Chromobacterium subtsugae PRAA4-1 and its comparison with that of Chromobacterium violaceum ATCC 12472. Genomics Data, 2016, 10, 1-3.	1.3	4
9	Insecticidal Activity of a Recently Described Bacterium, Chromobacterium sphagni 1. Journal of Entomological Science, 2018, 53, 333-338.	0.3	2
10	The response to cabbage looper infestation in Arabidopsis is altered by lowering levels of Zat18 a Q-type C2H2 zinc finger protein. Journal of Plant Interactions, 2022, 17, 198-205.	2.1	2
11	Crystalliferous <i>Bacillus cereus</i> group bacteria from a <scp>M</scp> aryland hardwood forest are dominated by psychrotolerant strains. MicrobiologyOpen, 2014, 3, 578-584.	3.0	O