

Anette Larsson

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2588677/publications.pdf>

Version: 2024-02-01

16
papers

466
citations

840776

11
h-index

1058476

14
g-index

16
all docs

16
docs citations

16
times ranked

591
citing authors

#	ARTICLE	IF	CITATIONS
1	Resistance exercise improves muscle strength, health status and pain intensity in fibromyalgia—a randomized controlled trial. <i>Arthritis Research and Therapy</i> , 2015, 17, 161.	3.5	122
2	Gene-to-gene interactions regulate endogenous pain modulation in fibromyalgia patients and healthy controls—antagonistic effects between opioid and serotonin-related genes. <i>Pain</i> , 2017, 158, 1194-1203.	4.2	54
3	Plasma Cytokine Levels in Fibromyalgia and Their Response to 15 Weeks of Progressive Resistance Exercise or Relaxation Therapy. <i>Mediators of Inflammation</i> , 2018, 2018, 1-14.	3.0	53
4	Resistance exercise improves physical fatigue in women with fibromyalgia: a randomized controlled trial. <i>Arthritis Research and Therapy</i> , 2016, 18, 176.	3.5	52
5	Comparison of the Levels of Pro-Inflammatory Cytokines Released in the Vastus Lateralis Muscle of Patients with Fibromyalgia and Healthy Controls during Contractions of the Quadriceps Muscle—A Microdialysis Study. <i>PLoS ONE</i> , 2015, 10, e0143856.	2.5	32
6	Decrease of fear avoidance beliefs following person-centered progressive resistance exercise contributes to reduced pain disability in women with fibromyalgia: secondary exploratory analyses from a randomized controlled trial. <i>Arthritis Research and Therapy</i> , 2016, 18, 116.	3.5	28
7	Controlled, cross-sectional, multi-center study of physical capacity and associated factors in women with fibromyalgia. <i>BMC Musculoskeletal Disorders</i> , 2018, 19, 121.	1.9	23
8	Effects of 15 weeks of resistance exercise on pro-inflammatory cytokine levels in the vastus lateralis muscle of patients with fibromyalgia. <i>Arthritis Research and Therapy</i> , 2016, 18, 137.	3.5	22
9	Benefits of resistance exercise in lean women with fibromyalgia: involvement of IGF-1 and leptin. <i>BMC Musculoskeletal Disorders</i> , 2017, 18, 106.	1.9	19
10	Perceived exertion at work in women with fibromyalgia: Explanatory factors and comparison with healthy women. <i>Journal of Rehabilitation Medicine</i> , 2014, 46, 773-780.	1.1	18
11	Pain and fear avoidance partially mediate change in muscle strength during resistance exercise in women with fibromyalgia. <i>Journal of Rehabilitation Medicine</i> , 2017, 49, 744-750.	1.1	15
12	Factors promoting physical activity in women with fibromyalgia: a qualitative interview study. <i>BMJ Open</i> , 2020, 10, e031693.	1.9	13
13	Acute effects of physical exercise on the serum insulin-like growth factor system in women with fibromyalgia. <i>BMC Musculoskeletal Disorders</i> , 2017, 18, 37.	1.9	10
14	Physical activity with person-centred guidance supported by a digital platform for persons with chronic widespread pain: A randomized controlled trial. <i>Journal of Rehabilitation Medicine</i> , 2021, 53, jrm00175.	1.1	4
15	Person-centred health plans for physical activity in persons with chronic widespread pain (CWP)—a retrospective descriptive review. <i>Disability and Rehabilitation</i> , 0, , 1-8.	1.8	1
16	Experience of co-creation of a health plan and support for sustainable physical activity among people with chronic widespread pain: a qualitative interview study in Sweden. <i>BMJ Open</i> , 2022, 12, e059432.	1.9	0