
## Kaj Sparle Christensen

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2590511/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Rasch analysis of the structural validity of the anxiety symptom scale in the lolland-falster health study. Journal of Affective Disorders, 2022, 303, 58-63.	2.0	5
2	Functional somatic symptoms in youths in general practice: A cross-sectional study on prevalence, clinical management and perceived burden. Journal of Psychosomatic Research, 2022, 156, 110765.	1.2	3
3	Cross-cultural validity of the WHO-5 Well-Being Index and Euthymia Scale: A clinimetric analysis. Journal of Affective Disorders, 2022, 311, 276-283.	2.0	12
4	Maternal depression and childhood injury risk: A populationâ€based cohort study in Denmark. Brain and Behavior, 2021, 11, e02029.	1.0	5
5	Clinimetric Criteria for Patient-Reported Outcome Measures. Psychotherapy and Psychosomatics, 2021, 90, 222-232.	4.0	84
6	Construct and criterion validity of patient-reported outcomes (PROs) for depression: A clinimetric comparison. Journal of Affective Disorders, 2021, 283, 30-35.	2.0	18
7	Improving the precision of depression diagnosis in general practice: a cluster-randomized trial. BMC Family Practice, 2021, 22, 88.	2.9	1
8	Construct validity of the Smoker Complaint Scale: A clinimetric analysis using Item Response Theory (IRT) models. Addictive Behaviors, 2021, 117, 106849.	1.7	2
9	Insufficient classification of anaemia in general practice: a Danish register-based observational study. Scandinavian Journal of Primary Health Care, 2021, 39, 364-372.	0.6	3
10	A clinimetric analysis of the euthymia, resilience, and positive mental health scales. Journal of Affective Disorders, 2021, 294, 71-76.	2.0	11
11	Clinimetric properties of the Smoking Abstinence Expectancies Questionnaire. Addictive Behaviors, 2021, 123, 107061.	1.7	5
12	Maternal depression and non-specific health complaints in the offspring: a cross-sectional study in Danish primary care. British Journal of General Practice, 2021, 71, e105-e112.	0.7	1
13	The Hopkins Symptom Checklist (SCL-90-R): A Patient-Reported Outcome Measure in Parkinson's Disease. Journal of Geriatric Psychiatry and Neurology, 2021, , 089198872110600.	1.2	0
14	Collaborative care for depression and anxiety disorders: results and lessons learned from the Danish cluster-randomized Collabri trials. BMC Family Practice, 2020, 21, 234.	2.9	10
15	Exploring Functional Impairment in Light of Prolonged Grief Disorder: A Prospective, Population-Based Cohort Study. Frontiers in Psychiatry, 2020, 11, 537674.	1.3	11
16	Factors associated with non-initiation of mental healthcare after detection of poor mental health at a scheduled health check: a cohort study. BMJ Open, 2020, 10, e037731.	0.8	0
17	Factors associated with non-initiation of mental healthcare after detection of poor mental health at a scheduled health check: a cohort study. BMJ Open, 2020, 10, e037731.	0.8	3
18	Diagnosing depression in primary care: a Rasch analysis of the major depression inventory. Scandinavian Journal of Primary Health Care, 2019, 37, 105-112.	0.6	10

#	Article	IF	CITATIONS
19	Incidence of medically treated depression in Denmark among individuals 15–44Âyears old: a comprehensive overview based on population registers. Acta Psychiatrica Scandinavica, 2019, 139, 548-557.	2.2	44
20	Diagnosing depression in primary care: a Rasch analysis of the Major Depression Inventory. Scandinavian Journal of Primary Health Care, 2019, 37, 256-263.	0.6	8
21	Maternal depression and primary healthcare use for children: a population-based cohort study in Denmark. British Journal of General Practice, 2019, 69, e182-e189.	0.7	9
22	Mental health assessment in health checks of participants aged 30–49 years: A large-scale cohort study. Preventive Medicine Reports, 2018, 9, 72-79.	0.8	4
23	Initiation and cessation of mental healthcare after mental health screening in primary care: a prospective cohort study. BMC Family Practice, 2018, 19, 176.	2.9	22
24	Barriers and facilitators to using a web-based tool for diagnosis and monitoring of patients with depression: a qualitative study among Danish general practitioners. BMC Health Services Research, 2018, 18, 503.	0.9	25
25	Screening for depression: Rasch analysis of the structural validity of the PHQ-9 in acutely injured trauma survivors. Journal of Psychosomatic Research, 2017, 97, 18-22.	1.2	17
26	The construct validity of the Major Depression Inventory: A Rasch analysis of a self-rating scale in primary care. Journal of Psychosomatic Research, 2017, 97, 70-81.	1.2	13
27	Collaborative care for depression in general practice: study protocol for a randomised controlled trial. Trials, 2017, 18, 344.	0.7	7
28	The criterion validity of the web-based Major Depression Inventory when used on clinical suspicion of depression in primary care. Clinical Epidemiology, 2017, Volume 9, 355-365.	1.5	12
29	The construct validity of the Perceived Stress Scale. Journal of Psychosomatic Research, 2016, 84, 22-30.	1.2	114
30	The Association Between Perceived Stress and Mortality Among People With Multimorbidity: A Prospective Population-Based Cohort Study. American Journal of Epidemiology, 2016, 184, 199-210.	1.6	119
31	Development and validation of a self-efficacy questionnaire (SE-12) measuring the clinical communication skills of health care professionals. BMC Medical Education, 2016, 16, 272.	1.0	61
32	Identifying depression among adolescents using three key questions: a validation study in primary care. British Journal of General Practice, 2016, 66, e65-e70.	0.7	10
33	Chlamydia trachomatis infection in young adults — association with concurrent partnerships and short gap length between partners. Infectious Diseases, 2015, 47, 838-845.	1.4	7
34	Diagnosis of depressed young people–criterion validity of WHO-5 and HSCL-6 in Denmark and Norway. Family Practice, 2015, 32, 359-363.	0.8	16
35	A new questionnaire to identify bodily distress in primary care: The †BDS checklist'. Journal of Psychosomatic Research, 2015, 78, 536-545.	1.2	78
36	Sexual behaviour among young Danes aged 15–29 years: a cross-sectional study of core indicators. Sexually Transmitted Infections, 2015, 91, 171-177.	0.8	23

#	Article	IF	CITATIONS
37	Prevalence of depression, quality of life and antidepressant treatment in the Danish General Suburban Population Study. Nordic Journal of Psychiatry, 2014, 68, 507-512.	0.7	60
38	Mental health status and risk of new cardiovascular events or death in patients with myocardial infarction: a population-based cohort study. BMJ Open, 2013, 3, e003045.	0.8	26
39	Psychological and social problems in primary care patients - general practitioners' assessment and classification. Scandinavian Journal of Primary Health Care, 2013, 31, 43-49.	0.6	34
40	Diagnosis of depressed young people in primary health carea validation of HSCL-10. Family Practice, 2011, 28, 233-237.	0.8	54
41	Case-finding and risk-group screening for depression in primary care. Scandinavian Journal of Primary Health Care, 2011, 29, 80-84.	0.6	31
42	The Outcome of Health Anxiety in Primary Care. A Two-Year Follow-up Study on Health Care Costs and Self-Rated Health. PLoS ONE, 2010, 5, e9873.	1.1	169
43	Do illness perceptions predict health outcomes in primary care patients? A 2-year follow-up study. Journal of Psychosomatic Research, 2007, 62, 129-138.	1.2	106
44	Diagnosis of depression among adolescents – a clinical validation study of key questions and questionnaire. BMC Family Practice, 2007, 8, 41.	2.9	6
45	Diagnosing depression. British Journal of General Practice, 2007, 57, 835-6; author reply 836.	0.7	3
46	A comment. Scandinavian Journal of Gastroenterology, 2006, 41, 995-995.	0.6	0
47	The Uncertain Consultation and Patient Satisfaction: The Impact of Patients' Illness Perceptions and a Randomized Controlled Trial on the Training of Physicians' Communication Skills. Psychosomatic Medicine, 2005, 67, 897-905.	1.3	85
48	The Patients' Illness Perceptions and the Use of Primary Health Care. Psychosomatic Medicine, 2005, 67, 997-1005.	1.3	120
49	Mental disorders in primary care: prevalence and co-morbidity among disorders. Results from the Functional Illness in Primary care (FIP) study. Psychological Medicine, 2005, 35, 1175-1184.	2.7	214
50	A brief case-finding questionnaire for common mental disorders: the CMDQ. Family Practice, 2005, 22, 448-457.	0.8	136
51	Screening for common mental disorders: who will benefit? Results from a randomised clinical trial. Family Practice, 2005, 22, 428-434.	0.8	28
52	Mental health screening may prove effective in primary care. British Journal of General Practice, 2005, 55, 967.	0.7	3
53	A New, Empirically Established Hypochondriasis Diagnosis. American Journal of Psychiatry, 2004, 161, 1680-1691.	4.0	178
54	The FIP study: a randomised, controlled trial of screening and recognition of psychiatric disorders. British Journal of General Practice, 2003, 53, 758-63.	0.7	34

#	Article	IF	CITATIONS
55	Diagnosis of somatisation: effect of an educational intervention in a cluster randomised controlled trial. British Journal of General Practice, 2003, 53, 917-22.	0.7	43