

Seung Hong Choi

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/2591932/seung-hong-choi-publications-by-year.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

187
papers

11,147
citations

43
h-index

103
g-index

191
ext. papers

12,744
ext. citations

7.9
avg, IF

6.04
L-index

| # | Paper | IF | Citations |
|-----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----------|
| 187 | Enhanced Chemodynamic Therapy by Cu-Fe Peroxide Nanoparticles: Tumor Microenvironment-Mediated Synergistic Fenton Reaction.. <i>ACS Nano</i> , 2022 , | 16.7 | 18 |
| 186 | Tumor-associated macrophages in cancer: recent advancements in cancer nanoimmunotherapies.. <i>Journal of Experimental and Clinical Cancer Research</i> , 2022 , 41, 68 | 12.8 | 6 |
| 185 | No Prognostic Impact of Staging Brain MRI in Patients with Stage IA Non-Small Cell Lung Cancer.. <i>Radiology</i> , 2022 , 212101 | 20.5 | 0 |
| 184 | Localized Delivery of Theranostic Nanoparticles and High-Energy Photons using Microneedles-on-Bioelectronics. <i>Advanced Materials</i> , 2021 , 33, e2100425 | 24 | 15 |
| 183 | Differentiation between glioblastoma and primary CNS lymphoma: application of DCE-MRI parameters based on arterial input function obtained from DSC-MRI. <i>European Radiology</i> , 2021 , 31, 9098-9109 ⁸ 9109 ¹ | | |
| 182 | Tissue-like skin-device interface for wearable bioelectronics by using ultrasoft, mass-permeable, and low-impedance hydrogels. <i>Science Advances</i> , 2021 , 7, | 14.3 | 56 |
| 181 | Multiparametric magnetic resonance imaging features of a canine glioblastoma model. <i>PLoS ONE</i> , 2021 , 16, e0254448 | 3.7 | 0 |
| 180 | Soft implantable drug delivery device integrated wirelessly with wearable devices to treat fatal seizures. <i>Science Advances</i> , 2021 , 7, | 14.3 | 36 |
| 179 | Comparison of Genetic Profiles and Prognosis of High-Grade Gliomas Using Quantitative and Qualitative MRI Features: A Focus on G3 Gliomas. <i>Korean Journal of Radiology</i> , 2021 , 22, 233-242 | 6.9 | 1 |
| 178 | Blood-Brain Barrier Disruption in Mild Traumatic Brain Injury Patients with Post-Concussion Syndrome: Evaluation with Region-Based Quantification of Dynamic Contrast-Enhanced MR Imaging Parameters Using Automatic Whole-Brain Segmentation. <i>Korean Journal of Radiology</i> , 2021 , 22, 118-130 | 6.9 | 4 |
| 177 | Application of T1 Map Information Based on Synthetic MRI for Dynamic Contrast-Enhanced Imaging: A Comparison Study with the Fixed Baseline T1 Value Method. <i>Korean Journal of Radiology</i> , 2021 , 22, 1352-1368 | 6.9 | |
| 176 | Prediction of Prognosis in Glioblastoma Using Radiomics Features of Dynamic Contrast-Enhanced MRI. <i>Korean Journal of Radiology</i> , 2021 , 22, 1514-1524 | 6.9 | 3 |
| 175 | Cognitive improvement effect of gintonin might be associated with blood-brain barrier permeability enhancement: dynamic contrast-enhanced MRI pilot study. <i>Translational and Clinical Pharmacology</i> , 2021 , 29, 21-32 | 2 | 0 |
| 174 | Quad-Contrast Imaging: Simultaneous Acquisition of Four Contrast-Weighted Images (PD-Weighted, T ₂ Weighted, PD-FLAIR and T ₂ FLAIR Images) With Synthetic T ₂ Weighted Image, T ₂ and T ₂ Maps. <i>IEEE Transactions on Medical Imaging</i> , 2021 , 40, 3617-3626 | 11.7 | 2 |
| 173 | Prognostic Prediction Based on Dynamic Contrast-Enhanced MRI and Dynamic Susceptibility Contrast-Enhanced MRI Parameters from Non-Enhancing, T2-High-Signal-Intensity Lesions in Patients with Glioblastoma. <i>Korean Journal of Radiology</i> , 2021 , 22, 1369-1378 | 6.9 | 0 |
| 172 | Promoter Methylation Status in Initial and Recurrent Glioblastoma: Correlation Study with DWI and DSC PWI Features. <i>American Journal of Neuroradiology</i> , 2021 , 42, 853-860 | 4.4 | 4 |
| 171 | Contrast-enhanced MRI T1 Mapping for Quantitative Evaluation of Putative Dynamic Glymphatic Activity in the Human Brain in Sleep-Wake States. <i>Radiology</i> , 2021 , 300, 661-668 | 20.5 | 5 |

| | | | |
|-----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|----|
| 170 | Cerebrovascular Reservoir and Arterial Transit Time Changes Assessed by Acetazolamide-Challenged Multi-Phase Arterial Spin Labeling Perfusion MRI in Chronic Cerebrovascular Steno-Occlusive Disease. <i>Journal of the Korean Society of Radiology</i> , 2021 , 82, 626 | 0.2 | 1 |
| 169 | Added Value of Computed Tomography to Ultrasonography for Assessing LN Metastasis in Preoperative Patients with Thyroid Cancer: Node-By-Node Correlation. <i>Cancers</i> , 2020 , 12, | 6.6 | 6 |
| 168 | Diagnostic Accuracy and Confidence of [18F] FDG PET/MRI in comparison with PET or MRI alone in Head and Neck Cancer. <i>Scientific Reports</i> , 2020 , 10, 9490 | 4.9 | 10 |
| 167 | Radiomics prognostication model in glioblastoma using diffusion- and perfusion-weighted MRI. <i>Scientific Reports</i> , 2020 , 10, 4250 | 4.9 | 21 |
| 166 | Intracranial Metaplastic Meningioma : Clinical and Radiological Characteristics of 11 Cases. <i>Journal of Korean Neurosurgical Society</i> , 2020 , 63, 657-663 | 2.3 | 1 |
| 165 | Ultrasonographic Indeterminate Lymph Nodes in Preoperative Thyroid Cancer Patients: Malignancy Risk and Ultrasonographic Findings Predictive of Malignancy. <i>Korean Journal of Radiology</i> , 2020 , 21, 598-604 | 6.0 | 6 |
| 164 | Prognostic Value of Dynamic Contrast-Enhanced MRI-Derived Pharmacokinetic Variables in Glioblastoma Patients: Analysis of Contrast-Enhancing Lesions and Non-Enhancing T2 High-Signal Intensity Lesions. <i>Korean Journal of Radiology</i> , 2020 , 21, 707-716 | 6.9 | 2 |
| 163 | Advances in drug delivery technology for the treatment of glioblastoma multiforme. <i>Journal of Controlled Release</i> , 2020 , 328, 350-367 | 11.7 | 25 |
| 162 | Advances in Soft Bioelectronics for Brain Research and Clinical Neuroengineering. <i>Matter</i> , 2020 , 3, 1923-1947 | 12.7 | 15 |
| 161 | Synthetic MRI: Technologies and Applications in Neuroradiology. <i>Journal of Magnetic Resonance Imaging</i> , 2020 , | 5.6 | 9 |
| 160 | Improving the Reliability of Pharmacokinetic Parameters at Dynamic Contrast-enhanced MRI in Astrocytomas: A Deep Learning Approach. <i>Radiology</i> , 2020 , 297, 178-188 | 20.5 | 3 |
| 159 | Dynamic Contrast-Enhanced MR Imaging of Nonenhancing T2 High-Signal-Intensity Lesions in Baseline and Posttreatment Glioblastoma: Temporal Change and Prognostic Value. <i>American Journal of Neuroradiology</i> , 2020 , 41, 49-56 | 4.4 | 8 |
| 158 | Prediction of IDH genotype in gliomas with dynamic susceptibility contrast perfusion MR imaging using an explainable recurrent neural network. <i>Neuro-Oncology</i> , 2019 , 21, 1197-1209 | 1 | 36 |
| 157 | Assessment of Early Therapeutic Response to Nitroxoline in Temozolomide-Resistant Glioblastoma by Amide Proton Transfer Imaging: A Preliminary Comparative Study with Diffusion-weighted Imaging. <i>Scientific Reports</i> , 2019 , 9, 5585 | 4.9 | 8 |
| 156 | Evaluation of Tumor Blood Flow Using Alternate Ascending/Descending Directional Navigation in Primary Brain Tumors: A Comparison Study with Dynamic Susceptibility Contrast Magnetic Resonance Imaging. <i>Korean Journal of Radiology</i> , 2019 , 20, 275-282 | 6.9 | 2 |
| 155 | Quantitative dynamic contrast-enhanced MR imaging shows widespread blood-brain barrier disruption in mild traumatic brain injury patients with post-concussion syndrome. <i>European Radiology</i> , 2019 , 29, 1308-1317 | 8 | 17 |
| 154 | Incorporating diffusion- and perfusion-weighted MRI into a radiomics model improves diagnostic performance for pseudoprogression in glioblastoma patients. <i>Neuro-Oncology</i> , 2019 , 21, 404-414 | 1 | 73 |
| 153 | Increased Antiangiogenic Effect by Blocking CCL2-dependent Macrophages in a Rodent Glioblastoma Model: Correlation Study with Dynamic Susceptibility Contrast Perfusion MRI. <i>Scientific Reports</i> , 2019 , 9, 11085 | 4.9 | 24 |

| | | | |
|-----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----|
| 152 | Altered Vascular Permeability in Migraine-associated Brain Regions: Evaluation with Dynamic Contrast-enhanced MRI. <i>Radiology</i> , 2019 , 292, 713-720 | 20.5 | 15 |
| 151 | Decreased APE-1 by Nitroxoline Enhances Therapeutic Effect in a Temozolomide-resistant Glioblastoma: Correlation with Diffusion Weighted Imaging. <i>Scientific Reports</i> , 2019 , 9, 16613 | 4.9 | 6 |
| 150 | Flexible, sticky, and biodegradable wireless device for drug delivery to brain tumors. <i>Nature Communications</i> , 2019 , 10, 5205 | 17.4 | 91 |
| 149 | Prognostic Predictions for Patients with Glioblastoma after Standard Treatment: Application of Contrast Leakage Information from DSC-MRI within Nonenhancing FLAIR High-Signal-Intensity Lesions. <i>American Journal of Neuroradiology</i> , 2019 , 40, 2052-2058 | 4.4 | 1 |
| 148 | Leakage correction improves prognosis prediction of dynamic susceptibility contrast perfusion MRI in primary central nervous system lymphoma. <i>Scientific Reports</i> , 2018 , 8, 456 | 4.9 | 5 |
| 147 | Application of 3D Fast Spin-Echo T1 Black-Blood Imaging in the Diagnosis and Prognostic Prediction of Patients with Leptomeningeal Carcinomatosis. <i>American Journal of Neuroradiology</i> , 2018 , 39, 1453-1459 | 4.4 | 13 |
| 146 | Can Amide Proton Transfer MRI Distinguish Benign and Malignant Head and Neck Tumors?. <i>Radiology</i> , 2018 , 288, 791-792 | 20.5 | 4 |
| 145 | Differentiation of High-Grade from Low-Grade Astrocytoma: Improvement in Diagnostic Accuracy and Reliability of Pharmacokinetic Parameters from DCE MR Imaging by Using Arterial Input Functions Obtained from DSC MR Imaging. <i>Radiology</i> , 2018 , 286, 981-991 | 20.5 | 11 |
| 144 | T1 Shortening in the Globus Pallidus after Multiple Administrations of Gadobutrol: Assessment with a Multidynamic Multiecho Sequence. <i>Radiology</i> , 2018 , 287, 258-266 | 20.5 | 24 |
| 143 | Can Arterial Spin-Labeling with Multiple Postlabeling Delays Predict Cerebrovascular Reserve?. <i>American Journal of Neuroradiology</i> , 2018 , 39, 84-90 | 4.4 | 7 |
| 142 | Loss of Pericytes in Radiation Necrosis after Glioblastoma Treatments. <i>Molecular Neurobiology</i> , 2018 , 55, 4918-4926 | 6.2 | 8 |
| 141 | . <i>American Journal of Neuroradiology</i> , 2018 , 39, E126 | 4.4 | 3 |
| 140 | Radiogenomics Profiling for Glioblastoma-related Immune Cells Reveals CD49d Expression Correlation with MRI parameters and Prognosis. <i>Scientific Reports</i> , 2018 , 8, 16022 | 4.9 | 15 |
| 139 | Prognosis prediction of non-enhancing T2 high signal intensity lesions in glioblastoma patients after standard treatment: application of dynamic contrast-enhanced MR imaging. <i>European Radiology</i> , 2017 , 27, 1176-1185 | 8 | 20 |
| 138 | Wearable/disposable sweat-based glucose monitoring device with multistage transdermal drug delivery module. <i>Science Advances</i> , 2017 , 3, e1601314 | 14.3 | 596 |
| 137 | Dynamic contrast-enhanced MR imaging in predicting progression of enhancing lesions persisting after standard treatment in glioblastoma patients: a prospective study. <i>European Radiology</i> , 2017 , 27, 3156-3166 | 8 | 23 |
| 136 | Comparison between the Prebolus T1 Measurement and the Fixed T1 Value in Dynamic Contrast-Enhanced MR Imaging for the Differentiation of True Progression from Pseudoprogession in Glioblastoma Treated with Concurrent Radiation Therapy and Temozolomide Chemotherapy. <i>American Journal of Neuroradiology</i> , 2017 , 38, 2243-2250 | 4.4 | 17 |
| 135 | Added Value of Arterial Spin-Labeling MR Imaging for the Differentiation of Cerebellar Hemangioblastoma From Metastasis. <i>American Journal of Neuroradiology</i> , 2017 , 38, 2052-2058 | 4.4 | 5 |

| | | | |
|-----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|------|
| 134 | Continuous O-Evolving MnFeO Nanoparticle-Anchored Mesoporous Silica Nanoparticles for Efficient Photodynamic Therapy in Hypoxic Cancer. <i>Journal of the American Chemical Society</i> , 2017 , 139, 10992-10995 | 16.4 | 486 |
| 133 | Iron oxide nanoclusters for T magnetic resonance imaging of non-human primates. <i>Nature Biomedical Engineering</i> , 2017 , 1, 637-643 | 19 | 117 |
| 132 | Investigation of control scans in pseudo-continuous arterial spin labeling (pCASL): Strategies for improving sensitivity and reliability of pCASL. <i>Magnetic Resonance in Medicine</i> , 2017 , 78, 917-929 | 4.4 | 8 |
| 131 | BCAT1 is a New MR Imaging-related Biomarker for Prognosis Prediction in IDH1-wildtype Glioblastoma Patients. <i>Scientific Reports</i> , 2017 , 7, 17740 | 4.9 | 13 |
| 130 | Combined use of susceptibility weighted magnetic resonance imaging sequences and dynamic susceptibility contrast perfusion weighted imaging to improve the accuracy of the differential diagnosis of recurrence and radionecrosis in high-grade glioma patients. <i>Oncotarget</i> , 2017 , 8, 20340-20353 | 3.3 | 11 |
| 129 | Electromechanical cardioplasty using a wrapped elasto-conductive epicardial mesh. <i>Science Translational Medicine</i> , 2016 , 8, 344ra86 | 17.5 | 136 |
| 128 | Antiangiogenic Effect of Bevacizumab: Application of Arterial Spin-Labeling Perfusion MR Imaging in a Rat Glioblastoma Model. <i>American Journal of Neuroradiology</i> , 2016 , 37, 1650-6 | 4.4 | 8 |
| 127 | Stretchable and Transparent Biointerface Using Cell-Sheet/Graphene Hybrid for Electrophysiology and Therapy of Skeletal Muscle. <i>Advanced Functional Materials</i> , 2016 , 26, 3207-3217 | 15.6 | 103 |
| 126 | A graphene-based electrochemical device with thermoresponsive microneedles for diabetes monitoring and therapy. <i>Nature Nanotechnology</i> , 2016 , 11, 566-572 | 28.7 | 1093 |
| 125 | Mapping blood flow directionality in the human brain. <i>Magnetic Resonance Imaging</i> , 2016 , 34, 754-764 | 3.3 | 5 |
| 124 | Assessment of bevacizumab resistance increased by expression of BCAT1 in IDH1 wild-type glioblastoma: application of DSC perfusion MR imaging. <i>Oncotarget</i> , 2016 , 7, 69606-69615 | 3.3 | 9 |
| 123 | Sclerosing Meningioma : Radiological and Clinical Characteristics of 21 Cases. <i>Journal of Korean Neurosurgical Society</i> , 2016 , 59, 584-589 | 2.3 | 2 |
| 122 | Recent Application of Advanced MR Imaging to Predict Pseudoprogression in High-grade Glioma Patients. <i>Magnetic Resonance in Medical Sciences</i> , 2016 , 15, 165-77 | 2.9 | 13 |
| 121 | Feasibility of Quantifying Arterial Cerebral Blood Volume Using Multiphase Alternate Ascending/Descending Directional Navigation (ALADDIN). <i>PLoS ONE</i> , 2016 , 11, e0156687 | 3.7 | 6 |
| 120 | Application of diffusion-weighted imaging and dynamic susceptibility contrast perfusion-weighted imaging for ganglioglioma in adults: Comparison study with oligodendroglioma. <i>Journal of Neuroradiology</i> , 2016 , 43, 331-8 | 3.1 | 8 |
| 119 | Glioblastoma treated with concurrent radiation therapy and temozolomide chemotherapy: differentiation of true progression from pseudoprogression with quantitative dynamic contrast-enhanced MR imaging. <i>Radiology</i> , 2015 , 274, 830-40 | 20.5 | 83 |
| 118 | Evaluation of the microenvironmental heterogeneity in high-grade gliomas with IDH1/2 gene mutation using histogram analysis of diffusion-weighted imaging and dynamic-susceptibility contrast perfusion imaging. <i>Journal of Neuro-Oncology</i> , 2015 , 121, 141-50 | 4.8 | 75 |
| 117 | Bioresorbable Electronic Stent Integrated with Therapeutic Nanoparticles for Endovascular Diseases. <i>ACS Nano</i> , 2015 , 9, 5937-46 | 16.7 | 158 |

| | | | |
|-----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----|
| 116 | Differentiation of Parkinsonism-Predominant Multiple System Atrophy from Idiopathic Parkinson Disease Using 3T Susceptibility-Weighted MR Imaging, Focusing on Putaminal Change and Lesion Asymmetry. <i>American Journal of Neuroradiology</i> , 2015 , 36, 2227-34 | 4.4 | 23 |
| 115 | Paradoxical perfusion metrics of high-grade gliomas with an oligodendroglioma component: quantitative analysis of dynamic susceptibility contrast perfusion MR imaging. <i>Neuroradiology</i> , 2015 , 57, 1111-20 | 3.2 | 7 |
| 114 | Hollow MnOxPy and Pt/MnOxPy yolk/shell nanoparticles as a T1 MRI contrast agent. <i>Journal of Colloid and Interface Science</i> , 2015 , 439, 134-8 | 9.3 | 6 |
| 113 | Hypofractionated chemoradiotherapy with temozolomide as a treatment option for glioblastoma patients with poor prognostic features. <i>International Journal of Clinical Oncology</i> , 2015 , 20, 21-8 | 4.2 | 7 |
| 112 | An NMR metabolomics approach for the diagnosis of leptomeningeal carcinomatosis in lung adenocarcinoma cancer patients. <i>International Journal of Cancer</i> , 2015 , 136, 162-71 | 7.5 | 32 |
| 111 | MRI molecular imaging using GLUT1 antibody-Fe ₃ O ₄ nanoparticles in the hemangioma animal model for differentiating infantile hemangioma from vascular malformation. <i>Nanomedicine: Nanotechnology, Biology, and Medicine</i> , 2015 , 11, 127-35 | 6 | 19 |
| 110 | IDH2 mutation in gliomas including novel mutation. <i>Neuropathology</i> , 2015 , 35, 236-44 | 2 | 16 |
| 109 | Physiological and Functional Magnetic Resonance Imaging Using Balanced Steady-state Free Precession. <i>Korean Journal of Radiology</i> , 2015 , 16, 550-9 | 6.9 | 20 |
| 108 | Investigation of inter-slice magnetization transfer effects as a new method for MTR imaging of the human brain. <i>PLoS ONE</i> , 2015 , 10, e0117101 | 3.7 | 9 |
| 107 | Inter-Slice Blood Flow and Magnetization Transfer Effects as A New Simultaneous Imaging Strategy. <i>PLoS ONE</i> , 2015 , 10, e0140560 | 3.7 | 10 |
| 106 | Squamous Cell Carcinoma of the Head and Neck: Comparison of Diffusion-weighted MRI at b-values of 1,000 and 2,000 s/mm ² to Predict Response to Induction Chemotherapy. <i>Magnetic Resonance in Medical Sciences</i> , 2015 , 14, 337-45 | 2.9 | 12 |
| 105 | Prediction of Response to Concurrent Chemoradiotherapy with Temozolomide in Glioblastoma: Application of Immediate Post-Operative Dynamic Susceptibility Contrast and Diffusion-Weighted MR Imaging. <i>Korean Journal of Radiology</i> , 2015 , 16, 1341-8 | 6.9 | 14 |
| 104 | Bright vessel appearance on arterial spin labeling MRI for localizing arterial occlusion in acute ischemic stroke. <i>Stroke</i> , 2015 , 46, 564-7 | 6.7 | 37 |
| 103 | An endoscope with integrated transparent bioelectronics and theranostic nanoparticles for colon cancer treatment. <i>Nature Communications</i> , 2015 , 6, 10059 | 17.4 | 122 |
| 102 | Polymeric embolization coil of bilayered polyvinyl alcohol strand for therapeutic vascular occlusion: a feasibility study in canine experimental vascular models. <i>Journal of Vascular and Interventional Radiology</i> , 2015 , 26, 117-23 | 2.4 | 2 |
| 101 | Multifunctional cell-culture platform for aligned cell sheet monitoring, transfer printing, and therapy. <i>ACS Nano</i> , 2015 , 9, 2677-88 | 16.7 | 58 |
| 100 | Metabolomic comparison between cells over-expressing isocitrate dehydrogenase 1 and 2 mutants and the effects of an inhibitor on the metabolism. <i>Journal of Neurochemistry</i> , 2015 , 132, 183-93 | 6 | 13 |
| 99 | Organized hematoma developed after suboccipital craniectomy. <i>Journal of Neuroimaging</i> , 2014 , 24, 610-612 | 6.82 | 1 |

| | | | |
|----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----|
| 98 | Contrast-enhanced FLAIR (fluid-attenuated inversion recovery) for evaluating mild traumatic brain injury. <i>PLoS ONE</i> , 2014 , 9, e102229 | 3.7 | 13 |
| 97 | Prognosis prediction of measurable enhancing lesion after completion of standard concomitant chemoradiotherapy and adjuvant temozolomide in glioblastoma patients: application of dynamic susceptibility contrast perfusion and diffusion-weighted imaging. <i>PLoS ONE</i> , 2014 , 9, e113587 | 3.7 | 12 |
| 96 | A case report of preoperative and postoperative 7.0T brain MRI in a patient with a small cell glioblastoma. <i>Journal of Korean Medical Science</i> , 2014 , 29, 1012-7 | 4.7 | 0 |
| 95 | Prognosis of Glioblastoma With Oligodendroglioma Component is Associated With the IDH1 Mutation and MGMT Methylation Status. <i>Translational Oncology</i> , 2014 , 7, 712-9 | 4.9 | 8 |
| 94 | Stretchable silicon nanoribbon electronics for skin prosthesis. <i>Nature Communications</i> , 2014 , 5, 5747 | 17.4 | 902 |
| 93 | Malignant glioma: MR imaging by using 5-aminolevulinic acid in an animal model. <i>Radiology</i> , 2014 , 272, 720-30 | 20.5 | 12 |
| 92 | Glioma grading using apparent diffusion coefficient map: application of histogram analysis based on automatic segmentation. <i>NMR in Biomedicine</i> , 2014 , 27, 1046-52 | 4.4 | 24 |
| 91 | Correlation of 11C-methionine PET and diffusion-weighted MRI: is there a complementary diagnostic role for gliomas?. <i>Nuclear Medicine Communications</i> , 2014 , 35, 720-6 | 1.6 | 12 |
| 90 | Segmentation-based MR attenuation correction including bones also affects quantitation in brain studies: an initial result of 18F-FP-CIT PET/MR for patients with parkinsonism. <i>Journal of Nuclear Medicine</i> , 2014 , 55, 1617-22 | 8.9 | 21 |
| 89 | Grading of cerebral glioma with multiparametric MR imaging and 18F-FDG-PET: concordance and accuracy. <i>European Radiology</i> , 2014 , 24, 380-9 | 8 | 46 |
| 88 | Glioma: application of whole-tumor texture analysis of diffusion-weighted imaging for the evaluation of tumor heterogeneity. <i>PLoS ONE</i> , 2014 , 9, e108335 | 3.7 | 132 |
| 87 | Clinicopathological and genetic characteristics of extraventricular neurocytomas. <i>Neuropathology</i> , 2013 , 33, 111-21 | 2 | 27 |
| 86 | Primary intracerebral malignant fibrous histiocytoma: CT, MRI, and PET-CT findings. <i>Journal of Neuroimaging</i> , 2013 , 23, 141-4 | 2.8 | 8 |
| 85 | Seizures during the management of high-grade gliomas: clinical relevance to disease progression. <i>Journal of Neuro-Oncology</i> , 2013 , 113, 101-9 | 4.8 | 19 |
| 84 | Early response evaluation for recurrent high grade gliomas treated with bevacizumab: a volumetric analysis using diffusion-weighted imaging. <i>Journal of Neuro-Oncology</i> , 2013 , 112, 427-35 | 4.8 | 13 |
| 83 | The value of temozolomide in combination with radiotherapy during standard treatment for newly diagnosed glioblastoma. <i>Journal of Neuro-Oncology</i> , 2013 , 112, 277-83 | 4.8 | 18 |
| 82 | Effect of carotid artery stenting on cerebral blood flow: evaluation of hemodynamic changes using arterial spin labeling. <i>Neuroradiology</i> , 2013 , 55, 271-81 | 3.2 | 19 |
| 81 | Desmoid-type fibromatosis in the head and neck: CT and MR imaging characteristics. <i>Neuroradiology</i> , 2013 , 55, 351-9 | 3.2 | 26 |

| | | | |
|----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----|
| 80 | A practical scoring system to determine whether to proceed with surgical resection in recurrent glioblastoma. <i>Neuro-Oncology</i> , 2013 , 15, 1096-101 | 1 | 48 |
| 79 | Radiosurgery for central neurocytoma: long-term outcome and failure pattern. <i>Journal of Neuro-Oncology</i> , 2013 , 115, 505-11 | 4.8 | 22 |
| 78 | Tumor blood flow from arterial spin labeling perfusion MRI: a key parameter in distinguishing high-grade gliomas from primary cerebral lymphomas, and in predicting genetic biomarkers in high-grade gliomas. <i>Journal of Magnetic Resonance Imaging</i> , 2013 , 38, 852-60 | 5.6 | 34 |
| 77 | In vivo magnetic resonance imaging of transgenic mice expressing human ferritin. <i>Molecular Imaging and Biology</i> , 2013 , 15, 48-57 | 3.8 | 14 |
| 76 | Correlation of apparent diffusion coefficient values measured by diffusion MRI and MGMT promoter methylation semiquantitatively analyzed with MS-MLPA in patients with glioblastoma multiforme. <i>Journal of Magnetic Resonance Imaging</i> , 2013 , 37, 351-8 | 5.6 | 33 |
| 75 | Nano-sized CT contrast agents. <i>Advanced Materials</i> , 2013 , 25, 2641-60 | 24 | 411 |
| 74 | Body fat assessment method using CT images with separation mask algorithm. <i>Journal of Digital Imaging</i> , 2013 , 26, 155-62 | 5.3 | 32 |
| 73 | A highly facile and specific assay for cancer-causing isocitrate dehydrogenase mutant using ¹³ C4-labeled β -ketoglutarate and heteronuclear NMR. <i>Analytical Chemistry</i> , 2013 , 85, 11987-92 | 7.8 | 3 |
| 72 | Shoulder sonography after intraarticular fluid injection for evaluation of anterior labral tears: comparison with conventional sonography. <i>Journal of Clinical Ultrasound</i> , 2013 , 41, 94-100 | 1 | 4 |
| 71 | Effect of delayed transit time on arterial spin labeling: correlation with dynamic susceptibility contrast perfusion magnetic resonance in moyamoya disease. <i>Investigative Radiology</i> , 2013 , 48, 795-802 | 10.1 | 23 |
| 70 | Differentiation of true progression from pseudoprogression in glioblastoma treated with radiation therapy and concomitant temozolomide: comparison study of standard and high-b-value diffusion-weighted imaging. <i>Radiology</i> , 2013 , 269, 831-40 | 20.5 | 124 |
| 69 | Benign lymphoepithelial tumor of the pituitary. <i>Neuropathology</i> , 2013 , 33, 413-7 | 2 | 1 |
| 68 | Central neurocytoma: long-term outcomes of multimodal treatments and management strategies based on 30 years experience in a single institute. <i>Neurosurgery</i> , 2013 , 72, 407-13; discussion 413-4 | 3.2 | 32 |
| 67 | Hyperfunction thyroid nodules: their risk for becoming or being associated with thyroid cancers. <i>Korean Journal of Radiology</i> , 2013 , 14, 643-52 | 6.9 | 13 |
| 66 | Early experience of pre- and post-contrast 7.0T MRI in brain tumors. <i>Journal of Korean Medical Science</i> , 2013 , 28, 1362-72 | 4.7 | 15 |
| 65 | Serial MR analysis of early permanent and transient ischemia in rats: diffusion tensor imaging and high b value diffusion weighted imaging. <i>Korean Journal of Radiology</i> , 2013 , 14, 307-15 | 6.9 | 11 |
| 64 | Noninvasive identification of viable cell populations in docetaxel-treated breast tumors using ferritin-based magnetic resonance imaging. <i>PLoS ONE</i> , 2013 , 8, e52931 | 3.7 | 16 |
| 63 | Gliomas: application of cumulative histogram analysis of normalized cerebral blood volume on 3 T MRI to tumor grading. <i>PLoS ONE</i> , 2013 , 8, e63462 | 3.7 | 32 |

| | | | |
|----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----|
| 62 | Cerebral blood volume analysis in glioblastomas using dynamic susceptibility contrast-enhanced perfusion MRI: a comparison of manual and semiautomatic segmentation methods. <i>PLoS ONE</i> , 2013 , 8, e69323 | 3.7 | 36 |
| 61 | True progression versus pseudoprogression in the treatment of glioblastomas: a comparison study of normalized cerebral blood volume and apparent diffusion coefficient by histogram analysis. <i>Korean Journal of Radiology</i> , 2013 , 14, 662-72 | 6.9 | 68 |
| 60 | Hydrotropic magnetic micelles for combined magnetic resonance imaging and cancer therapy. <i>Journal of Controlled Release</i> , 2012 , 160, 692-8 | 11.7 | 32 |
| 59 | Imaging and quantification of metastatic melanoma cells in lymph nodes with a ferritin MR reporter in living mice. <i>NMR in Biomedicine</i> , 2012 , 25, 737-45 | 4.4 | 26 |
| 58 | Augmentation of chemotherapeutic infusion effect by TSU-68, an oral targeted antiangiogenic agent, in a rabbit VX2 liver tumor model. <i>CardioVascular and Interventional Radiology</i> , 2012 , 35, 168-75 | 2.7 | 3 |
| 57 | MR imaging findings of extraventricular neurocytoma: a series of ten patients confirmed by immunohistochemistry of IDH1 gene mutation. <i>Acta Neurochirurgica</i> , 2012 , 154, 1973-79; discussion 1980 | | 19 |
| 56 | Histogram analysis of apparent diffusion coefficient map of standard and high B-value diffusion MR imaging in head and neck squamous cell carcinoma: a correlation study with histological grade. <i>Academic Radiology</i> , 2012 , 19, 1233-40 | 4.3 | 48 |
| 55 | The Changes in MGMT Promoter Methylation Status in Initial and Recurrent Glioblastomas. <i>Translational Oncology</i> , 2012 , 5, 393-7 | 4.9 | 34 |
| 54 | Protective effects of cleavage agents on INS-1 cells against h-IAPP-induced apoptosis. <i>Chemical Communications</i> , 2012 , 48, 588-90 | 5.8 | 6 |
| 53 | Diffusion-weighted MR imaging for the differentiation of true progression from pseudoprogression following concomitant radiotherapy with temozolomide in patients with newly diagnosed high-grade gliomas. <i>Academic Radiology</i> , 2012 , 19, 1353-61 | 4.3 | 79 |
| 52 | An NMR metabolomics approach for the diagnosis of leptomeningeal carcinomatosis. <i>Cancer Research</i> , 2012 , 72, 5179-87 | 10.1 | 14 |
| 51 | Magnetic resonance enhancement pattern and diagnostic accuracy of gadofluorine M in a rabbit VX2 tumor model: Comparison with gadopentetate dimeglumine. <i>European Journal of Radiology</i> , 2012 , 81, 1751-7 | 4.7 | |
| 50 | Theranostic probe based on lanthanide-doped nanoparticles for simultaneous in vivo dual-modal imaging and photodynamic therapy. <i>Advanced Materials</i> , 2012 , 24, 5755-61 | 24 | 334 |
| 49 | Macrophages homing to metastatic lymph nodes can be monitored with ultrasensitive ferromagnetic iron-oxide nanocubes and a 1.5T clinical MR scanner. <i>PLoS ONE</i> , 2012 , 7, e29575 | 3.7 | 10 |
| 48 | Evaluation of lymph node metastases: comparison of gadofluorine M-enhanced MRI and diffusion-weighted MRI in a rabbit VX2 rectal cancer model. <i>Journal of Magnetic Resonance Imaging</i> , 2012 , 35, 1179-86 | 5.6 | 5 |
| 47 | Synthesis of Uniformly Sized Manganese Oxide Nanocrystals with Various Sizes and Shapes and Characterization of Their T1 Magnetic Resonance Relaxivity. <i>European Journal of Inorganic Chemistry</i> , 2012 , 2012, 2148-2155 | 2.3 | 62 |
| 46 | Water-dispersible ferrimagnetic iron oxide nanocubes with extremely high r ₁ relaxivity for highly sensitive in vivo MRI of tumors. <i>Nano Letters</i> , 2012 , 12, 3127-31 | 11.5 | 238 |
| 45 | Multifunctional Fe ₃ O ₄ /TaO(x) core/shell nanoparticles for simultaneous magnetic resonance imaging and X-ray computed tomography. <i>Journal of the American Chemical Society</i> , 2012 , 134, 10309-12 | 16.4 | 193 |

| | | | |
|----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----|
| 44 | Development of endovascular vibrating polymer actuator probe for mechanical thrombolysis: in vivo study. <i>ASAIO Journal</i> , 2012 , 58, 503-8 | 3.6 | 2 |
| 43 | Lymph node metastases from gastric cancer: gadofluorine M and gadopentetate dimeglumine MR imaging in a rabbit model. <i>Radiology</i> , 2012 , 263, 391-400 | 20.5 | 9 |
| 42 | Computational analysis of blood clot dissolution using a vibrating catheter tip. <i>Proceedings of the Institution of Mechanical Engineers, Part H: Journal of Engineering in Medicine</i> , 2012 , 226, 337-40 | 1.7 | |
| 41 | Comparison between transauricular and transfemoral arterial access for hepatic artery angiography in a rabbit model. <i>Journal of Vascular and Interventional Radiology</i> , 2011 , 22, 1181-7 | 2.4 | 12 |
| 40 | Transformation of hydrophobic iron oxide nanoparticles to hydrophilic and biocompatible maghemite nanocrystals for use as highly efficient MRI contrast agent. <i>Journal of Materials Chemistry</i> , 2011 , 21, 11472 | | 42 |
| 39 | Large-Scale Synthesis of Ultrathin Manganese Oxide Nanoplates and Their Applications to T1 MRI Contrast Agents. <i>Chemistry of Materials</i> , 2011 , 23, 3318-3324 | 9.6 | 83 |
| 38 | Large-scale synthesis of uniform and extremely small-sized iron oxide nanoparticles for high-resolution T1 magnetic resonance imaging contrast agents. <i>Journal of the American Chemical Society</i> , 2011 , 133, 12624-31 | 16.4 | 691 |
| 37 | Development of endovascular vibrating polymer actuator probe for mechanical thrombolysis: a phantom study. <i>ASAIO Journal</i> , 2011 , 57, 286-92 | 3.6 | 1 |
| 36 | Multiple-interaction ligands inspired by mussel adhesive protein: synthesis of highly stable and biocompatible nanoparticles. <i>Angewandte Chemie - International Edition</i> , 2011 , 50, 11360-5 | 16.4 | 108 |
| 35 | Multifunctional mesoporous silica nanocomposite nanoparticles for pH controlled drug release and dual modal imaging. <i>Journal of Materials Chemistry</i> , 2011 , 21, 16869 | | 69 |
| 34 | Large-scale synthesis of bioinert tantalum oxide nanoparticles for X-ray computed tomography imaging and bimodal image-guided sentinel lymph node mapping. <i>Journal of the American Chemical Society</i> , 2011 , 133, 5508-15 | 16.4 | 270 |
| 33 | Gliomas: Histogram analysis of apparent diffusion coefficient maps with standard- or high-b-value diffusion-weighted MR imaging--correlation with tumor grade. <i>Radiology</i> , 2011 , 261, 882-90 | 20.5 | 248 |
| 32 | Clinical and radiological features of pandemic H1N1 2009 influenza virus infection manifesting as acute febrile respiratory illness at their initial presentations: comparison with contemporaneous non-H1N1 patients. <i>Acta Radiologica</i> , 2011 , 52, 410-6 | 2 | 7 |
| 31 | Correlation of 18F-FDG uptake with apparent diffusion coefficient ratio measured on standard and high b value diffusion MRI in head and neck cancer. <i>Journal of Nuclear Medicine</i> , 2011 , 52, 1056-62 | 8.9 | 46 |
| 30 | Brain death: evaluation of cerebral blood flow by use of arterial spin labeling. <i>Circulation</i> , 2011 , 124, 2572-3 | 16.7 | 7 |
| 29 | Usefulness of MS-MLPA for detection of MGMT promoter methylation in the evaluation of pseudoprogression in glioblastoma patients. <i>Neuro-Oncology</i> , 2011 , 13, 195-202 | 1 | 47 |
| 28 | Magnetosome-like ferrimagnetic iron oxide nanocubes for highly sensitive MRI of single cells and transplanted pancreatic islets. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011 , 108, 2662-7 | 11.5 | 166 |
| 27 | The role of perfusion CT as a follow-up modality after transcatheter arterial chemoembolization: an experimental study in a rabbit model. <i>Investigative Radiology</i> , 2010 , 45, 427-36 | 10.1 | 51 |

| | | | |
|----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----|
| 26 | Contrast-enhanced MR imaging of lymph nodes in cancer patients. <i>Korean Journal of Radiology</i> , 2010 , 11, 383-94 | 6.9 | 19 |
| 25 | Radiological and clinical characteristics of a military outbreak of pandemic H1N1 2009 influenza virus infection. <i>Korean Journal of Radiology</i> , 2010 , 11, 417-24 | 6.9 | 9 |
| 24 | In vivo imaging of tumor transduced with bimodal lentiviral vector encoding human ferritin and green fluorescent protein on a 1.5T clinical magnetic resonance scanner. <i>Cancer Research</i> , 2010 , 70, 7315-24 | 19.1 | 56 |
| 23 | Uniform mesoporous dye-doped silica nanoparticles decorated with multiple magnetite nanocrystals for simultaneous enhanced magnetic resonance imaging, fluorescence imaging, and drug delivery. <i>Journal of the American Chemical Society</i> , 2010 , 132, 552-7 | 16.4 | 645 |
| 22 | Comparison of lymph node metastases assessment with the use of USPIO-enhanced MR imaging at 1.5 T versus 3.0 T in a rabbit model. <i>Journal of Magnetic Resonance Imaging</i> , 2010 , 31, 134-41 | 5.6 | 8 |
| 21 | MRI of magnetically labeled mesenchymal stem cells in hepatic failure model. <i>World Journal of Gastroenterology</i> , 2010 , 16, 5611-5 | 5.6 | 6 |
| 20 | Nonblinking and Nonbleaching Upconverting Nanoparticles as an Optical Imaging Nanoprobe and T1 Magnetic Resonance Imaging Contrast Agent. <i>Advanced Materials</i> , 2009 , 21, 4467-4471 | 24 | 501 |
| 19 | Bandgap engineered reverse type-I CdTe/InP/ZnS core-shell nanocrystals for the near-infrared. <i>Chemical Communications</i> , 2009 , 1267-9 | 5.8 | 25 |
| 18 | The efficacy of magnetic resonance imaging for the diagnosis of testicular rupture: a prospective preliminary study. <i>Journal of Trauma</i> , 2009 , 66, 239-42 | | 32 |
| 17 | Enhancement characteristics of cholangiocarcinomas on multiphasic helical CT: emphasis on morphologic subtypes. <i>Clinical Imaging</i> , 2008 , 32, 114-20 | 2.7 | 25 |
| 16 | Hepatocellular carcinoma in liver transplantation candidates: detection with gadobenate dimeglumine-enhanced MRI. <i>American Journal of Roentgenology</i> , 2008 , 191, 529-36 | 5.4 | 78 |
| 15 | Lymph node metastasis: ultrasmall superparamagnetic iron oxide-enhanced MR imaging versus PET/CT in a rabbit model. <i>Radiology</i> , 2007 , 242, 137-43 | 20.5 | 32 |
| 14 | Hepatocellular carcinoma with internal mammary artery supply: feasibility and efficacy of transarterial chemoembolization and factors affecting patient prognosis. <i>Journal of Vascular and Interventional Radiology</i> , 2007 , 18, 611-9; quiz 620 | 2.4 | 27 |
| 13 | Assessment of lymph node metastases by contrast-enhanced MR imaging in a head and neck cancer model. <i>Korean Journal of Radiology</i> , 2007 , 8, 9-14 | 6.9 | 11 |
| 12 | Internal mammary arteries supplying hepatocellular carcinoma: vascular anatomy at digital subtraction angiography in 97 patients. <i>Radiology</i> , 2007 , 242, 925-32 | 20.5 | 30 |
| 11 | Cervical lymph node metastases: MR imaging of gadofluorine M and monocrySTALLINE iron oxide nanoparticle-47 in a rabbit model of head and neck cancer. <i>Radiology</i> , 2006 , 241, 753-62 | 20.5 | 25 |
| 10 | CT Differentiation of cholangiocarcinoma from periductal fibrosis in patients with hepatolithiasis. <i>American Journal of Roentgenology</i> , 2006 , 187, 445-53 | 5.4 | 39 |
| 9 | Hepatic radiofrequency ablation using multiple probes: ex vivo and in vivo comparative studies of monopolar versus multipolar modes. <i>Korean Journal of Radiology</i> , 2006 , 7, 106-17 | 6.9 | 24 |

| | | | |
|---|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|----|
| 8 | Computed tomography features of an intraductal polypoid mass: differentiation between hepatocellular carcinoma with bile duct tumor invasion and intraductal papillary cholangiocarcinoma. <i>Journal of Computer Assisted Tomography</i> , 2006 , 30, 18-24 | 2.2 | 5 |
| 7 | CT features of an intraductal polypoid mass: Differentiation between hepatocellular carcinoma with bile duct tumor invasion and intraductal papillary cholangiocarcinoma. <i>Journal of Computer Assisted Tomography</i> , 2006 , 30, 173-81 | 2.2 | 30 |
| 6 | Bipolar radiofrequency ablation in ex vivo bovine liver with the open-perfused system versus the cooled-wet system. <i>European Radiology</i> , 2005 , 15, 759-64 | 8 | 26 |
| 5 | Differentiating malignant from benign common bile duct stricture with multiphasic helical CT. <i>Radiology</i> , 2005 , 236, 178-83 | 20.5 | 92 |
| 4 | Postbiopsy splenic bleeding in a dog model: comparison of cauterization, embolization, and plugging of the needle tract. <i>American Journal of Roentgenology</i> , 2005 , 185, 878-84 | 5.4 | 20 |
| 3 | Endobronchial metastasis from renal cell carcinoma: CT findings in four patients. <i>European Journal of Radiology</i> , 2004 , 51, 155-9 | 4.7 | 17 |
| 2 | Imaging of gastrointestinal stromal tumors. <i>Journal of Computer Assisted Tomography</i> , 2004 , 28, 596-604 | 2.2 | 24 |
| 1 | Preoperative magnetic resonance imaging staging of uterine cervical carcinoma: results of prospective study. <i>Journal of Computer Assisted Tomography</i> , 2004 , 28, 620-7 | 2.2 | 71 |