

Ricardo Araya

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2595917/publications.pdf>

Version: 2024-02-01

268
papers

18,935
citations

13865

67
h-index

15732

125
g-index

290
all docs

290
docs citations

290
times ranked

19190
citing authors

#	ARTICLE	IF	CITATIONS
1	How much change is enough? Evidence from a longitudinal study on depression in UK primary care. <i>Psychological Medicine</i> , 2022, 52, 1875-1882.	4.5	51
2	Psychometric properties of the PHQ-9 measure of depression among Brazilian older adults. <i>Aging and Mental Health</i> , 2022, 26, 2285-2290.	2.8	10
3	Economic threshold analysis of delivering a task-sharing treatment for common mental disorders at scale: the Friendship Bench, Zimbabwe. <i>Evidence-Based Mental Health</i> , 2022, 25, 47-53.	4.5	2
4	Depressive and subthreshold depressive symptomatology among older adults in a socioeconomically deprived area in Brazil. <i>International Journal of Geriatric Psychiatry</i> , 2022, 37, .	2.7	5
5	Effect of Sleep Disturbance Symptoms on Treatment Outcome in Blended Cognitive Behavioral Therapy for Depression (E-COMPARED Study): Secondary Analysis. <i>Journal of Medical Internet Research</i> , 2022, 24, e30231.	4.3	2
6	HeAlth System StrEngThening in four sub-Saharan African countries (ASSET) to achieve high-quality, evidence-informed surgical, maternal and newborn, and primary care: protocol for pre-implementation phase studies. <i>Global Health Action</i> , 2022, 15, 1987044.	1.9	11
7	The relationship between multidimensional poverty, income poverty and youth depressive symptoms: cross-sectional evidence from Mexico, South Africa and Colombia. <i>BMJ Global Health</i> , 2022, 7, e006960.	4.7	8
8	Burden of HIV-related stigma and associated factors among women living with depression accessing PMTCT services in Dar es Salaam, Tanzania. <i>AIDS Care - Psychological and Socio-Medical Aspects of AIDS/HIV</i> , 2022, 34, 1572-1579.	1.2	1
9	A Data-Driven Clustering Method for Discovering Profiles in the Dynamics of Major Depressive Disorder Using a Smartphone-Based Ecological Momentary Assessment of Mood. <i>Frontiers in Psychiatry</i> , 2022, 13, 755809.	2.6	1
10	A Validation Study of the EQ-5D-5L and ICEpop Capability Measure for Older People Among Older Individuals With Depressive Symptoms in Brazil. <i>Value in Health Regional Issues</i> , 2022, 30, 91-99.	1.2	2
11	Association of Task-Shared Psychological Interventions With Depression Outcomes in Low- and Middle-Income Countries. <i>JAMA Psychiatry</i> , 2022, 79, 430.	11.0	22
12	Relationship between the Bolsa Família national cash transfer programme and suicide incidence in Brazil: A quasi-experimental study. <i>PLoS Medicine</i> , 2022, 19, e1004000.	8.4	9
13	Adolescent Lifestyle Profile-Revised 2: validity and reliability among adolescents in Chile. <i>Jornal De Pediatria</i> , 2021, 97, 52-60.	2.0	7
14	The emergence of digital mental health in low-income and middle-income countries: A review of recent advances and implications for the treatment and prevention of mental disorders. <i>Journal of Psychiatric Research</i> , 2021, 133, 223-246.	3.1	71
15	Recruitment, training and supervision of nurses and nurse assistants for a task-shifting depression intervention in two RCTs in Brazil and Peru. <i>Human Resources for Health</i> , 2021, 19, 16.	3.1	7
16	Integrating youth mental health into cash transfer programmes in response to the COVID-19 crisis in low-income and middle-income countries. <i>Lancet Psychiatry</i> , 2021, 8, 340-346.	7.4	45
17	Validation of the Revised Olweus Bully/Victim Questionnaire (OBVQ-R) Among Adolescents in Chile. <i>Frontiers in Psychology</i> , 2021, 12, 578661.	2.1	18
18	Brief Research Report: The Association Between Educational Experiences and Covid-19 Pandemic-Related Variables, and Mental Health Among Children and Adolescents. <i>Frontiers in Psychiatry</i> , 2021, 12, 647456.	2.6	8

#	ARTICLE	IF	CITATIONS
19	Effect of a Digital Intervention on Depressive Symptoms in Patients With Comorbid Hypertension or Diabetes in Brazil and Peru. JAMA - Journal of the American Medical Association, 2021, 325, 1852.	7.4	40
20	Effectiveness and cost-effectiveness of integrating the management of depression into routine HIV Care in Uganda (the HIV+â€”D trial): A protocol for a cluster-randomised trial. International Journal of Mental Health Systems, 2021, 15, 45.	2.7	4
21	Examining the dynamics between young peopleâ€™s mental health, poverty and life chances in six low- and middle-income countries: protocol for the CHANCES-6 study. Social Psychiatry and Psychiatric Epidemiology, 2021, 56, 1687-1703.	3.1	4
22	Implementation science protocol for a participatory, theory-informed implementation research programme in the context of health system strengthening in sub-Saharan Africa (ASSET-ImplementER). BMJ Open, 2021, 11, e048742.	1.9	8
23	A collaborative, computer-assisted, psycho-educational intervention for depressed patients with chronic disease at primary care: protocol for a cluster randomized controlled trial. BMC Psychiatry, 2021, 21, 418.	2.6	4
24	Optimising implementation strategies of the first scaleup of a primary care psychological intervention for common mental disorders in Sub-Saharan Africa: a mixed methods study protocol for the optimised Friendship Bench (OptFB). BMJ Open, 2021, 11, e045481.	1.9	3
25	Psychometric Evaluation of the Health State Description Questionnaire in Chile: A Proposal for a Latent Variable Approach for Valuating Health States. Value in Health Regional Issues, 2021, 26, 142-149.	1.2	1
26	Global Mental Health and Nutrition: Moving Toward a Convergent Research Agenda. Frontiers in Public Health, 2021, 9, 722290.	2.7	8
27	A Mobile Health Intervention for Patients With Depressive Symptoms: Protocol for an Economic Evaluation Alongside Two Randomized Trials in Brazil and Peru. JMIR Research Protocols, 2021, 10, e26164.	1.0	0
28	Conditional cash transfers and adolescent mental health in Brazil: Evidence from the 2004 Pelotas Birth Cohort. Journal of Global Health, 2021, 11, 04066.	2.7	6
29	An intervention to improve teacher well-being support and training to support students in UK high schools (the WISE study): A cluster randomised controlled trial. PLoS Medicine, 2021, 18, e1003847.	8.4	12
30	Mental health support and training to improve secondary school teachersâ€™ well-being: the WISE cluster RCT. Public Health Research, 2021, 9, 1-138.	1.3	2
31	A guide to systems-level, participatory, theory-informed implementation research in global health. BMJ Global Health, 2021, 6, e005365.	4.7	15
32	Acceptability and fidelity of a psychosocial intervention (PROACTIVE) for older adults with depression in a basic health unit in SÃ£o Paulo, Brazil: a qualitative study. BMC Public Health, 2021, 21, 2278.	2.9	3
33	Examining the Theoretical Framework of Behavioral Activation for Major Depressive Disorder: Smartphone-Based Ecological Momentary Assessment Study. JMIR Mental Health, 2021, 8, e32007.	3.3	5
34	Effect of a differentiated service delivery model on virological failure in adolescents with HIV in Zimbabwe (Zvandiri): a cluster-randomised controlled trial. The Lancet Global Health, 2020, 8, e264-e275.	6.3	113
35	Cost-Effectiveness of Sertraline in Primary Care According to Initial Severity and Duration of Depressive Symptoms: Findings from the PANDA RCT. Pharmacoeconomics - Open, 2020, 4, 427-438.	1.8	3
36	A collaborative care psychosocial intervention to improve late life depression in socioeconomically deprived areas of Guarulhos, Brazil: the PROACTIVE cluster randomised controlled trial protocol. Trials, 2020, 21, 914.	1.6	19

#	ARTICLE	IF	CITATIONS
37	Towards a conceptual framework of the working alliance in a blended low-intensity cognitive behavioural therapy intervention for depression in primary mental health care: a qualitative study. <i>BMJ Open</i> , 2020, 10, e036299.	1.9	9
38	Effectiveness of clinical decision support systems and telemedicine on outcomes of depression: a cluster randomized trial in general practice. <i>Family Practice</i> , 2020, 37, 731-737.	1.9	12
39	Delivery of a Mental Health First Aid training package and staff peer support service in secondary schools: a process evaluation of uptake and fidelity of the WISE intervention. <i>Trials</i> , 2020, 21, 745.	1.6	5
40	The effect of co-morbid anxiety on remission from depression for people participating in a randomised controlled trial of the Friendship Bench intervention in Zimbabwe. <i>EClinicalMedicine</i> , 2020, 23, 100333.	7.1	6
41	Major Depressive Disorder Among HIV Infected Youth in Uganda: Incidence, Persistence and Their Predictors. <i>AIDS and Behavior</i> , 2020, 24, 2588-2596.	2.7	6
42	Added value or added burden? A qualitative investigation of blending internet self-help with face-to-face cognitive behaviour therapy for depression. <i>Psychotherapy Research</i> , 2020, 30, 998-1010.	1.8	21
43	Healthy Options: study protocol and baseline characteristics for a cluster randomized controlled trial of group psychotherapy for perinatal women living with HIV and depression in Tanzania. <i>BMC Public Health</i> , 2020, 20, 80.	2.9	12
44	Mental health research capacity building in sub-Saharan Africa: the African Mental Health Research Initiative. <i>Global Mental Health (Cambridge, England)</i> , 2020, 7, e8.	2.5	16
45	Building a Platform for Translational Research in Chronic Noncommunicable Diseases to Address Population Health: Lessons From NHLBI Supported CRONICAS in Peru. <i>Global Heart</i> , 2020, 10, 13.	2.3	4
46	Is teachers' mental health and wellbeing associated with students' mental health and wellbeing?. <i>Journal of Affective Disorders</i> , 2019, 242, 180-187.	4.1	155
47	Computer-Assisted Cognitive-Behavioral Therapy to Treat Adolescents With Depression in Primary Health Care Centers in Santiago, Chile: A Randomized Controlled Trial. <i>Frontiers in Psychiatry</i> , 2019, 10, 552.	2.6	14
48	Pre- and post-natal maternal anxiety and early childhood weight gain. <i>Journal of Affective Disorders</i> , 2019, 257, 136-142.	4.1	9
49	Exploring Differences Between Adolescents and Adults With Perinatal Depression" Data From the Expanding Care for Perinatal Women With Depression Trial in Nigeria. <i>Frontiers in Psychiatry</i> , 2019, 10, 761.	2.6	15
50	Internet-Based Interventions for the Prevention and Treatment of Mental Disorders in Latin America: A Scoping Review. <i>Frontiers in Psychiatry</i> , 2019, 10, 664.	2.6	54
51	Pilot study of a two-arm non-randomized controlled cluster trial of a psychosocial intervention to improve late life depression in socioeconomically deprived areas of São Paulo, Brazil (PROACTIVE): feasibility study of a psychosocial intervention for late life depression in São Paulo. <i>BMC Public Health</i> , 2019, 19, 1152.	2.9	22
52	Validation and calibration of the patient health questionnaire (PHQ-9) in Argentina. <i>BMC Psychiatry</i> , 2019, 19, 291.	2.6	70
53	The clinical effectiveness of sertraline in primary care and the role of depression severity and duration (PANDA): a pragmatic, double-blind, placebo-controlled randomised trial. <i>Lancet Psychiatry</i> , 2019, 6, 903-914.	7.4	78
54	Effect of a stepped-care intervention delivered by lay health workers on major depressive disorder among primary care patients in Nigeria (STPEPCARE): a cluster-randomised controlled trial. <i>The Lancet Global Health</i> , 2019, 7, e951-e960.	6.3	47

#	ARTICLE	IF	CITATIONS
55	Is teachers' mental health and wellbeing associated with students' mental health and wellbeing?. Journal of Affective Disorders, 2019, 253, 460-466.	4.1	28
56	Women's circles as a culturally safe psychosocial intervention in Guatemalan indigenous communities: a community-led pilot randomised trial. BMC Women's Health, 2019, 19, 53.	2.0	21
57	Rates, types and co-occurrence of emotional and behavioural disorders among perinatally HIV-infected youth in Uganda: the CHAKA study. Social Psychiatry and Psychiatric Epidemiology, 2019, 54, 415-425.	3.1	32
58	Psychotherapy adaptation in aging populations. , 2019, , 321-340.		2
59	High- versus low-intensity interventions for perinatal depression delivered by non-specialist primary maternal care providers in Nigeria: cluster randomised controlled trial (the EXPONATE trial). British Journal of Psychiatry, 2019, 215, 528-535.	2.8	38
60	Investigating Software Requirements for Systems Supporting Task-Shifted Interventions: Usability Study. Journal of Medical Internet Research, 2019, 21, e11346.	4.3	9
61	Use of a Mobile Phone App to Treat Depression Comorbid With Hypertension or Diabetes: A Pilot Study in Brazil and Peru. JMIR Mental Health, 2019, 6, e11698.	3.3	36
62	Addressing Depression Comorbid With Diabetes or Hypertension in Resource-Poor Settings: A Qualitative Study About User Perception of a Nurse-Supported Smartphone App in Peru. JMIR Mental Health, 2019, 6, e11701.	3.3	20
63	Antidepressant treatment with sertraline for adults with depressive symptoms in primary care: the PANDA research programme including RCT. Programme Grants for Applied Research, 2019, 7, 1-108.	1.0	6
64	Determinants of access to health care for depression in 49 countries: A multilevel analysis. Journal of Affective Disorders, 2018, 234, 80-88.	4.1	42
65	Resilience in high-risk adolescents of mothers with recurrent depressive disorder: The contribution of fathers. Journal of Adolescence, 2018, 65, 207-218.	2.4	11
66	Major Depressive Disorder: Longitudinal Analysis of Impact on Clinical and Behavioral Outcomes in Uganda. Journal of Acquired Immune Deficiency Syndromes (1999), 2018, 78, 136-143.	2.1	20
67	A cluster randomised controlled trial of the Wellbeing in Secondary Education (WISE) Project – an intervention to improve the mental health support and training available to secondary school teachers: protocol for an integrated process evaluation. Trials, 2018, 19, 270.	1.6	61
68	Trained lay health workers reduce common mental disorder symptoms of adults with suicidal ideation in Zimbabwe: a cohort study. BMC Public Health, 2018, 18, 227.	2.9	33
69	The Impact of Universal Health Care Programmes on Improving ‘Realized Access’ to Care for Depression in Chile. Administration and Policy in Mental Health and Mental Health Services Research, 2018, 45, 790-799.	2.1	14
70	Mental health and associated factors among young offenders in Chile: a cross-sectional study. Criminal Behaviour and Mental Health, 2018, 28, 13-27.	0.8	8
71	Collaborative meta-analysis finds no evidence of a strong interaction between stress and 5-HTTLPR genotype contributing to the development of depression. Molecular Psychiatry, 2018, 23, 133-142.	7.9	247
72	Impact of a Blended Web-Based Mindfulness Programme for General Practitioners: a Pilot Study. Mindfulness, 2018, 9, 129-139.	2.8	11

#	ARTICLE	IF	CITATIONS
73	Impact of the New Mental Health Services on Rates of Suicide and Hospitalisations by Attempted Suicide, Psychiatric Problems, and Alcohol Problems in Brazil. Administration and Policy in Mental Health and Mental Health Services Research, 2018, 45, 381-391.	2.1	9
74	Conditional cash transfer programme: Impact on homicide rates and hospitalisations from violence in Brazil. PLoS ONE, 2018, 13, e0208925.	2.5	35
75	Association between School Membership and Substance Use among Adolescents. Frontiers in Psychiatry, 2018, 9, 25.	2.6	7
76	Mental health among children and adolescents: Construct validity, reliability, and parent-adolescent agreement on the "Strengths and Difficulties Questionnaire"™ in Chile. PLoS ONE, 2018, 13, e0191809.	2.5	26
77	Provision of mental health care within primary care in Peru: A qualitative study exploring the perspectives of psychologists, primary health care providers, and patients. Wellcome Open Research, 2018, 3, 9.	1.8	8
78	Remote Collaborative Depression Care Program for Adolescents in Araucanãa Region, Chile: Randomized Controlled Trial. Journal of Medical Internet Research, 2018, 20, e38.	4.3	27
79	Integration of a Technology-Based Mental Health Screening Program Into Routine Practices of Primary Health Care Services in Peru (The Allillanchu Project): Development and Implementation. Journal of Medical Internet Research, 2018, 20, e100.	4.3	59
80	The Healthy Activity Program (HAP), a lay counsellor-delivered brief psychological treatment for severe depression, in primary care in India: a randomised controlled trial. Lancet, The, 2017, 389, 176-185.	13.7	292
81	Transitioning mental health into primary care. Lancet Psychiatry,the, 2017, 4, 90-92.	7.4	27
82	Attitudes towards digital treatment for depression: A European stakeholder survey. Internet Interventions, 2017, 8, 1-9.	2.7	201
83	How to reduce the number of rating scale items without predictability loss?. Scientometrics, 2017, 111, 581-593.	3.0	19
84	Telephone-supported computerised cognitive"behavioural therapy: REEACT-2 large-scale pragmatic randomised controlled trial. British Journal of Psychiatry, 2017, 210, 362-367.	2.8	94
85	Individual and contextual factors associated with tobacco, alcohol, and cannabis use among Chilean adolescents: A multilevel study. Journal of Adolescence, 2017, 56, 166-178.	2.4	32
86	Digital technology for treating and preventing mental disorders in low-income and middle-income countries: a narrative review of the literature. Lancet Psychiatry,the, 2017, 4, 486-500.	7.4	363
87	The KiVa antibullying program in primary schools in Chile, with and without the digital game component: study protocol for a randomized controlled trial. Trials, 2017, 18, 75.	1.6	17
88	Incidence and Persistence of Major Depressive Disorder Among People Living with HIV in Uganda. AIDS and Behavior, 2017, 21, 1641-1654.	2.7	20
89	Substance Use among Adolescents Involved in Bullying: A Cross-Sectional Multilevel Study. Frontiers in Psychology, 2017, 8, 1056.	2.1	42
90	Evaluating a multi-component, community-based program to improve adherence and retention in care among adolescents living with HIV in Zimbabwe: study protocol for a cluster randomized controlled trial. Trials, 2017, 18, 478.	1.6	31

#	ARTICLE	IF	CITATIONS
91	A randomised controlled trial assessing the severity and duration of depressive symptoms associated with a clinically significant response to sertraline versus placebo, in people presenting to primary care with depression (PANDA trial): study protocol for a randomised controlled trial. <i>Trials</i> , 2017, 18, 496.	1.6	15
92	Sustained effectiveness and cost-effectiveness of the Healthy Activity Programme, a brief psychological treatment for depression delivered by lay counsellors in primary care: 12-month follow-up of a randomised controlled trial. <i>PLoS Medicine</i> , 2017, 14, e1002385.	8.4	68
93	Towards sustainable partnerships in global health: the case of the CRONICAS Centre of Excellence in Chronic Diseases in Peru. <i>Globalization and Health</i> , 2016, 12, 29.	4.9	10
94	Validation of the Spanish Version of the Psychological Sense of School Membership (PSSM) Scale in Chilean Adolescents and Its Association with School-Related Outcomes and Substance Use. <i>Frontiers in Psychology</i> , 2016, 7, 1901.	2.1	16
95	Patterns of health-related behaviours among adolescents: a cross-sectional study based on the National Survey of School Health Brazil 2012. <i>BMJ Open</i> , 2016, 6, e011571.	1.9	29
96	Effect of a Primary Care-Based Psychological Intervention on Symptoms of Common Mental Disorders in Zimbabwe. <i>JAMA - Journal of the American Medical Association</i> , 2016, 316, 2618.	7.4	297
97	Validation of screening tools for depression and anxiety disorders in a primary care population with high HIV prevalence in Zimbabwe. <i>Journal of Affective Disorders</i> , 2016, 198, 50-55.	4.1	110
98	The Healthy Activity Program lay counsellor delivered treatment for severe depression in India: Systematic development and randomised evaluation. <i>British Journal of Psychiatry</i> , 2016, 208, 381-388.	2.8	115
99	Brief report: Association between psychological sense of school membership and mental health among early adolescents. <i>Journal of Adolescence</i> , 2016, 50, 1-5.	2.4	14
100	Hispanic immigrants in the USA: social and mental health perspectives. <i>Lancet Psychiatry</i> , 2016, 3, 860-870.	7.4	66
101	Influence of school-related factors on smoking among Chilean adolescents: a cross-sectional multilevel study. <i>BMC Pediatrics</i> , 2016, 16, 79.	1.7	8
102	Anxiety, depression and risk of cannabis use: Examining the internalising pathway to use among Chilean adolescents. <i>Drug and Alcohol Dependence</i> , 2016, 166, 109-115.	3.2	33
103	Epidemiology of maternal depression, risk factors, and child outcomes in low-income and middle-income countries. <i>Lancet Psychiatry</i> , 2016, 3, 973-982.	7.4	673
104	Indicated school-based intervention to improve depressive symptoms among at risk Chilean adolescents: a randomized controlled trial. <i>BMC Psychiatry</i> , 2016, 16, 276.	2.6	24
105	A pilot cluster randomised controlled trial of a support and training intervention to improve the mental health of secondary school teachers and students – the WISE (Wellbeing in Secondary) Tj ETQq1 1 0.784214 rgBT 10/Overlock 1	2.4	10
106	Protocol for a cluster randomised controlled trial of an intervention to improve the mental health support and training available to secondary school teachers – the WISE (Wellbeing in Secondary) Tj ETQq0 0 0 rgBT 10/Overlock 10 Tf 5	2.4	10
107	European COMPARative Effectiveness research on blended Depression treatment versus treatment-as-usual (E-COMPARED): study protocol for a randomized controlled, non-inferiority trial in eight European countries. <i>Trials</i> , 2016, 17, 387.	1.6	118
108	Longitudinal association between different levels of alcohol consumption and a new onset of depression and generalized anxiety disorder: Results from an international study in primary care. <i>Psychiatry Research</i> , 2016, 243, 30-34.	3.3	62

#	ARTICLE	IF	CITATIONS
109	Drinking to Cope: a Latent Class Analysis of Coping Motives for Alcohol Use in a Large Cohort of Adolescents. <i>Prevention Science</i> , 2016, 17, 584-594.	2.6	48
110	“Opening up the mind”: problem-solving therapy delivered by female lay health workers to improve access to evidence-based care for depression and other common mental disorders through the Friendship Bench Project in Zimbabwe. <i>International Journal of Mental Health Systems</i> , 2016, 10, 39.	2.7	58
111	Building Research Capacity Across and Within Low- and Middle-Income Countries: The Collaborative Hubs for International Research on Mental Health. <i>Academic Psychiatry</i> , 2016, 40, 686-691.	0.9	16
112	Teachers' wellbeing and depressive symptoms, and associated risk factors: A large cross sectional study in English secondary schools. <i>Journal of Affective Disorders</i> , 2016, 192, 76-82.	4.1	183
113	An Internet-Based Intervention for Depression in Primary Care in Spain: A Randomized Controlled Trial. <i>Journal of Medical Internet Research</i> , 2016, 18, e231.	4.3	59
114	Clinical effectiveness and cost-effectiveness of collaborative care for depression in UK primary care (CADET): a cluster randomised controlled trial. <i>Health Technology Assessment</i> , 2016, 20, 1-192.	2.8	41
115	The second Randomised Evaluation of the Effectiveness, cost-effectiveness and Acceptability of Computerised Therapy (REEACT-2) trial: does the provision of telephone support enhance the effectiveness of computer-delivered cognitive behaviour therapy? A randomised controlled trial. <i>Health Technology Assessment</i> , 2016, 20, 1-64.	2.8	31
116	Towards Clarity in Research on “Non-Cognitive” Skills: Linking Executive Functions, Self-Regulation, and Economic Development to Advance Life Outcomes for Children, Adolescents and Youth Globally. <i>Human Development</i> , 2015, 58, 313-317.	2.0	8
117	The Friendship Bench programme: a cluster randomised controlled trial of a brief psychological intervention for common mental disorders delivered by lay health workers in Zimbabwe. <i>International Journal of Mental Health Systems</i> , 2015, 9, 21.	2.7	99
118	“I was thinking too much”: experiences of HIV-positive adults with common mental disorders and poor adherence to antiretroviral therapy in Zimbabwe. <i>Tropical Medicine and International Health</i> , 2015, 20, 903-913.	2.3	58
119	Exposure to socioeconomic adversity in early life and risk of depression at 18 years: The mediating role of locus of control. <i>Journal of Affective Disorders</i> , 2015, 183, 269-278.	4.1	78
120	Computerised cognitive behaviour therapy (cCBT) as treatment for depression in primary care (REEACT) Tj ETQq0 0.0 rgBT /Overlock 10	6.0	313
121	Early Childhood Father Absence and Depressive Symptoms in Adolescent Girls from a UK Cohort: The Mediating Role of Early Menarche. <i>Journal of Abnormal Child Psychology</i> , 2015, 43, 921-931.	3.5	17
122	“Simulation-based learning in psychiatry for undergraduates at the University of Zimbabwe medical school”. <i>BMC Medical Education</i> , 2015, 15, 23.	2.4	22
123	A pilot randomized controlled trial of a stepped care intervention package for depression in primary care in Nigeria. <i>BMC Psychiatry</i> , 2015, 15, 96.	2.6	24
124	Individual and contextual factors associated with verbal bullying among Brazilian adolescents. <i>BMC Pediatrics</i> , 2015, 15, 49.	1.7	28
125	The Association Between School Bonding and Smoking Amongst Chilean Adolescents. <i>Substance Abuse</i> , 2015, 36, 515-523.	2.3	5
126	A cluster randomized clinical trial of a stepped care intervention for depression in primary care (STEP CARE)- study protocol. <i>BMC Psychiatry</i> , 2015, 15, 148.	2.6	14

#	ARTICLE	IF	CITATIONS
127	Expanding care for perinatal women with depression (EXPONATE): study protocol for a randomized controlled trial of an intervention package for perinatal depression in primary care. BMC Psychiatry, 2015, 15, 136.	2.6	28
128	Peer victimization during adolescence: concurrent and prospective impact on symptoms of depression and anxiety. Anxiety, Stress and Coping, 2015, 28, 105-120.	2.9	121
129	A randomised controlled trial of computerised cognitive behaviour therapy for the treatment of depression in primary care: the Randomised Evaluation of the Effectiveness and Acceptability of Computerised Therapy (REEACT) trial. Health Technology Assessment, 2015, 19, 1-174.	2.8	55
130	The Latin American treatment and innovation network in mental health h (LATINMH): rationale and scope. Revista De La Facultad De Ciencias Medicas De Cordoba, 2015, 72, 321-30.	0.3	6
131	Outreach to internally displaced persons in Bogotá, Colombia: challenges and potential solutions. Disaster Health, 2014, 2, 75-81.	0.6	7
132	The need for a behavioural science focus in research on mental health and mental disorders. International Journal of Methods in Psychiatric Research, 2014, 23, 28-40.	2.1	38
133	PEER VICTIMIZATION DURING ADOLESCENCE AND RISK FOR ANXIETY DISORDERS IN ADULTHOOD: A PROSPECTIVE COHORT STUDY. Depression and Anxiety, 2014, 31, 574-582.	4.1	128
134	The methods and outcomes of cultural adaptations of psychological treatments for depressive disorders: a systematic review. Psychological Medicine, 2014, 44, 1131-1146.	4.5	214
135	Do depressed and anxious men do groups? What works and what are the barriers to help seeking?. Primary Health Care Research and Development, 2014, 15, 287-301.	1.2	18
136	Factors associated with the development of self-harm amongst a socio-economically deprived cohort of adolescents in Santiago, Chile. Social Psychiatry and Psychiatric Epidemiology, 2014, 49, 629-637.	3.1	20
137	Advancing psychotherapy and evidence-based psychological interventions. International Journal of Methods in Psychiatric Research, 2014, 23, 58-91.	2.1	126
138	Prevention of anxiety and depression in Chinese: A randomized clinical trial testing the effectiveness of a stepped care program in primary care. Journal of Affective Disorders, 2014, 169, 212-220.	4.1	22
139	Mental health technologies and the needs of cultural groups. Lancet Psychiatry,the, 2014, 1, 326-327.	7.4	11
140	Internally Displaced "Victims of Armed Conflict" in Colombia: The Trajectory and Trauma Signature of Forced Migration. Current Psychiatry Reports, 2014, 16, 475.	4.5	105
141	ASSOCIATION BETWEEN EARLY TEMPERAMENT AND DEPRESSION AT 18 YEARS. Depression and Anxiety, 2014, 31, 729-736.	4.1	35
142	Computer-assisted cognitive-behavioral therapy for adolescent depression in primary care clinics in Santiago, Chile (YPSA-M): study protocol for a randomized controlled trial. Trials, 2014, 15, 309.	1.6	6
143	World Region of Origin and Common Mental Disorders Among Migrant Women in Spain. Journal of Immigrant and Minority Health, 2014, 16, 1111-1120.	1.6	11
144	The effectiveness and cost-effectiveness of lay counsellor-delivered psychological treatments for harmful and dependent drinking and moderate to severe depression in primary care in India: PREMIUM study protocol for randomized controlled trials. Trials, 2014, 15, 101.	1.6	57

#	ARTICLE	IF	CITATIONS
145	Using consumer perspectives to inform the cultural adaptation of psychological treatments for depression: A mixed methods study from South Asia. <i>Journal of Affective Disorders</i> , 2014, 163, 88-101.	4.1	40
146	Father absence and timing of menarche in adolescent girls from a UK cohort: The mediating role of maternal depression and major financial problems. <i>Journal of Adolescence</i> , 2014, 37, 291-301.	2.4	45
147	Cortisol awakening response and subsequent depression: prospective longitudinal study. <i>British Journal of Psychiatry</i> , 2014, 204, 137-143.	2.8	49
148	Cost-Effectiveness of Collaborative Care for Depression in UK Primary Care: Economic Evaluation of a Randomised Controlled Trial (CADET). <i>PLoS ONE</i> , 2014, 9, e104225.	2.5	38
149	Atendiendo la salud mental de las personas con enfermedades cr�nicas no transmisibles en el Per�: retos y oportunidades para la integraci�n de cuidados en el primer nivel de atenci�n. <i>Revista Peruana De Medicina De Experimental Y Salud Publica</i> , 2014, 31, .	0.4	18
150	Detecting depression among adolescents in Santiago, Chile: sex differences. <i>BMC Psychiatry</i> , 2013, 13, 122.	2.6	27
151	Expectations, experiences and attitudes of patients and primary care health professionals regarding online psychotherapeutic interventions for depression: protocol for a qualitative study. <i>BMC Psychiatry</i> , 2013, 13, 64.	2.6	19
152	Low intensity vs. self-guided Internet-delivered psychotherapy for major depression: a multicenter, controlled, randomized study. <i>BMC Psychiatry</i> , 2013, 13, 21.	2.6	22
153	Cross-cultural patterns of the association between varying levels of alcohol consumption and the common mental disorders of depression and anxiety: Secondary analysis of the WHO Collaborative Study on Psychological Problems in General Health Care. <i>Drug and Alcohol Dependence</i> , 2013, 133, 825-831.	3.2	53
154	The Emotionality Activity Sociability Temperament Survey: Factor analysis and temporal stability in a longitudinal cohort. <i>Personality and Individual Differences</i> , 2013, 54, 628-633.	2.9	38
155	Early Menarche and Depressive Symptoms From Adolescence to Young Adulthood in�� UK Cohort. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i> , 2013, 52, 591-598.e2.	0.5	39
156	Clinical effectiveness of collaborative care for depression in UK primary care (CADET): cluster randomised controlled trial. <i>BMJ</i> , The, 2013, 347, f4913-f4913.	6.0	173
157	Early life socio�economic position and later alcohol use: birth cohort study. <i>Addiction</i> , 2013, 108, 516-525.	3.3	35
158	Sleep disturbances and depressive symptoms: an investigation of their longitudinal association in a representative sample of the UK general population. <i>Psychological Medicine</i> , 2013, 43, 329-339.	4.5	25
159	Father absence and depressive symptoms in adolescence: findings from a UK cohort. <i>Psychological Medicine</i> , 2013, 43, 2615-2626.	4.5	43
160	School Intervention to Improve Mental Health of Students in Santiago, Chile. <i>JAMA Pediatrics</i> , 2013, 167, 1004.	6.2	58
161	Country- and individual-level socioeconomic determinants of depression: multilevel cross-national comparison. <i>British Journal of Psychiatry</i> , 2013, 202, 195-203.	2.8	175
162	Enhancing Psychosocial Support for HIV Positive Adolescents in Harare, Zimbabwe. <i>PLoS ONE</i> , 2013, 8, e70254.	2.5	119

#	ARTICLE	IF	CITATIONS
163	Depression Symptom Trajectories and Associated Risk Factors among Adolescents in Chile. PLoS ONE, 2013, 8, e78323.	2.5	20
164	A cluster randomised controlled trial to determine the clinical effectiveness and cost-effectiveness of classroom-based cognitive-behavioural therapy (CBT) in reducing symptoms of depression in high-risk adolescents. Health Technology Assessment, 2013, 17, vii-xvii, 1-109.	2.8	51
165	Economic evaluation of a task-shifting intervention for common mental disorders in India. Bulletin of the World Health Organization, 2012, 90, 813-821.	3.3	128
166	Classroom based cognitive behavioural therapy in reducing symptoms of depression in high risk adolescents: pragmatic cluster randomised controlled trial. BMJ, The, 2012, 345, e6058-e6058.	6.0	92
167	The Effect of the School Environment on the Emotional Health of Adolescents: A Systematic Review. Pediatrics, 2012, 129, 925-949.	2.1	186
168	Psychological distress and risk of long-term disability: population-based longitudinal study. Journal of Epidemiology and Community Health, 2012, 66, 586-592.	3.7	78
169	Folic acid supplementation during pregnancy may protect against depression 21 months after pregnancy, an effect modified by MTHFR C677T genotype. European Journal of Clinical Nutrition, 2012, 66, 97-103.	2.9	50
170	Association between pubertal development and depressive symptoms in girls from a UK cohort. Psychological Medicine, 2012, 42, 2579-2589.	4.5	57
171	Group psychological therapies for depression in the community: systematic review and meta-analysis. British Journal of Psychiatry, 2012, 200, 184-190.	2.8	105
172	Morale in the English mental health workforce: questionnaire survey. British Journal of Psychiatry, 2012, 201, 239-246.	2.8	48
173	Understanding burnout according to individual differences: ongoing explanatory power evaluation of two models for measuring burnout types. BMC Public Health, 2012, 12, 922.	2.9	32
174	Parental Socioeconomic Status and Risk of Offspring Autism Spectrum Disorders in a Swedish Population-Based Study. Journal of the American Academy of Child and Adolescent Psychiatry, 2012, 51, 467-476.e6.	0.5	188
175	The Relaxation Exercise and Social Support Trial (RESST): a community-based randomized controlled trial to alleviate medically unexplained vaginal discharge symptoms. BMC Psychiatry, 2012, 12, 195.	2.6	3
176	Bullying behaviour in schools, socioeconomic position and psychiatric morbidity: a cross-sectional study in late adolescents in Greece. Child and Adolescent Psychiatry and Mental Health, 2012, 6, 8.	2.5	46
177	Lessons from scaling up a depression treatment program in primary care in Chile. Revista Panamericana De Salud Publica/Pan American Journal of Public Health, 2012, 32, 234-240.	1.1	31
178	Scale up of services for mental health in low-income and middle-income countries. Lancet, The, 2011, 378, 1592-1603.	13.7	564
179	Smoking Is Associated with, but Does Not Cause, Depressed Mood in Pregnancy – A Mendelian Randomization Study. PLoS ONE, 2011, 6, e21689.	2.5	48
180	A practical approach to assess depression risk and to guide risk reduction strategies in later life. International Psychogeriatrics, 2011, 23, 280-291.	1.0	51

#	ARTICLE	IF	CITATIONS
181	Postnatal depression in women after unsuccessful attempted abortion. British Journal of Psychiatry, 2011, 198, 237-238.	2.8	7
182	The relaxation exercise and social support trial-resst: study protocol for a randomized community based trial. BMC Psychiatry, 2011, 11, 142.	2.6	8
183	The association between bullying and early stages of suicidal ideation in late adolescents in Greece. BMC Psychiatry, 2011, 11, 22.	2.6	50
184	Group cognitive behavioural therapy for women with depression: pilot and feasibility study for a randomised controlled trial using mixed methods. BMC Psychiatry, 2011, 11, 82.	2.6	40
185	Problem-solving therapy for depression and common mental disorders in Zimbabwe: piloting a task-shifting primary mental health care intervention in a population with a high prevalence of people living with HIV. BMC Public Health, 2011, 11, 828.	2.9	219
186	Towards a brief definition of burnout syndrome by subtypes: Development of the "Burnout Clinical Subtypes Questionnaire" (BCSQ-12). Health and Quality of Life Outcomes, 2011, 9, 74.	2.4	40
187	School-based intervention to improve the mental health of low-income, secondary school students in Santiago, Chile (YPSA): study protocol for a randomized controlled trial. Trials, 2011, 12, 49.	1.6	15
188	Common somatic symptoms, causal attributions of somatic symptoms and psychiatric morbidity in a cross-sectional community study in Santiago, Chile. BMC Research Notes, 2011, 4, 155.	1.4	11
189	Lay health worker led intervention for depressive and anxiety disorders in India: impact on clinical and disability outcomes over 12 months. British Journal of Psychiatry, 2011, 199, 459-466.	2.8	214
190	Timing of menarche and depressive symptoms in adolescent girls from a UK cohort. British Journal of Psychiatry, 2011, 198, 17-23.	2.8	91
191	Adolescent Alcohol and Tobacco Use and Early Socioeconomic Position: The ALSPAC Birth Cohort. Pediatrics, 2011, 127, e948-e955.	2.1	139
192	Depression in primary care: Chile's Programme for Screening, Diagnosis and Comprehensive Treatment of Depression. International Psychiatry: Bulletin of the Board of International Affairs of the Royal College of Psychiatrists, 2011, 8, 6-7.	0.1	2
193	Depression in primary care: Chile's Programme for Screening, Diagnosis and Comprehensive Treatment of Depression. International Psychiatry: Bulletin of the Board of International Affairs of the Royal College of Psychiatrists, 2011, 8, 6-7.	0.1	0
194	The Effect of Meditation on Psychological Distress among Buddhist Monks and Nuns. International Journal of Psychiatry in Medicine, 2010, 40, 461-468.	1.8	6
195	Job strain and other work conditions: relationships with psychological distress among civil servants in Rio de Janeiro, Brazil. Social Psychiatry and Psychiatric Epidemiology, 2010, 45, 345-354.	3.1	26
196	Cognitive ability, parental socioeconomic position and internalising and externalising problems in adolescence: Findings from two European cohort studies. European Journal of Epidemiology, 2010, 25, 569-580.	5.7	45
197	Protocol for a randomised controlled trial of a school based cognitive behaviour therapy (CBT) intervention to prevent depression in high risk adolescents (PROMISE). Trials, 2010, 11, 114.	1.6	16
198	Common mental disorders, subthreshold symptoms and disability: longitudinal study. British Journal of Psychiatry, 2010, 197, 411-412.	2.8	55

#	ARTICLE	IF	CITATIONS
199	Cigarette smoking and depression: a question of causation. British Journal of Psychiatry, 2010, 196, 425-426.	2.8	89
200	FC03-01 - The Association Between School Bonding and Smoking Amongst Adolescents in Chilean Schools. European Psychiatry, 2010, 25, .	0.2	0
201	Cost-effectiveness of improved primary care treatment of depression in women in Chile. British Journal of Psychiatry, 2010, 197, 291-296.	2.8	17
202	Violence against women by their intimate partner during pregnancy and postnatal depression: a prospective cohort study. Lancet, The, 2010, 376, 903-910.	13.7	249
203	Effectiveness of an intervention led by lay health counsellors for depressive and anxiety disorders in primary care in Goa, India (MANAS): a cluster randomised controlled trial. Lancet, The, 2010, 376, 2086-2095.	13.7	566
204	Common mental disorders during pregnancy and adverse obstetric outcomes. Journal of Psychosomatic Obstetrics and Gynaecology, 2010, 31, 229-235.	2.1	23
205	Epidemiología de la asociación entre los trastornos somatomorfos y los trastornos de ansiedad y depresivos. , 2010, , 1-8.		0
206	Invited commentary on "Mental health research priorities in low- and middle-income countries. British Journal of Psychiatry, 2009, 195, 364-365.	2.8	4
207	Packages of Care for Depression in Low- and Middle-Income Countries. PLoS Medicine, 2009, 6, e1000159.	8.4	161
208	Collaborative Depression Trial (CADET): multi-centre randomised controlled trial of collaborative care for depression - study protocol. BMC Health Services Research, 2009, 9, 188.	2.2	30
209	Emotional symptoms in children: The effect of maternal depression, life events, and COMT genotype. American Journal of Medical Genetics Part B: Neuropsychiatric Genetics, 2009, 150B, 209-218.	1.7	21
210	Effects of stressful life events, maternal depression and 5-HTTLPR genotype on emotional symptoms in pre-adolescent children. American Journal of Medical Genetics Part B: Neuropsychiatric Genetics, 2009, 150B, 670-682.	1.7	53
211	Common mental disorders during pregnancy: prevalence and associated factors among low-income women in São Paulo, Brazil. Archives of Women's Mental Health, 2009, 12, 335-343.	2.6	85
212	Validity of the self reporting questionnaire-20 in epidemiological studies with older adults. Social Psychiatry and Psychiatric Epidemiology, 2009, 44, 247-254.	3.1	50
213	Is socioeconomic status in early life associated with drug use? A systematic review of the evidence. Drug and Alcohol Review, 2009, 28, 142-153.	2.1	110
214	Therapist-delivered internet psychotherapy for depression in primary care: a randomised controlled trial. Lancet, The, 2009, 374, 628-634.	13.7	340
215	Chile: an ongoing mental health revolution. Lancet, The, 2009, 374, 597-598.	13.7	51
216	Improving the outcomes of primary care attenders with common mental disorders in developing countries: a cluster randomized controlled trial of a collaborative stepped care intervention in Goa, India. Trials, 2008, 9, 4.	1.6	56

#	ARTICLE	IF	CITATIONS
217	Smoking patterns during pregnancy and postnatal period and depressive symptoms. <i>Nicotine and Tobacco Research</i> , 2008, 10, 1609-1620.	2.6	60
218	Detecting common mental disorders in primary care in India: a comparison of five screening questionnaires. <i>Psychological Medicine</i> , 2008, 38, 221-228.	4.5	279
219	Risk factors across the life course and dementia in a Brazilian population: results from the Sao Paulo Ageing & Health Study (SPAH). <i>International Journal of Epidemiology</i> , 2008, 37, 879-890.	1.9	79
220	Integrating evidence-based treatments for common mental disorders in routine primary care: feasibility and acceptability of the MANAS intervention in Goa, India. <i>World Psychiatry</i> , 2008, 7, 39-46.	10.4	84
221	Experience counts. <i>World Psychiatry</i> , 2008, 7, 93-94.	10.4	1
222	Alcohol consumption as a risk factor for non-recovery from common mental disorder: results from the longitudinal follow-up of the National Psychiatric Morbidity Survey. <i>Psychological Medicine</i> , 2008, 38, 451-455.	4.5	33
223	Common mental disorders and the built environment in Santiago, Chile. <i>British Journal of Psychiatry</i> , 2007, 190, 394-401.	2.8	67
224	Epidemiology of the Association Between Somatoform Disorders and Anxiety and Depressive Disorders: An Update. <i>Psychosomatic Medicine</i> , 2007, 69, 860-863.	2.0	160
225	Mental health and quality of residential environment. <i>British Journal of Psychiatry</i> , 2007, 191, 500-505.	2.8	44
226	Treatment and prevention of mental disorders in low-income and middle-income countries. <i>Lancet</i> , The, 2007, 370, 991-1005.	13.7	737
227	Treatment of postnatal depression in low-income mothers in primary-care clinics in Santiago, Chile: a randomised controlled trial. <i>Lancet</i> , The, 2007, 370, 1629-1637.	13.7	156
228	Socio-economic status in childhood and later alcohol use: a systematic review*. <i>Addiction</i> , 2007, 102, 1546-1563.	3.3	69
229	Smoking and common mental disorders: a population-based survey in Santiago, Chile. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2007, 42, 874-880.	3.1	20
230	Un ensayo clínico aleatorizado de farmacoterapia con monitorización telefónica para mejorar el tratamiento de la depresión en la atención primaria en Santiago, Chile. <i>Revista Medica De Chile</i> , 2007, 135, .	0.2	14
231	Socio-economic position and common mental disorders. <i>British Journal of Psychiatry</i> , 2006, 189, 109-117.	2.8	170
232	Cost-Effectiveness of a Primary Care Treatment Program for Depression in Low-Income Women in Santiago, Chile. <i>American Journal of Psychiatry</i> , 2006, 163, 1379-1387.	7.2	103
233	Worry content across the lifespan: an analysis of 16- to 74-year-old participants in the British National Survey of Psychiatric Morbidity 2000. <i>Psychological Medicine</i> , 2006, 36, 1625-1633.	4.5	47
234	Inequities in Mental Health Care After Health Care System Reform in Chile. <i>American Journal of Public Health</i> , 2006, 96, 109-113.	2.7	38

#	ARTICLE	IF	CITATIONS
235	The thermolabile variant of MTHFR is associated with depression in the British Women's Heart and Health Study and a meta-analysis. <i>Molecular Psychiatry</i> , 2006, 11, 352-360.	7.9	126
236	Perceptions of social capital and the built environment and mental health. <i>Social Science and Medicine</i> , 2006, 62, 3072-3083.	3.8	156
237	Alcohol consumption as a risk factor for anxiety and depression. <i>British Journal of Psychiatry</i> , 2005, 187, 544-551.	2.8	94
238	Mental health inequalities in Wales, UK: Multi-level investigation of the effect of area deprivation. <i>British Journal of Psychiatry</i> , 2005, 186, 417-422.	2.8	74
239	An observation tool to assist with the assessment of urban residential environments. <i>Journal of Environmental Psychology</i> , 2005, 25, 293-305.	5.1	85
240	Comparing sex inequalities in common affective disorders across countries: Great Britain and Chile. <i>Social Science and Medicine</i> , 2005, 60, 1693-1703.	3.8	27
241	Negative self-schemas and the onset of depression in women: longitudinal study. <i>British Journal of Psychiatry</i> , 2005, 186, 302-307.	2.8	111
242	Use of psychotropic medication in Santiago, Chile. <i>Journal of Mental Health</i> , 2005, 14, 407-414.	1.9	9
243	Self-help books for depression: how can practitioners and patients make the right choice?. <i>British Journal of General Practice</i> , 2005, 55, 387-92.	1.4	73
244	Editorial: Treating depression in the developing world. <i>Tropical Medicine and International Health</i> , 2004, 9, 539-541.	2.3	61
245	Learning from low income countries: mental health. <i>BMJ: British Medical Journal</i> , 2004, 329, 1138-1140.	2.3	63
246	International and regional variation in the prevalence of common mental disorders: do we need more surveys?. <i>British Journal of Psychiatry</i> , 2004, 184, 289-290.	2.8	35
247	Treating depression in primary care in low-income women in Santiago, Chile: a randomised controlled trial. <i>Lancet, The</i> , 2003, 361, 995-1000.	13.7	388
248	Education and income: which is more important for mental health?. <i>Journal of Epidemiology and Community Health</i> , 2003, 57, 501-505.	3.7	173
249	Two-year follow-up after a randomised controlled trial of self- and clinician-accompanied exposure for phobia/panic disorders. <i>British Journal of Psychiatry</i> , 2001, 178, 543-548.	2.8	33
250	"Patient knows best"—detection of common mental disorders in Santiago, Chile: cross sectional study Commentary: Clinical practice is as important as diagnosis. <i>BMJ: British Medical Journal</i> , 2001, 322, 79-81.	2.3	19
251	Classification, disability and the public health agenda. <i>British Medical Bulletin</i> , 2001, 57, 3-15.	6.9	14
252	Common mental disorders in Santiago, Chile. <i>British Journal of Psychiatry</i> , 2001, 178, 228-233.	2.8	213

#	ARTICLE	IF	CITATIONS
253	The growing public health importance of common mental disorders. Epidemiology and Psychiatric Sciences, 2000, 9, 241-248.	3.9	1
254	Women, poverty and common mental disorders in four restructuring societies. Social Science and Medicine, 1999, 49, 1461-1471.	3.8	332
255	The management of depression in primary health care. Current Opinion in Psychiatry, 1999, 12, 103-107.	6.3	6
256	The comparison of latent variable models of non-psychotic psychiatric morbidity in four culturally diverse populations. Psychological Medicine, 1998, 28, 145-152.	4.5	52
257	Is the General Health Questionnaire (12 item) a culturally biased measure of psychiatric disorder?. Social Psychiatry and Psychiatric Epidemiology, 1995, 30, 20-25.	3.1	30
258	A cautious convert Ricardo Araya. Addiction, 1995, 90, 758-759.	3.3	5
259	Validation of the Edinburgh Postnatal Depression Scale (EPDS) in Chilean Postpartum Women. Journal of Psychosomatic Obstetrics and Gynaecology, 1995, 16, 187-191.	2.1	132
260	Women's drinking and society. Addiction, 1994, 89, 954-955.	3.3	3
261	Psychiatric Morbidity in Primary Health Care in Santiago, Chile Preliminary Findings. British Journal of Psychiatry, 1994, 165, 530-533.	2.8	34
262	Measuring psychiatric disorder in the community: a standardized assessment for use by lay interviewers. Psychological Medicine, 1992, 22, 465-486.	4.5	1,588
263	Comparison of two self administered psychiatric questionnaires (GHQ-12 and SRQ-20) in primary care in Chile. Social Psychiatry and Psychiatric Epidemiology, 1992, 27, 168-173.	3.1	127
264	Role of exposure homework in phobia reduction: A controlled study. Behavior Therapy, 1992, 23, 599-621.	2.4	77
265	Overseas doctors. Psychiatric Bulletin, 1990, 14, 312-312.	0.3	0
266	“Achieving a Balance” and visiting doctors: some unanswered questions. Psychiatric Bulletin, 1989, 13, 636-637.	0.3	0
267	Globalization and Mental Health. , 0, , 57-78.		10
268	Provision of mental health care within primary care in Peru: A qualitative study exploring the perspectives of psychologists, primary health care providers, and patients. Wellcome Open Research, 0, 3, 9.	1.8	2