
## Mari Nerhus

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2618729/publications.pdf Version: 2024-02-01



MADI NEDHIIS

#	Article	IF	CITATIONS
1	Resting-state high-frequency heart rate variability is related to respiratory frequency in individuals with severe mental illness but not healthy controls. Scientific Reports, 2016, 6, 37212.	1.6	52
2	Inflammatory evidence for the psychosis continuum model. Psychoneuroendocrinology, 2016, 67, 189-197.	1.3	39
3	Altered systemic cortisol metabolism in bipolar disorder and schizophrenia spectrum disorders. Journal of Psychiatric Research, 2014, 52, 57-62.	1.5	36
4	Low vitamin D is associated with negative and depressive symptoms in psychotic disorders. Schizophrenia Research, 2016, 178, 44-49.	1.1	29
5	Migrant background and ethnic minority status as predictors for duration of untreated psychosis. Microbial Biotechnology, 2015, 9, 61-65.	0.9	25
6	Persistent increase in TNF and ILâ€1 markers in severe mental disorders suggests traitâ€related inflammation: a one year followâ€up study. Acta Psychiatrica Scandinavica, 2017, 136, 400-408.	2.2	23
7	Vitamin D Deficiency Associated With Cognitive Functioning in Psychotic Disorders. Journal of Clinical Psychiatry, 2017, 78, e750-e757.	1.1	20
8	Childhood trauma mediates the association between ethnic minority status and more severe hallucinations in psychotic disorder. Psychological Medicine, 2015, 45, 133-142.	2.7	19
9	Vitamin D status in psychotic disorder patients and healthy controls – The influence of ethnic background. Psychiatry Research, 2015, 230, 616-621.	1.7	19
10	Serum levels of second-generation antipsychotics are associated with cognitive function in psychotic disorders. World Journal of Biological Psychiatry, 2017, 18, 471-482.	1.3	18
11	<i>VRK2</i> gene expression in schizophrenia, bipolar disorder and healthy controls. British Journal of Psychiatry, 2016, 209, 114-120.	1.7	17
12	Contribution of oxytocin receptor polymorphisms to amygdala activation in schizophrenia spectrum disorders. BJPsych Open, 2016, 2, 353-358.	0.3	11
13	Vitamin D levels, brain volume, and genetic architecture in patients with psychosis. PLoS ONE, 2018, 13, e0200250.	1.1	11
14	Higher vitamin B12 levels in neurodevelopmental disorders than in healthy controls and schizophrenia. FASEB Journal, 2020, 34, 8114-8124.	0.2	11
15	Serum level of venlafaxine is associated with better memory in psychotic disorders. Schizophrenia Research, 2015, 169, 386-392.	1.1	10
16	Serum concentrations of mood stabilizers are associated with memory, but not other cognitive domains in psychosis spectrum disorders; explorative analyses in a naturalistic setting. International Journal of Bipolar Disorders, 2016, 4, 24.	0.8	7
17	Hippocampus volume reduction in psychosis spectrum could be ameliorated by vitamin D. Schizophrenia Research, 2018, 199, 433-435.	1.1	7
18	Modelling difficulties in abstract thinking in psychosis: the importance of socio-developmental background. Cognitive Neuropsychiatry, 2017, 22, 39-52.	0.7	6

Mari Nerhus

#	Article	IF	CITATIONS
19	Psychosis: clinical insight and beliefs in immigrants in their first episode. Microbial Biotechnology, 2018, 12, 185-192.	0.9	6
20	Substance Misuse Trajectories and Risk of Relapse in the Early Course of Bipolar Disorder. Frontiers in Psychiatry, 2021, 12, 656912.	1.3	4
21	The Influence of Mental Health Literacy, Migration, and Education on the Duration of Untreated Psychosis. Frontiers in Public Health, 2021, 9, 705397.	1.3	3
22	M85. Age-Related Effect of Vitamin D on Hippocampus Volume in Schizophrenia and Bipolar Disorder. Schizophrenia Bulletin, 2017, 43, S241-S242.	2.3	2
23	Sex differences in antipsychotic-related triglyceride levels are associated with metabolic hormone differences in patients with severe mental disorders. Schizophrenia Research, 2022, 243, 55-63.	1.1	2
24	Vitamin D, Folate and the Intracranial Volume in Schizophrenia and Bipolar Disorder and Healthy Controls. Scientific Reports, 2018, 8, 10817.	1.6	1