

Tamie Nakajima

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2621725/publications.pdf>

Version: 2024-02-01

22
papers

533
citations

758635

12
h-index

713013

21
g-index

22
all docs

22
docs citations

22
times ranked

933
citing authors

#	ARTICLE	IF	CITATIONS
1	Bisphenol A may cause testosterone reduction by adversely affecting both testis and pituitary systems similar to estradiol. <i>Toxicology Letters</i> , 2010, 194, 16-25.	0.4	202
2	8-Hydroxydeoxyguanosine levels in human leukocyte and urine according to exposure to organophosphorus pesticides and paraoxonase 1 genotype. <i>International Archives of Occupational and Environmental Health</i> , 2007, 80, 217-227.	1.1	46
3	Dysregulated Bile Acid Synthesis, Metabolism and Excretion in a High Fat-Cholesterol Diet-Induced Fibrotic Steatohepatitis in Rats. <i>Digestive Diseases and Sciences</i> , 2013, 58, 2212-2222.	1.1	30
4	A Possible Role of Chenodeoxycholic Acid and Glycine-Conjugated Bile Acids in Fibrotic Steatohepatitis in a Dietary Rat Model. <i>Digestive Diseases and Sciences</i> , 2014, 59, 1490-1501.	1.1	28
5	Modulation of ammonium perfluorooctanoate-induced hepatic damage by genetically different PPAR α in mice. <i>Archives of Toxicology</i> , 2012, 86, 63-74.	1.9	27
6	“Hypothesis of Seven Balances” Molecular Mechanisms behind Alcoholic Liver Diseases and Association with PPAR α . <i>Journal of Occupational Health</i> , 2009, 51, 391-403.	1.0	24
7	Exposure to DEHP decreased four fatty acid levels in plasma of prepartum mice. <i>Toxicology</i> , 2013, 309, 52-60.	2.0	24
8	The modulation of hepatic adenosine triphosphate and inflammation by eicosapentaenoic acid during severe fibrotic progression in the SHRSP5/Dmcr rat model. <i>Life Sciences</i> , 2012, 90, 934-943.	2.0	21
9	Evidence for diazinon-mediated inhibition of cis-permethrin metabolism and its effects on reproductive toxicity in adult male mice. <i>Reproductive Toxicology</i> , 2012, 34, 489-497.	1.3	20
10	Bile acid detoxifying enzymes limit susceptibility to liver fibrosis in female SHRSP5/Dmcr rats fed with a high-fat-cholesterol diet. <i>PLoS ONE</i> , 2018, 13, e0192863.	1.1	18
11	Differences in metabolite burden of di(2-ethylhexyl)phthalate in pregnant and postpartum dams and their offspring in relation to drug-metabolizing enzymes in mice. <i>Archives of Toxicology</i> , 2012, 86, 563-569.	1.9	14
12	Differential Effects of Aging, Drinking and Exercise on Serum Cholesterol Levels Dependent on the PPAR α -V227A Polymorphism. <i>Journal of Occupational Health</i> , 2007, 49, 353-362.	1.0	13
13	Trichloroethylene and trichloroethanol induce skin sensitization with focal hepatic necrosis in guinea pigs. <i>Journal of Occupational Health</i> , 2020, 62, e12142.	1.0	13
14	High-fat and high-cholesterol diet decreases phosphorylated inositol-requiring kinase-1 and inhibits autophagy process in rat liver. <i>Scientific Reports</i> , 2019, 9, 12514.	1.6	11
15	In utero exposure to di(2-ethylhexyl)phthalate suppresses blood glucose and leptin levels in the offspring of wild-type mice. <i>Toxicology</i> , 2019, 415, 49-55.	2.0	11
16	Efficacy of Dietary Lipid Control in Healing High-Fat and High-Cholesterol Diet-Induced Fibrotic Steatohepatitis in Rats. <i>PLoS ONE</i> , 2016, 11, e0145939.	1.1	10
17	Increased serum anti-CYP2E1 IgG autoantibody levels may be involved in the pathogenesis of occupational trichloroethylene hypersensitivity syndrome: a case-control study. <i>Archives of Toxicology</i> , 2022, 96, 2785-2797.	1.9	6
18	Association of maternal whole blood fatty acid status during the prenatal period with term birth dimensions: a cross-sectional study. <i>Journal of Perinatal Medicine</i> , 2015, 43, 565-75.	0.6	5

#	ARTICLE	IF	CITATIONS
19	Increased risk of occupational trichloroethylene hypersensitivity syndrome at exposure levels higher than 15Âmg/L of urinary trichloroacetic acid, regardless of whether the patients had the HLA-B*13:01 allele. <i>Environmental Research</i> , 2020, 191, 109972.	3.7	5
20	The antihypertensive agent hydralazine reduced extracellular matrix synthesis and liver fibrosis in nonalcoholic steatohepatitis exacerbated by hypertension. <i>PLoS ONE</i> , 2020, 15, e0243846.	1.1	4
21	Simple method to detect triclofos and its metabolites in plasma of children by combined use of liquid chromatography tandem-mass spectrometry and gas chromatography-mass spectrometry. <i>Scientific Reports</i> , 2019, 9, 9294.	1.6	1
22	Associations between maternal mono-(2-ethylhexyl) phthalate levels, nuclear receptor gene polymorphisms, and fatty acid levels in pregnant Japanese women in the Hokkaido study. <i>Reproductive Toxicology</i> , 2022, 107, 22-32.	1.3	0