

Kyoichi Kaira

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/262341/publications.pdf>

Version: 2024-02-01

260
papers

6,084
citations

87843

38
h-index

123376

61
g-index

266
all docs

266
docs citations

266
times ranked

6773
citing authors

#	ARTICLE	IF	CITATIONS
1	The Risk of Cytotoxic Chemotherapy-Related Exacerbation of Interstitial Lung Disease with Lung Cancer. <i>Journal of Thoracic Oncology</i> , 2011, 6, 1242-1246.	0.5	177
2	Metabolic activity by ^{18}F -FDG-PET/CT is predictive of early response after nivolumab in previously treated NSCLC. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2018, 45, 56-66.	3.3	166
3	CD4+ T-cell Immunity in the Peripheral Blood Correlates with Response to Anti-PD-1 Therapy. <i>Cancer Immunology Research</i> , 2020, 8, 334-344.	1.6	155
4	Biologic Correlation of ^{18}F -Fluoro-2-Deoxy-D-Glucose Uptake on Positron Emission Tomography in Thymic Epithelial Tumors. <i>Journal of Clinical Oncology</i> , 2010, 28, 3746-3753.	0.8	143
5	Efficacy of gefitinib for non-adenocarcinoma non-small-cell lung cancer patients harboring epidermal growth factor receptor mutations: A pooled analysis of published reports. <i>Cancer Science</i> , 2011, 102, 1032-1037.	1.7	128
6	L-type amino acid transporter 1 and CD98 expression in primary and metastatic sites of human neoplasms. <i>Cancer Science</i> , 2008, 99, 2380-2386.	1.7	126
7	Usefulness of FDG-PET for early prediction of the response to gefitinib in non-small cell lung cancer. <i>Lung Cancer</i> , 2008, 59, 203-210.	0.9	125
8	Transport of 3-Fluoro-L-Methyl-Tyrosine by Tumor-Upregulated L-Type Amino Acid Transporter 1: A Cause of the Tumor Uptake in PET. <i>Journal of Nuclear Medicine</i> , 2012, 53, 1253-1261.	2.8	120
9	Diagnostic Usefulness of Fluorine- ^{18}F -Methyltyrosine Positron Emission Tomography in Combination With ^{18}F -Fluorodeoxyglucose in Sarcoidosis Patients. <i>Chest</i> , 2007, 131, 1019-1027.	0.4	119
10	Pulmonary Pleomorphic Carcinoma: A Clinicopathological Study Including EGFR Mutation Analysis. <i>Journal of Thoracic Oncology</i> , 2010, 5, 460-465.	0.5	107
11	Fluorine- ^{18}F -Methyltyrosine Positron Emission Tomography for Diagnosis and Staging of Lung Cancer: A Clinicopathologic Study. <i>Clinical Cancer Research</i> , 2007, 13, 6369-6378.	3.2	99
12	Biological significance of ^{18}F -FDG uptake on PET in patients with non-small-cell lung cancer. <i>Lung Cancer</i> , 2014, 83, 197-204.	0.9	98
13	Inhibition of L-type amino acid transporter 1 has antitumor activity in non-small cell lung cancer. <i>Anticancer Research</i> , 2010, 30, 4819-28.	0.5	95
14	Clinical significance of L-type amino acid transporter 1 expression as a prognostic marker and potential of new targeting therapy in biliary tract cancer. <i>BMC Cancer</i> , 2013, 13, 482.	1.1	81
15	Efficacy of chemotherapy with carboplatin and paclitaxel for unresectable thymic carcinoma. <i>Lung Cancer</i> , 2010, 67, 194-197.	0.9	79
16	2-Deoxy-2-[fluorine- ^{18}F] fluoro-d-glucose uptake on positron emission tomography is associated with programmed death ligand-1 expression in patients with pulmonary adenocarcinoma. <i>European Journal of Cancer</i> , 2018, 101, 181-190.	1.3	77
17	Transient IGF-1R inhibition combined with osimertinib eradicates AXL-low expressing EGFR mutated lung cancer. <i>Nature Communications</i> , 2020, 11, 4607.	5.8	69
18	CD98 Expression Is Associated with Poor Prognosis in Resected Non-Small-Cell Lung Cancer with Lymph Node Metastases. <i>Annals of Surgical Oncology</i> , 2009, 16, 3473-3481.	0.7	65

#	ARTICLE	IF	CITATIONS
19	Prognostic significance of L-type amino acid transporter 1 (LAT1) and 4F2 heavy chain (CD98) expression in stage I pulmonary adenocarcinoma. <i>Lung Cancer</i> , 2009, 66, 120-126.	0.9	65
20	Pooled analysis of the reports of erlotinib after failure of gefitinib for non-small cell lung cancer. <i>Lung Cancer</i> , 2010, 68, 99-104.	0.9	64
21	Improved efficacy of ramucirumab plus docetaxel after nivolumab failure in previously treated non-small cell lung cancer patients. <i>Thoracic Cancer</i> , 2019, 10, 775-781.	0.8	64
22	Expression of L-type amino acid transporter 1 (LAT1) as a prognostic and therapeutic indicator in multiple myeloma. <i>Cancer Science</i> , 2014, 105, 1496-1502.	1.7	54
23	Expression of L-type amino acid transporter 1 (LAT1) in neuroendocrine tumors of the lung. <i>Pathology Research and Practice</i> , 2008, 204, 553-561.	1.0	53
24	A phase II study of amrubicin, a synthetic 9-aminoanthracycline, in patients with previously treated lung cancer. <i>Lung Cancer</i> , 2010, 69, 99-104.	0.9	53
25	Prognostic significance of L-type amino acid transporter 1 (LAT1) expression in cutaneous melanoma. <i>Melanoma Research</i> , 2015, 25, 399-405.	0.6	52
26	Expression of amino acid transporters (LAT1, ASCT2 and xCT) as clinical significance in hepatocellular carcinoma. <i>Hepatology Research</i> , 2015, 45, 1014-1022.	1.8	51
27	LAT1 expression is closely associated with hypoxic markers and mTOR in resected non-small cell lung cancer. <i>American Journal of Translational Research (discontinued)</i> , 2011, 3, 468-78.	0.0	51
28	Correlation of angiogenesis with ¹⁸ F-FMT and ¹⁸ F-FDG uptake in non-small cell lung cancer. <i>Cancer Science</i> , 2009, 100, 753-758.	1.7	50
29	L-type amino acid transporter 1 expression is a prognostic marker in patients with surgically resected stage I non-small cell lung cancer. <i>Histopathology</i> , 2009, 54, 804-813.	1.6	49
30	¹⁸ F-FMT Uptake Seen Within Primary Cancer on PET Helps Predict Outcome of Non-Small Cell Lung Cancer. <i>Journal of Nuclear Medicine</i> , 2009, 50, 1770-1776.	2.8	47
31	Correlation Between ¹⁸ F-FDG Uptake on PET and Molecular Biology in Metastatic Pulmonary Tumors. <i>Journal of Nuclear Medicine</i> , 2011, 52, 705-711.	2.8	46
32	Relationship between ¹⁸ F-FDG uptake on positron emission tomography and molecular biology in malignant pleural mesothelioma. <i>European Journal of Cancer</i> , 2012, 48, 1244-1254.	1.3	46
33	Clinical significance of post-progression survival in lung cancer. <i>Thoracic Cancer</i> , 2017, 8, 379-386.	0.8	46
34	High STMN1 level is associated with chemo-resistance and poor prognosis in gastric cancer patients. <i>British Journal of Cancer</i> , 2017, 116, 1177-1185.	2.9	46
35	Correlation of tumor-related immunity with ¹⁸ F-FDG-PET in pulmonary squamous-cell carcinoma. <i>Lung Cancer</i> , 2018, 119, 71-77.	0.9	46
36	Potential of FDG-PET as Prognostic Significance after anti-PD-1 Antibody against Patients with Previously Treated Non-Small Cell Lung Cancer. <i>Journal of Clinical Medicine</i> , 2020, 9, 725.	1.0	46

#	ARTICLE	IF	CITATIONS
37	Relationship between LAT1 expression and resistance to chemotherapy in pancreatic ductal adenocarcinoma. <i>Cancer Chemotherapy and Pharmacology</i> , 2018, 81, 141-153.	1.1	45
38	Radiotherapy is an independent prognostic marker of favorable prognosis in non-small cell lung cancer patients after treatment with the immune checkpoint inhibitor, nivolumab. <i>Thoracic Cancer</i> , 2019, 10, 992-1000.	0.8	44
39	Long-term survivors of more than 5 years in advanced non-small cell lung cancer. <i>Lung Cancer</i> , 2010, 67, 120-123.	0.9	43
40	Prognostic significance of L-type amino acid transporter 1 (LAT1) and 4F2 heavy chain (CD98) expression in early stage squamous cell carcinoma of the lung. <i>Cancer Science</i> , 2009, 100, 249-254.	1.7	42
41	Efficacy of system L-type amino acid transporter 1 inhibition as a therapeutic target in esophageal squamous cell carcinoma. <i>Cancer Science</i> , 2016, 107, 1499-1505.	1.7	40
42	System L-type amino acid transporter 1 (LAT1) is frequently expressed in thymic carcinomas but is absent in thymomas. <i>Journal of Surgical Oncology</i> , 2009, 99, 433-438.	0.8	39
43	A phase II study of afatinib treatment for elderly patients with previously untreated advanced non-small-cell lung cancer harboring EGFR mutations. <i>Lung Cancer</i> , 2018, 126, 41-47.	0.9	39
44	Biologic correlates of 18F-FDG uptake on PET in pulmonary pleomorphic carcinoma. <i>Lung Cancer</i> , 2011, 71, 144-150.	0.9	38
45	Diagnostic usefulness of 18F-FAMT PET and L-type amino acid transporter 1 (LAT1) expression in oral squamous cell carcinoma. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2013, 40, 1692-1700.	3.3	38
46	Clinicopathological significance of LAT1 and ASCT2 in patients with surgically resected esophageal squamous cell carcinoma. <i>Journal of Surgical Oncology</i> , 2016, 113, 381-389.	0.8	38
47	Diagnostic value of 18F-FDG-PET to predict the tumour immune status defined by tumoural PD-L1 and CD8+ tumour-infiltrating lymphocytes in oral squamous cell carcinoma. <i>British Journal of Cancer</i> , 2020, 122, 1686-1694.	2.9	38
48	Clinicopathological significance of system L-type amino acid transporter 1 expression in pancreatic ductal carcinoma. <i>Histopathology</i> , 2015, 66, 234-243.	1.6	37
49	Clinical and Pathological Significance of ER Stress Marker (BiP/GRP78 and PERK) Expression in Malignant Melanoma. <i>Pathology and Oncology Research</i> , 2017, 23, 111-116.	0.9	37
50	Chemoradiotherapy followed by durvalumab in patients with unresectable advanced non-small cell lung cancer: Management of adverse events. <i>Thoracic Cancer</i> , 2020, 11, 1280-1287.	0.8	37
51	Evaluation of thoracic tumors with ¹⁸ F-FAMT and ¹⁸ F-FDG PET-CT: A clinicopathological study. <i>International Journal of Cancer</i> , 2009, 124, 1152-1160.	2.3	36
52	Specific transport of ³ fluoro-L-type amino acid methyl-L-tyrosine by LAT1 explains its specificity to malignant tumors in imaging. <i>Cancer Science</i> , 2016, 107, 347-352.	1.7	35
53	High Stromal TGFBI in Lung Cancer and Intratumoral CD8-Positive T Cells were Associated with Poor Prognosis and Therapeutic Resistance to Immune Checkpoint Inhibitors. <i>Annals of Surgical Oncology</i> , 2020, 27, 933-942.	0.7	35
54	Expression of Amino Acid Transporters (LAT1 and ASCT2) in Patients with Stage III/IV Laryngeal Squamous Cell Carcinoma. <i>Pathology and Oncology Research</i> , 2015, 21, 1175-1181.	0.9	34

#	ARTICLE	IF	CITATIONS
55	Positive PD-L1 Expression Predicts Worse Outcome in Cutaneous Angiosarcoma. <i>Journal of Global Oncology</i> , 2017, 3, 360-369.	0.5	34
56	Clinical difference between discontinuation and retreatment with nivolumab after immune-related adverse events in patients with lung cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2019, 84, 873-880.	1.1	32
57	Efficacy and safety of immune checkpoint inhibitor monotherapy in pretreated elderly patients with non-small cell lung cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2020, 85, 761-771.	1.1	32
58	Prognostic significance of diabetes mellitus in locally advanced non-small cell lung cancer. <i>BMC Cancer</i> , 2015, 15, 989.	1.1	31
59	High STMN1 Expression is Associated with Cancer Progression and Chemo-Resistance in Lung Squamous Cell Carcinoma. <i>Annals of Surgical Oncology</i> , 2017, 24, 4017-4024.	0.7	31
60	Post-treatment Glasgow Prognostic Score Predicts Efficacy in Advanced Non-small-cell Lung Cancer Treated With Anti-PD1. <i>Anticancer Research</i> , 2019, 39, 1455-1461.	0.5	31
61	Skeletal muscle mass predicts the outcome of nivolumab treatment for non-small cell lung cancer. <i>Medicine (United States)</i> , 2020, 99, e19059.	0.4	30
62	Progression-free survival, post-progression survival, and tumor response as surrogate markers for overall survival in patients with extensive small cell lung cancer. <i>Annals of Thoracic Medicine</i> , 2015, 10, 61-6.	0.7	30
63	L-type amino acid transporter 1 (LAT1) expression in malignant pleural mesothelioma. <i>Anticancer Research</i> , 2011, 31, 4075-82.	0.5	30
64	The role of β -tubulin in non-small cell lung cancer patients treated by taxane-based chemotherapy. <i>International Journal of Clinical Oncology</i> , 2013, 18, 371-379.	1.0	27
65	Clinical and Biological Significance of PD-L1 Expression Within the Tumor Microenvironment of Oral Squamous Cell Carcinoma. <i>Anticancer Research</i> , 2019, 39, 3039-3046.	0.5	27
66	Prognostic significance of L-type amino acid transporter 1 (LAT1) and 4F2 heavy chain (CD98) expression in surgically resectable stage III non-small cell lung cancer. <i>Experimental and Therapeutic Medicine</i> , 2010, 1, 799-808.	0.8	26
67	^{18}F -FDG uptake on PET helps predict outcome and response after treatment in unresectable thymic epithelial tumors. <i>Annals of Nuclear Medicine</i> , 2011, 25, 247-253.	1.2	26
68	Thymidylate synthase expression is closely associated with outcome in patients with pulmonary adenocarcinoma. <i>Medical Oncology</i> , 2012, 29, 1663-1672.	1.2	26
69	Expression of amino acid transporter (LAT1 and 4F2hc) in pulmonary pleomorphic carcinoma. <i>Human Pathology</i> , 2019, 84, 142-149.	1.1	26
70	Biological correlation of ^{18}F -FDG uptake on PET in pulmonary neuroendocrine tumors. <i>Anticancer Research</i> , 2013, 33, 4219-28.	0.5	26
71	MUC1 expression in thymic epithelial tumors: MUC1 may be useful marker as differential diagnosis between type B3 thymoma and thymic carcinoma. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2011, 458, 615-620.	1.4	25
72	Depolarized MUC1 Expression Is Closely Associated With Hypoxic Markers and Poor Outcome in Resected Non-Small Cell Lung Cancer. <i>International Journal of Surgical Pathology</i> , 2012, 20, 223-232.	0.4	25

#	ARTICLE	IF	CITATIONS
73	Expression of ER stress markers (GRP78/BiP and PERK) in adenoid cystic carcinoma. <i>Acta Oto-Laryngologica</i> , 2016, 136, 1-7.	0.3	25
74	Fluorodeoxyglucose uptake is associated with low tumor-infiltrating lymphocyte levels in patients with small cell lung cancer. <i>Lung Cancer</i> , 2019, 134, 180-186.	0.9	25
75	Different incidence of interstitial lung disease according to different kinds of EGFR tyrosine kinase inhibitors administered immediately before and/or after anti-PD-L1 antibodies in lung cancer. <i>Thoracic Cancer</i> , 2019, 10, 975-979.	0.8	25
76	FDG uptake reflects breast cancer immunological features: the PD-L1 expression and degree of TILs in primary breast cancer. <i>Breast Cancer Research and Treatment</i> , 2020, 181, 331-338.	1.1	25
77	Value of FDG Positron Emission Tomography in Monitoring the Effects of Therapy in Progressive Pulmonary Sarcoidosis. <i>Clinical Nuclear Medicine</i> , 2007, 32, 114-116.	0.7	24
78	Barium sulphate aspiration. <i>Lancet</i> , The, 2004, 364, 2220.	6.3	23
79	Comparison of L-type amino acid transporter 1 expression and L-[3-18F]-methyl tyrosine uptake in outcome of non-small cell lung cancer. <i>Nuclear Medicine and Biology</i> , 2010, 37, 911-916.	0.3	23
80	Tumor metabolic volume by 18F-FDG-PET as a prognostic predictor of first-line pembrolizumab for NSCLC patients with PD-L1 ≥ 50%. <i>Scientific Reports</i> , 2020, 10, 14990.	1.6	23
81	Clinical significance of coexpression of L-type amino acid transporter 1 (LAT1) and ASC amino acid transporter 2 (ASCT2) in lung adenocarcinoma. <i>American Journal of Translational Research (discontinued)</i> , 2015, 7, 1126-39.	0.0	23
82	Prognostic significance of L-type amino acid transporter 1 (LAT1) expression in patients with ovarian tumors. <i>American Journal of Translational Research (discontinued)</i> , 2015, 7, 1161-71.	0.0	23
83	CD98 as a novel prognostic indicator for patients with stage III/IV hypopharyngeal squamous cell carcinoma. <i>Head and Neck</i> , 2015, 37, 1569-1574.	0.9	22
84	Prognostic role of BiP/GRP78 expression as ER stress in patients with gastric adenocarcinoma. <i>Cancer Biomarkers</i> , 2017, 20, 273-281.	0.8	22
85	β2-Adrenergic receptor expression is associated with biomarkers of tumor immunity and predicts poor prognosis in estrogen receptor-negative breast cancer. <i>Breast Cancer Research and Treatment</i> , 2019, 177, 603-610.	1.1	22
86	First-line gefitinib treatment in elderly patients (aged ≥ 75 years) with non-small cell lung cancer harboring EGFR mutations. <i>Cancer Chemotherapy and Pharmacology</i> , 2015, 76, 761-769.	1.1	21
87	Role of Amino Acid Transporter Expression as a Prognostic Marker in Patients With Surgically Resected Colorectal Cancer. <i>Anticancer Research</i> , 2019, 39, 2535-2543.	0.5	21
88	Efficacy and safety of first-line pembrolizumab monotherapy in elderly patients (aged ≥ 75 years) with non-small cell lung cancer. <i>Journal of Cancer Research and Clinical Oncology</i> , 2020, 146, 457-466.	1.2	21
89	Incidence and dose-volume relationship of radiation pneumonitis after concurrent chemoradiotherapy followed by durvalumab for locally advanced non-small cell lung cancer. <i>Clinical and Translational Radiation Oncology</i> , 2020, 23, 85-88.	0.9	21
90	Pretreatment Glasgow prognostic score predicts survival among patients with high PD-L1 expression administered first-line pembrolizumab monotherapy for non-small cell lung cancer. <i>Cancer Medicine</i> , 2021, 10, 6971-6984.	1.3	21

#	ARTICLE	IF	CITATIONS
91	Value of 18F-FDG-PET to predict PD-L1 expression and outcomes of PD-1 inhibition therapy in human cancers. <i>Cancer Imaging</i> , 2021, 21, 11.	1.2	21
92	Management of Malignant Pericardial Effusion with Instillation of Mitomycin C in Non-small Cell Lung Cancer. <i>Japanese Journal of Clinical Oncology</i> , 2005, 35, 57-60.	0.6	20
93	Fulminant hepatic failure resulting from small-cell lung cancer and dramatic response of chemotherapy. <i>World Journal of Gastroenterology</i> , 2006, 12, 2466.	1.4	20
94	Expression of thymidylate synthase, orotate phosphoribosyltransferase and dihydropyrimidine dehydrogenase in thymic epithelial tumors. <i>Lung Cancer</i> , 2011, 74, 419-425.	0.9	20
95	Phase II study of oral S-1 and cisplatin with concurrent radiotherapy for locally advanced non-small-cell lung cancer. <i>Lung Cancer</i> , 2013, 82, 449-454.	0.9	20
96	Surrogate endpoints for overall survival in advanced non-small-cell lung cancer patients with mutations of the epidermal growth factor receptor gene. <i>Molecular and Clinical Oncology</i> , 2014, 2, 731-736.	0.4	20
97	L-type amino-acid transporter 1 expression predicts the response to preoperative hyperthermo-chemoradiotherapy for advanced rectal cancer. <i>Anticancer Research</i> , 2010, 30, 4223-7.	0.5	20
98	Expression of 4F2hc (CD98) in pulmonary neuroendocrine tumors. <i>Oncology Reports</i> , 2011, 26, 931-7.	1.2	19
99	18F-FAMT in patients with multiple myeloma: clinical utility compared to 18F-FDG. <i>Annals of Nuclear Medicine</i> , 2012, 26, 811-816.	1.2	19
100	The role of [¹⁸ F]fluorodeoxyglucose positron emission tomography in thymic epithelial tumors. <i>Cancer Imaging</i> , 2011, 11, 195-201.	1.2	19
101	18F-FDG uptake on PET in primary mediastinal non-thymic neoplasm: A clinicopathological study. <i>European Journal of Radiology</i> , 2012, 81, 2423-2429.	1.2	18
102	Biological evaluation of 3-[18F]fluoro- β -methyl-d-tyrosine (d-[18F]FAMT) as a novel amino acid tracer for positron emission tomography. <i>Annals of Nuclear Medicine</i> , 2013, 27, 314-324.	1.2	18
103	Forearm Muscle Metastasis as an Initial Clinical Manifestation of Lung Cancer. <i>Southern Medical Journal</i> , 2009, 102, 79-81.	0.3	17
104	High-grade neuroendocrine carcinoma of the lung shows increased thymidylate synthase expression compared to other histotypes. <i>Journal of Surgical Oncology</i> , 2010, 102, 11-17.	0.8	17
105	Decreased expression of class III β -tubulin is associated with unfavourable prognosis in patients with malignant melanoma. <i>Melanoma Research</i> , 2016, 26, 29-34.	0.6	17
106	Caspase14 expression is associated with triple negative phenotypes and cancer stem cell marker expression in breast cancer patients. <i>Journal of Surgical Oncology</i> , 2017, 116, 706-715.	0.8	17
107	Pre-existing interstitial lung disease does not affect prognosis in non-small cell lung cancer patients with <sc>PD-L1</sc> expression $\geq 50\%$ on first-line pembrolizumab. <i>Thoracic Cancer</i> , 2021, 12, 304-313.	0.8	17
108	A systemic review of PET and biology in lung cancer. <i>American Journal of Translational Research (discontinued)</i> , 2011, 3, 383-91.	0.0	17

#	ARTICLE	IF	CITATIONS
109	Assessment of Therapy Response in Lung Cancer With ^{18}F -Methyl Tyrosine PET. <i>American Journal of Roentgenology</i> , 2010, 195, 1204-1211.	1.0	16
110	Clinicopathological Significance of L-type Amino Acid Transporter 1 (LAT1) Expression in Patients with Adenoid Cystic Carcinoma. <i>Pathology and Oncology Research</i> , 2013, 19, 649-656.	0.9	16
111	^{18}F -FDG uptake on PET correlates with biological potential in early oral squamous cell carcinoma. <i>Acta Oto-Laryngologica</i> , 2015, 135, 494-499.	0.3	16
112	Clinical significance of β^2 -adrenergic receptor expression in patients with surgically resected gastric adenocarcinoma. <i>Tumor Biology</i> , 2016, 37, 13885-13892.	0.8	16
113	Prognostic significance of β^2 -adrenergic receptor expression in malignant melanoma. <i>Tumor Biology</i> , 2016, 37, 5971-5978.	0.8	16
114	Evaluation of gefitinib efficacy according to body mass index, body surface area, and body weight in patients with EGFR-mutated advanced non-small cell lung cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2017, 79, 497-505.	1.1	16
115	High expression of carcinoembryonic antigen and telomerase reverse transcriptase in circulating tumor cells is associated with poor clinical response to the immune checkpoint inhibitor nivolumab. <i>Oncology Letters</i> , 2018, 15, 3061-3067.	0.8	16
116	Fucosylated β^1 -acid glycoprotein as a biomarker to predict prognosis following tumor immunotherapy of patients with lung cancer. <i>Scientific Reports</i> , 2019, 9, 14503.	1.6	16
117	Prognostic Impact of β^2 Adrenergic Receptor Expression in Surgically Resected Pulmonary Pleomorphic Carcinoma. <i>Anticancer Research</i> , 2019, 39, 395-403.	0.5	16
118	Prognostic significance of β^2 -adrenergic receptor expression in non-small cell lung cancer. <i>American Journal of Translational Research (discontinued)</i> , 2016, 8, 5059-5070.	0.0	16
119	Prognostic significance of PD-L1 expression and tumor infiltrating lymphocytes in large cell neuroendocrine carcinoma of lung. <i>American Journal of Translational Research (discontinued)</i> , 2018, 10, 3243-3253.	0.0	16
120	Expression of Excision Repair Cross-Complementation Group 1, Breast Cancer Susceptibility 1, and β^2 III-Tubulin in Thymic Epithelial Tumors. <i>Journal of Thoracic Oncology</i> , 2011, 6, 606-613.	0.5	15
121	MUC1 Expression in Pulmonary Metastatic Tumors: A Comparison of Primary Lung Cancer. <i>Pathology and Oncology Research</i> , 2012, 18, 439-447.	0.9	15
122	Individual-level data on the relationships of progression-free survival and post-progression survival with overall survival in patients with advanced non-squamous non-small cell lung cancer patients who received second-line chemotherapy. <i>Medical Oncology</i> , 2014, 31, 88.	1.2	15
123	Decreasing expression of glucose-regulated protein GRP78/BiP as a significant prognostic predictor in patients with advanced laryngeal squamous cell carcinoma. <i>Head and Neck</i> , 2016, 38, 1539-1544.	0.9	15
124	Transport of 3-fluoro- ^{18}F -methyl-tyrosine (FAMT) by organic ion transporters explains renal background in [^{18}F]FAMT positron emission tomography. <i>Journal of Pharmacological Sciences</i> , 2016, 130, 101-109.	1.1	15
125	STXBP4 Drives Tumor Growth and Is Associated with Poor Prognosis through PDGF Receptor Signaling in Lung Squamous Cell Carcinoma. <i>Clinical Cancer Research</i> , 2017, 23, 3442-3452.	3.2	15
126	Sodium Glucose Cotransporter 2 Inhibition Combined With Cetuximab Significantly Reduced Tumor Size and Carcinoembryonic Antigen Level in Colon Cancer Metastatic to Liver. <i>Clinical Colorectal Cancer</i> , 2018, 17, e45-e48.	1.0	15

#	ARTICLE	IF	CITATIONS
127	Clinical Outcomes of Second-Line Chemotherapy in Patients with Previously Treated Advanced Thymic Carcinoma: A Retrospective Analysis of 191 Patients from the NEJ023 Study. <i>Oncologist</i> , 2020, 25, e668-e674.	1.9	15
128	Effect of durvalumab on local control after concurrent chemoradiotherapy for locally advanced non-small cell lung cancer in comparison with chemoradiotherapy alone. <i>Thoracic Cancer</i> , 2021, 12, 245-250.	0.8	15
129	Relationship between CD147 and expression of amino acid transporters (LAT1 and ASCT2) in patients with pancreatic cancer. <i>American Journal of Translational Research (discontinued)</i> , 2015, 7, 356-63.	0.0	15
130	Relationship between LAT1 expression and response to platinum-based chemotherapy in non-small cell lung cancer patients with postoperative recurrence. <i>Anticancer Research</i> , 2011, 31, 3775-82.	0.5	15
131	Laryngeal Sarcoidosis Detected by FDG Positron Emission Tomography. <i>Clinical Nuclear Medicine</i> , 2008, 33, 878-879.	0.7	14
132	Ratio of standardized uptake value on PET helps predict response and outcome after chemotherapy in advanced non-small cell lung cancer. <i>Annals of Nuclear Medicine</i> , 2010, 24, 697-705.	1.2	14
133	CD98 is a promising prognostic biomarker in biliary tract cancer. <i>Hepatobiliary and Pancreatic Diseases International</i> , 2014, 13, 654-657.	0.6	14
134	High stathmin 1 expression is associated with poor prognosis and chemoradiation resistance in esophageal squamous cell carcinoma. <i>International Journal of Oncology</i> , 2017, 50, 1184-1190.	1.4	14
135	Severe hepatotoxicity due to osimertinib after nivolumab therapy in patients with non-small cell lung cancer harboring EGFR mutation. <i>Thoracic Cancer</i> , 2020, 11, 1045-1051.	0.8	14
136	CD98 expression is associated with the grade of malignancy in thymic epithelial tumors. <i>Oncology Reports</i> , 2010, 24, 861-7.	1.2	14
137	Dose-Escalation Study of Three-Dimensional Conformal Thoracic Radiotherapy With Concurrent S-1 and Cisplatin for Inoperable Stage III Non-Small-Cell Lung Cancer. <i>Clinical Lung Cancer</i> , 2013, 14, 440-445.	1.1	13
138	Successful afatinib treatment of advanced non-small-cell lung cancer patients undergoing hemodialysis. <i>Cancer Chemotherapy and Pharmacology</i> , 2017, 79, 209-213.	1.1	13
139	A retrospective study of amrubicin monotherapy for the treatment of relapsed small cell lung cancer in elderly patients. <i>Cancer Chemotherapy and Pharmacology</i> , 2017, 80, 615-622.	1.1	13
140	Accumulation of periostin in acute exacerbation of familial idiopathic pulmonary fibrosis. <i>Journal of Thoracic Disease</i> , 2018, 10, E587-E591.	0.6	13
141	Re-challenge of afatinib after 1st generation EGFR-TKI failure in patients with previously treated non-small cell lung cancer harboring EGFR mutation. <i>Cancer Chemotherapy and Pharmacology</i> , 2019, 83, 817-825.	1.1	13
142	Uptake of positron emission tomography tracers reflects the tumor immune status in esophageal squamous cell carcinoma. <i>Cancer Science</i> , 2020, 111, 1969-1978.	1.7	13
143	Association of L-type amino acid transporter 1 (LAT1) with the immune system and prognosis in invasive breast cancer. <i>Scientific Reports</i> , 2022, 12, 2742.	1.6	13
144	Real-world data of atezolizumab plus carboplatin and etoposide in elderly patients with extensive-stage small cell lung cancer. <i>Cancer Medicine</i> , 2023, 12, 73-83.	1.3	13

#	ARTICLE	IF	CITATIONS
145	A phase I dose-escalation study of S-1 plus carboplatin in patients with advanced non-small-cell lung cancer. <i>Anti-Cancer Drugs</i> , 2007, 18, 471-476.	0.7	12
146	Pulmonary traumatic pneumatocele and hematoma. <i>Japanese Journal of Radiology</i> , 2009, 27, 100-102.	1.0	12
147	Small-cell lung cancer with voltage-gated calcium channel antibody-positive paraneoplastic limbic encephalitis: a case report. <i>Journal of Medical Case Reports</i> , 2014, 8, 119.	0.4	12
148	Clinical Significance of the Relationship between Progression-Free Survival or Postprogression Survival and Overall Survival in Patients with Extensive Disease-Small-Cell Lung Cancer Treated with Carboplatin plus Etoposide. <i>Canadian Respiratory Journal</i> , 2016, 2016, 1-8.	0.8	12
149	Phase I study of nab-paclitaxel plus carboplatin and concurrent thoracic radiotherapy in patients with locally advanced non-small cell lung cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2017, 79, 165-171.	1.1	12
150	High expression of GRP78/BiP as a novel predictor of favorable outcomes in patients with advanced thymic carcinoma. <i>International Journal of Clinical Oncology</i> , 2017, 22, 872-879.	1.0	11
151	Osimertinib induced cardiomyopathy. <i>Medicine (United States)</i> , 2020, 99, e22301.	0.4	11
152	Glasgow prognostic score predicts efficacy and prognosis in patients with advanced non-small cell lung cancer receiving EGFR-TKI treatment. <i>Thoracic Cancer</i> , 2020, 11, 2188-2195.	0.8	11
153	Prognostic significance of β 2-adrenergic receptor expression in patients with surgically resected colorectal cancer. <i>International Journal of Clinical Oncology</i> , 2020, 25, 1137-1144.	1.0	11
154	Effect of Systemic Steroid Use for Immune-Related Adverse Events in Patients with Non-Small Cell Lung Cancer Receiving PD-1 Blockade Drugs. <i>Journal of Clinical Medicine</i> , 2021, 10, 3744.	1.0	11
155	Phase II study of weekly docetaxel and cisplatin in patients with non-small cell lung cancer. <i>Anti-Cancer Drugs</i> , 2005, 16, 455-460.	0.7	10
156	Phase 2 study of S-1 plus carboplatin in patients with advanced non-small cell lung cancer. <i>Lung Cancer</i> , 2010, 68, 253-257.	0.9	10
157	Post-Progression Survival Associated with Overall Survival in Patients with Advanced Non-Small-Cell Lung Cancer Receiving Docetaxel Monotherapy as Second-Line Chemotherapy. <i>Chemotherapy</i> , 2017, 62, 205-213.	0.8	10
158	A new enzyme immunoassay for the determination of highly sialylated and fucosylated human β 1-acid glycoprotein as a biomarker of tumorigenesis. <i>Clinica Chimica Acta</i> , 2018, 478, 120-128.	0.5	10
159	Severe Thrombocytopenia Associated With Pembrolizumab in Patients With Non-small Cell Lung Cancer (NSCLC): A Case Report and Literature Review. <i>In Vivo</i> , 2020, 34, 877-880.	0.6	10
160	Effectiveness of EGFR-TKI rechallenge immediately after PD-1 blockade failure. <i>Thoracic Cancer</i> , 2021, 12, 864-873.	0.8	10
161	Efficacy and Feasibility of Programmed Death-1/Programmed Death Ligand-1 Blockade Therapy in Non-Small Cell Lung Cancer Patients With High Antinuclear Antibody Titers. <i>Frontiers in Oncology</i> , 2021, 11, 610952.	1.3	10
162	Distinctive roles of syntaxin binding protein 4 and its action target, TP63, in lung squamous cell carcinoma: a theranostic study for the precision medicine. <i>BMC Cancer</i> , 2020, 20, 935.	1.1	10

#	ARTICLE	IF	CITATIONS
163	Primary Mediastinal Synovial Sarcoma. <i>Journal of Computer Assisted Tomography</i> , 2008, 32, 238-241.	0.5	9
164	Phase I Study of Oral S-1 Plus Cisplatin With Concurrent Radiotherapy for Locally Advanced Non-Small-Cell Lung Cancer. <i>International Journal of Radiation Oncology Biology Physics</i> , 2009, 75, 109-114.	0.4	9
165	High expression of nucleobindin 2 is associated with poor prognosis in gastric cancer. <i>Tumor Biology</i> , 2017, 39, 101042831770381.	0.8	9
166	Clinical significance of primary prophylactic pegylated-granulocyte colony stimulating factor after the administration of ramucirumab plus docetaxel in patients with previously treated non-small cell lung cancer. <i>Thoracic Cancer</i> , 2019, 10, 1005-1008.	0.8	9
167	Usefulness of ^{18}F -Methyltyrosine PET for Therapeutic Monitoring of Patients with Advanced Lung Cancer. <i>Anticancer Research</i> , 2016, 36, 6481-6490.	0.5	9
168	Prognostic Significance of the Expression of CD98 (4F2hc) in Gastric Cancer. <i>Anticancer Research</i> , 2017, 37, 631-636.	0.5	9
169	Clinical Significance and Phenotype of MTA1 Expression in Esophageal Squamous Cell Carcinoma. <i>Anticancer Research</i> , 2017, 37, 4147-4155.	0.5	9
170	Prognostic significance of thymidylate synthase expression in the adjuvant chemotherapy after resection for pulmonary metastases from colorectal cancer. <i>Anticancer Research</i> , 2011, 31, 2763-71.	0.5	9
171	Efficacy and safety of amrubicin monotherapy after atezolizumab plus carboplatin and etoposide in patients with relapsed small-cell lung cancer. <i>Investigational New Drugs</i> , 2022, 40, 1066-1079.	1.2	9
172	Prognostic and predictive factors in resected non-small-cell lung cancer. <i>Expert Opinion on Medical Diagnostics</i> , 2010, 4, 373-381.	1.6	8
173	The efficacy of amrubicin on central nervous system metastases originating from small-cell lung cancer: a case series of eight patients. <i>Investigational New Drugs</i> , 2015, 33, 755-760.	1.2	8
174	Prognostic value of metabolic tumor volume of pretreatment ^{18}F -FAMT PET/CT in non-small cell lung Cancer. <i>BMC Medical Imaging</i> , 2018, 18, 46.	1.4	8
175	Stathmin-1 Is a Useful Diagnostic Marker for High-Grade Lung Neuroendocrine Tumors. <i>Annals of Thoracic Surgery</i> , 2019, 108, 235-243.	0.7	8
176	Post-progression survival is highly linked to overall survival in patients with non-small cell lung cancer harboring sensitive EGFR mutations treated with first-line epidermal growth factor receptor tyrosine kinase inhibitors. <i>Thoracic Cancer</i> , 2019, 10, 2200-2208.	0.8	8
177	Severe gastritis due to pembrolizumab treatment in a lung cancer patient. <i>Respirology Case Reports</i> , 2020, 8, e00636.	0.3	8
178	Administration of docetaxel plus ramucirumab with primary prophylactic pegylated-granulocyte colony-stimulating factor for pretreated non-small cell lung cancer: a phase II study. <i>Supportive Care in Cancer</i> , 2020, 28, 4825-4831.	1.0	8
179	Progression of Idiopathic Pulmonary Fibrosis Is Associated with Silica/Silicate Inhalation. <i>Environmental Science and Technology Letters</i> , 2021, 8, 903-910.	3.9	8
180	Clinical Effectiveness of Immune Checkpoint Inhibitors in Non-Small-Cell Lung Cancer with a Poor Performance Status. <i>Medicina (Lithuania)</i> , 2021, 57, 1273.	0.8	8

#	ARTICLE	IF	CITATIONS
181	A Phase I Dose Escalation Study of Weekly Docetaxel and Carboplatin in Elderly Patients With Nonsmall Cell Lung Cancer. <i>American Journal of Clinical Oncology: Cancer Clinical Trials</i> , 2007, 30, 51-56.	0.6	7
182	Phase I trial of oral S-1 plus gemcitabine in elderly patients with nonsmall cell lung cancer. <i>Anti-Cancer Drugs</i> , 2008, 19, 289-294.	0.7	7
183	High expression of topoisomerase-II predicts favorable clinical outcomes in patients with relapsed small cell lung cancers receiving amrubicin. <i>Lung Cancer</i> , 2018, 115, 42-48.	0.9	7
184	Intrapericardial carboplatin in the management of malignant pericardial effusion in breast cancer: a pilot study. <i>Cancer Chemotherapy and Pharmacology</i> , 2019, 84, 655-660.	1.1	7
185	Drastic Response of Re-challenge of EGFR-TKIs Immediately After Nivolumab Therapy in EGFR-TKI-Resistant Patients. <i>Journal of Thoracic Oncology</i> , 2019, 14, e135-e136.	0.5	7
186	Relationship Between Tumor Immune Markers and Fluorine-18- β -Methyltyrosine ([18 F]FAMT) Uptake in Patients with Lung Cancer. <i>Molecular Imaging and Biology</i> , 2020, 22, 1078-1086.	1.3	7
187	Occurrence of Ventricular Fibrillation in a Patient With Lung Cancer Receiving Osimertinib. <i>Journal of Thoracic Oncology</i> , 2020, 15, e54-e55.	0.5	7
188	Course of postoperative relapse in non-small cell lung cancer is strongly associated with post-progression survival. <i>Thoracic Cancer</i> , 2021, 12, 2740-2748.	0.8	7
189	B-Cell Non-Hodgkin Lymphoma Presenting As an Endobronchial Polypoid Mass. <i>Journal of Thoracic Oncology</i> , 2008, 3, 530-531.	0.5	6
190	Phase II study of oral S-1 plus cisplatin with bevacizumab for advanced non-squamous non-small cell lung cancer. <i>Lung Cancer</i> , 2013, 82, 103-108.	0.9	6
191	Prognostic effect of class III β -tubulin and Topoisomerase-II in patients with advanced thymic carcinoma who received combination chemotherapy, including taxanes or topoisomerase-II inhibitors. <i>Oncology Letters</i> , 2017, 14, 2369-2378.	0.8	6
192	Dual inhibition of MEK and p38 impairs tumor growth in KRAS-mutated non-small cell lung cancer. <i>Oncology Letters</i> , 2019, 17, 3569-3575.	0.8	6
193	Pattern of Local Failure and its Risk Factors of Locally Advanced Non-small Cell Lung Cancer Treated With Concurrent Chemo-radiotherapy. <i>Anticancer Research</i> , 2020, 40, 3513-3517.	0.5	6
194	Expression of LAT1 and 4F2hc in Gastroenteropancreatic Neuroendocrine Neoplasms. <i>In Vivo</i> , 2021, 35, 2425-2432.	0.6	6
195	Incidence and risk factors for pneumonitis among patients with lung cancer who received immune checkpoint inhibitors after palliative thoracic radiotherapy. <i>Journal of Radiation Research</i> , 2021, 62, 669-675.	0.8	6
196	Feasibility of intensity modulated radiotherapy with involved field radiotherapy for Japanese patients with locally advanced non-small cell lung cancer. <i>Journal of Radiation Research</i> , 2021, 62, 894-900.	0.8	6
197	Pretreatment body mass index predicts survival among patients administered nivolumab monotherapy for pretreated non-small cell lung cancer. <i>Thoracic Cancer</i> , 2022, 13, 1479-1489.	0.8	6
198	Coinfection of invasive pulmonary aspergillosis and pneumocystis jiroveci pneumonia in a non-HIV patient. <i>Acta Medica Okayama</i> , 2007, 61, 235-8.	0.1	6

#	ARTICLE	IF	CITATIONS
199	Rapid Growth of Malignant Pleural Mesothelioma. <i>Journal of Thoracic Oncology</i> , 2007, 2, 966-967.	0.5	5
200	Thymic squamous cell carcinoma producing granulocyte colony-stimulating factor associated with a high serum level of interleukin 6. <i>International Journal of Clinical Oncology</i> , 2009, 14, 534-536.	1.0	5
201	Prognostic impact of 18F-FDG uptake on PET in non-small cell lung cancer patients with postoperative recurrence following platinum-based chemotherapy. <i>Respiratory Investigation</i> , 2014, 52, 121-128.	0.9	5
202	18F-FDG uptake on PET is a predictive marker of thymidylate synthase expression in patients with thoracic neoplasms. <i>Oncology Reports</i> , 2014, 31, 209-215.	1.2	5
203	Efficacy and safety of cytotoxic drug chemotherapy after first-line EGFR TKI treatment in elderly patients with non-small-cell lung cancer harboring sensitive EGFR mutations. <i>Cancer Chemotherapy and Pharmacology</i> , 2018, 82, 119-127.	1.1	5
204	Pirfenidone Improves Familial Idiopathic Pulmonary Fibrosis without Affecting Serum Periostin Levels. <i>Medicina (Lithuania)</i> , 2019, 55, 161.	0.8	5
205	Treatment with Tumor Necrosis Factor- α Inhibitors, History of Allergy, and Hypercalcemia Are Risk Factors of Immune Reconstitution Inflammatory Syndrome in HIV-Negative Pulmonary Tuberculosis Patients. <i>Journal of Clinical Medicine</i> , 2020, 9, 96.	1.0	5
206	Perspective of Immune Checkpoint Inhibitors in Thymic Carcinoma. <i>Cancers</i> , 2021, 13, 1065.	1.7	5
207	Comprehensive expressional analysis of chemosensitivity-related markers in large cell neuroendocrine carcinoma of the lung. <i>Thoracic Cancer</i> , 2021, 12, 2666-2679.	0.8	5
208	Tumor immunity is related to ¹⁸ F-FDG uptake in thymic epithelial tumor. <i>Cancer Medicine</i> , 2021, 10, 6317-6326.	1.3	5
209	Interstitial lung disease secondary to alectinib after interstitial injury induced by crizotinib. <i>Journal of Cancer Research and Therapeutics</i> , 2020, 16, 919.	0.3	5
210	Phase I Study of Biweekly Paclitaxel and Carboplatin for Frail Patients With Advanced Non-Small Cell Lung Cancer. <i>American Journal of Clinical Oncology: Cancer Clinical Trials</i> , 2007, 30, 487-491.	0.6	4
211	18F-FDG uptake on PET could be a predictive marker of Excision Repair Cross-Complementation Group 1 (ERCC1) expression in patients with thoracic neoplasms?. <i>Neoplasma</i> , 2012, 59, 257-263.	0.7	4
212	Clinical Significance of Various Drug-Sensitivity Markers in Patients with Surgically Resected Pulmonary Pleomorphic Carcinoma. <i>Cancers</i> , 2019, 11, 1636.	1.7	4
213	Prognostic Significance of Tumor Immunity in Surgically Resected Pulmonary Pleomorphic Carcinoma. <i>Anticancer Research</i> , 2020, 40, 261-269.	0.5	4
214	Clinical significance of topoisomerase II expression in patients with advanced non-small cell lung cancer treated with amrubicin. <i>Thoracic Cancer</i> , 2020, 11, 426-435.	0.8	4
215	Primary thymic adenocarcinoma with an aggressive clinical course: An autopsy case showing signet ring cell-like features. <i>Thoracic Cancer</i> , 2020, 11, 3609-3613.	0.8	4
216	Efficacy and safety of S-1 monotherapy in previously treated elderly patients (aged ≥ 75 years) with non-small cell lung cancer: A retrospective analysis. <i>Thoracic Cancer</i> , 2020, 11, 2867-2876.	0.8	4

#	ARTICLE	IF	CITATIONS
217	Prognostic Significance of Glucose Metabolism as GLUT1 in Patients with Pulmonary Pleomorphic Carcinoma. <i>Journal of Clinical Medicine</i> , 2020, 9, 413.	1.0	4
218	Efficacy and Safety of Anti-Programed Death-1 Blockade in Previously Treated Large-Cell Neuroendocrine Carcinoma. <i>Chemotherapy</i> , 2021, 66, 65-71.	0.8	4
219	Plasma platin ³ : A tumor marker in patients with non ³ small ³ cell lung cancer treated with nivolumab. <i>Oncology Letters</i> , 2020, 21, 1-1.	0.8	4
220	Salvage Chemotherapy in Patients with Previously Treated Thymic Carcinoma. <i>Cancers</i> , 2021, 13, 5441.	1.7	4
221	Visual Assessment of 18F-FDG Uptake on PET to Predict Survival Benefit to PD-1 Blockade in Non ³ Small Cell Lung Cancer. <i>Clinical Nuclear Medicine</i> , 2022, 47, 108-116.	0.7	4
222	A retrospective study of the efficacy and safety of naldemedine for opioid ³ induced constipation in thoracic cancer patients. <i>Thoracic Cancer</i> , 2022, 13, 2301-2308.	0.8	4
223	Small cell carcinoma of the parotid gland. <i>Otolaryngology - Head and Neck Surgery</i> , 2007, 136, 330-331.	1.1	3
224	Phase I dose escalation study of amrubicin plus paclitaxel in previously treated advanced non-small cell lung cancer. <i>International Journal of Clinical Oncology</i> , 2016, 21, 240-247.	1.0	3
225	Prospective exploratory study of gemcitabine and S-1 against elderly patients with advanced non-small cell lung cancer. <i>Oncology Letters</i> , 2017, 14, 1123-1128.	0.8	3
226	The effect of post-progression survival on overall survival among patients with sensitive relapse of small cell lung cancer. <i>Medical Oncology</i> , 2018, 35, 45.	1.2	3
227	Topotecan monotherapy for the treatment of relapsed small cell lung cancer in elderly patients: A retrospective analysis. <i>Thoracic Cancer</i> , 2018, 9, 1699-1706.	0.8	3
228	An Exploratory Randomized Phase II Trial Comparing CDDP Plus S-1 With Bevacizumab and CDDP Plus Pemetrexed With Bevacizumab Against Patients With Advanced Non-squamous Non-small Cell Lung Cancer. <i>Anticancer Research</i> , 2019, 39, 2483-2491.	0.5	3
229	Efficacy of PD-1 blockade therapy and T cell immunity in lung cancer patients. <i>Immunological Medicine</i> , 2020, 43, 10-15.	1.4	3
230	Stereotactic body radiation therapy using CyberKnife for T1N0M0 lung cancer patients with severe pulmonary dysfunction. <i>Journal of Radiation Research</i> , 2020, 61, 903-907.	0.8	3
231	A phase II study of daily carboplatin plus irradiation followed by durvalumab for stage III non-small cell lung cancer patients with PS 2 up to 74 ³ %years old and patients with PS 0 or 1 from 75 ³ %years: NEJ039A.1.1 (trial in progress). <i>BMC Cancer</i> , 2020, 20, 961.		3
232	Phase II Study of Weekly Nanoparticle Albumin-Bound Paclitaxel as Second- or Third-Line Therapy in Patients with Advanced Non-Small Cell Lung Cancer. <i>Chemotherapy</i> , 2020, 65, 21-28.	0.8	3
233	Recovery of the Sensitivity to Anti-PD-1 Antibody by Celecoxib in Lung Cancer. <i>Anticancer Research</i> , 2020, 40, 5309-5311.	0.5	3
234	Post-Progression Survival Influences Overall Survival among Patients with Advanced Non-Small Cell Lung Cancer Undergoing First-Line Pembrolizumab Monotherapy. <i>Oncology</i> , 2021, 99, 562-570.	0.9	3

#	ARTICLE	IF	CITATIONS
235	Combination of immune check inhibitor and immunomodulatory arabinomannan extracted from <i>Mycobacterium tuberculosis</i> : A case report. <i>Molecular and Clinical Oncology</i> , 2021, 15, 227.	0.4	3
236	Effectiveness and Safety of EGFR-TKI Rechallenge Treatment in Elderly Patients with Advanced Non-Small-Cell Lung Cancer Harboring Drug-Sensitive EGFR Mutations. <i>Medicina (Lithuania)</i> , 2021, 57, 929.	0.8	3
237	Carbonic anhydrase 9 expression is associated with poor prognosis, tumor proliferation, and radiosensitivity of thymic carcinomas. <i>Oncotarget</i> , 2019, 10, 1306-1319.	0.8	3
238	Management of Lung Cancer-Associated Malignant Pericardial Effusion with Intrapericardial Administration of Carboplatin: A Retrospective Study. <i>Current Oncology</i> , 2022, 29, 163-172.	0.9	3
239	Prospective assessment using 18F-FDG PET/CT as a novel predictor for early response to PD-1 blockade in non-small-cell lung cancer. <i>Scientific Reports</i> , 2022, 12, .	1.6	3
240	Dramatic Response of S-1 Administration to Chemorefractory Advanced Thymic Cancer. <i>Chemotherapy</i> , 2014, 60, 356-359.	0.8	2
241	Pseudoprogression mimicking hyperprogressive disease after pembrolizumab treatment in a patient with lung cancer. <i>Lung Cancer</i> , 2020, 139, 221-223.	0.9	2
242	Occurrence of Hematological Malignancy in Long-term Survivors With Advanced Thymic Cancer. <i>In Vivo</i> , 2020, 34, 1511-1513.	0.6	2
243	A phase II study of S-1 and cisplatin with concurrent thoracic radiotherapy followed by durvalumab for unresectable, locally advanced non-small-cell lung cancer in Japan (SAMURAI study). <i>Therapeutic Advances in Medical Oncology</i> , 2021, 13, 175883592199858.	1.4	2
244	Synchronous dilemma of sarcoid-like reaction and drastic response after PD-1 blockade administration in lung cancer. <i>Japanese Journal of Clinical Oncology</i> , 2021, 51, 1179-1180.	0.6	2
245	Post-Progression Survival Is Strongly Associated with Overall Survival in Patients Exhibiting Postoperative Relapse of Non-Small-Cell Lung Cancer Harboring Sensitizing EGFR Mutations. <i>Medicina (Lithuania)</i> , 2021, 57, 508.	0.8	2
246	Recurrent intimal sarcoma mimicking pulmonary embolism. <i>Japanese Journal of Clinical Oncology</i> , 2015, 45, 695-696.	0.6	1
247	Prognostic value of morphological characteristics assessed by CT scan in patients with non-small cell lung cancer treated with nivolumab. <i>Thoracic Cancer</i> , 2020, 11, 3521-3527.	0.8	1
248	Detection of pseudoprogression with [18F]-FDG-PET in a patient with pulmonary large cell neuroendocrine carcinoma who received anti-PD-1 treatment. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2021, 48, 1268-1270.	3.3	1
249	Clinical impact of post-progression survival on overall survival in patients receiving nivolumab monotherapy as a second-line treatment for advanced non-small cell lung cancer. <i>Thoracic Cancer</i> , 2021, 12, 1171-1179.	0.8	1
250	Post-Progression Survival Highly Influences Overall Survival in Driver Gene Mutation/Translocation Negative or Unknown Type of Non-Small Cell Lung Cancer. <i>Oncology</i> , 2022, 100, 89-100.	0.9	1
251	The Relationship between Tumor Development and Sarcoidosis in Aspects of Carcinogenesis before and after the Onset of Sarcoidosis. <i>Medicina (Lithuania)</i> , 2022, 58, 768.	0.8	1
252	Pulmonary Pleomorphic Carcinoma: Diagnosis Using Small Biopsy Specimens. <i>Journal of Thoracic Oncology</i> , 2010, 5, 1492-1493.	0.5	0

#	ARTICLE	IF	CITATIONS
253	Reply to: Relationship between LAT1 expression and resistance to chemotherapy in pancreatic ductal adenocarcinoma. <i>Cancer Chemotherapy and Pharmacology</i> , 2018, 82, 369-369.	1.1	0
254	Spontaneous aspiration of a long tree twig as foreign body. <i>Respirology Case Reports</i> , 2019, 7, e00401.	0.3	0
255	Verrucous skin lesion caused by neuropathy misdiagnosed as squamous cell carcinoma and treated with amputation surgery. <i>Journal of Dermatology</i> , 2020, 47, e188-e190.	0.6	0
256	Prospective Feasibility Study of Amrubicin and Bevacizumab Therapy for Patients With Previously Treated Advanced NSCLC. <i>Anticancer Research</i> , 2020, 40, 1571-1578.	0.5	0
257	A 54ÂGy in three fractions of stereotactic body radiotherapy using CyberKnife for T1b-2aN0M0 pathologically confirmed non-small cell lung cancer. <i>Japanese Journal of Clinical Oncology</i> , 2021, 51, 1723-1728.	0.6	0
258	PET-CT, Bio-imaging for Predicting Prognosis and Response to Chemotherapy in Patients with Lung Cancer. , 2017, , 45-61.		0
259	The Role of 18F-FDG-PET as Therapeutic Monitoring in Patients with Lung Cancer. , 2021, , 23-33.		0
260	Clinical impact of post-progression survival in patients with locally advanced non-small cell lung cancer after chemoradiotherapy. <i>Radiology and Oncology</i> , 2022, 56, 228-237.	0.6	0