Silvia Tabacchioni

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2623988/publications.pdf

Version: 2024-02-01

40 papers

1,953 citations

257357 24 h-index 38 g-index

41 all docs

41 docs citations

41 times ranked

2284 citing authors

#	Article	IF	CITATIONS
1	Identification of Beneficial Microbial Consortia and Bioactive Compounds with Potential as Plant Biostimulants for a Sustainable Agriculture. Microorganisms, 2021, 9, 426.	1.6	37
2	Does the Introduction of N2-Fixing Trees in Forest Plantations on Tropical Soils Ameliorate Low Fertility and Enhance Carbon Sequestration via Interactions Between Biota and Nutrient Availability? Case Studies From Central Africa and South America. Frontiers in Soil Science, 2021, 1, .	0.8	6
3	Influence of Acacia mangium on Soil Fertility and Bacterial Community in Eucalyptus Plantations in the Congolese Coastal Plains. Sustainability, 2020, 12, 8763.	1.6	13
4	Towards the development of a biobased economy in Europe and India. Critical Reviews in Biotechnology, 2019, 39, 779-799.	5.1	46
5	Anaerobic digestion of the above ground biomass of Jerusalem Artichoke in a pilot plant: Impact of the preservation method on the biogas yield and microbial community. Biomass and Bioenergy, 2018, 108, 190-197.	2.9	17
6	New Insights in Plant-Associated Paenibacillus Species: Biocontrol and Plant Growth-Promoting Activity., 2016,, 237-279.		5
7	Bio-Methane Production from Wastes: Focus on Feedstock Sources and Microbial Communities. , 2015, , 333-353.		0
8	Dynamics of hydrogen-producing bacteria in a repeated batch fermentation process using lake sediment as inoculum. Archives of Microbiology, 2014, 196, 97-107.	1.0	8
9	Vertical distribution of bacterioplankton in Lake Averno in relation to water chemistry. FEMS Microbiology Ecology, 2013, 84, 176-188.	1.3	14
10	The Paenibacillus polymyxa species is abundant among hydrogen-producing facultative anaerobic bacteria in Lake Averno sediment. Archives of Microbiology, 2012, 194, 345-351.	1.0	10
11	Pathogenicity and biotechnological applications of the genus Burkholderia. Open Life Sciences, 2011, 6, 997-1005.	0.6	5
12	Genetic relationships among Italian and Mexican maize-rhizosphere Burkholderia cepacia complex (BCC) populations belonging to Burkholderia cenocepacia IIIB and BCC6 group. BMC Microbiology, 2011, 11, 228.	1.3	5
13	Application of multiplex single nucleotide primer extension (mSNuPE) to the identification of bacteria: The Burkholderia cepacia complex case. Journal of Microbiological Methods, 2010, 80, 251-256.	0.7	12
14	Ecology and biotechnological potential of Paenibacillus polymyxa: a minireview. Indian Journal of Microbiology, 2009, 49, 2-10.	1.5	211
15	<i>RecA</i> gene sequence and Multilocus Sequence Typing for species-level resolution of <i>Burkholderia cepacia</i> complex isolates. Letters in Applied Microbiology, 2009, 49, 580-588.	1.0	32
16	Sustainable power production in a membrane-less and mediator-less synthetic wastewater microbial fuel cell. Bioresource Technology, 2009, 100, 3252-3260.	4.8	106
17	Use of the gyrB gene to discriminate among species of the Burkholderia cepacia complex. FEMS Microbiology Letters, 2008, 281, 175-182.	0.7	20
18	InvestigatingBurkholderia cepaciacomplex populations recovered from Italian maize rhizosphere by multilocus sequence typing. Environmental Microbiology, 2007, 9, 1632-1639.	1.8	35

#	Article	IF	CITATIONS
19	Burkholderia cepacia complex species: health hazards and biotechnological potential. Trends in Microbiology, 2006, 14, 277-286.	3.5	176
20	Perturbation of maize rhizosphere microflora following seed bacterization with Burkholderia cepacia MCI 7. FEMS Microbiology Ecology, 2006, 23, 183-193.	1.3	60
21	Detection of cultured and uncultured Burkholderia cepacia complex bacteria naturally occurring in the maize rhizosphere. Environmental Microbiology, 2005, 7, 1734-1742.	1.8	28
22	Metabolic Profiling of Burkholderia cenocepacia, Burkholderia ambifaria, and Burkholderia pyrrocinia Isolates from Maize Rhizosphere. Microbial Ecology, 2005, 50, 385-395.	1.4	24
23	Efficacy of species-specificrecAPCR tests in the identification ofBurkholderia cepaciacomplex environmental isolates. FEMS Microbiology Letters, 2005, 246, 39-45.	0.7	24
24	Effect of Fusarium verticillioides on maize-root-associated Burkholderia cenocepacia populations. Research in Microbiology, 2005, 156, 974-983.	1.0	28
25	Epidemiology and Clinical Course of Burkholderia cepacia Complex Infections, Particularly Those Caused by Different Burkholderia cenocepacia Strains, among Patients Attending an Italian Cystic Fibrosis Center. Journal of Clinical Microbiology, 2004, 42, 1491-1497.	1.8	59
26	Exopolysaccharides produced by Burkholderia cenocepacia recA lineages IIIA and IIIB. Journal of Cystic Fibrosis, 2004, 3, 165-172.	0.3	46
27	A rhizospheric Burkholderia cepacia complex population: genotypic and phenotypic diversity of Burkholderia cenocepacia and Burkholderia ambifaria. FEMS Microbiology Ecology, 2003, 46, 179-187.	1.3	24
28	Burkholderia cepacia Complex Bacteria from Clinical and Environmental Sources in Italy: Genomovar Status and Distribution of Traits Related to Virulence and Transmissibility. Journal of Clinical Microbiology, 2002, 40, 846-851.	1.8	87
29	Effects of two different application methods of Burkholderia ambifaria MCI 7 on plant growth and rhizospheric bacterial diversity. Environmental Microbiology, 2002, 4, 238-245.	1.8	69
30	Burkholderia cepacia complex: distribution of genomovars among isolates from the maize rhizosphere in Italy. Environmental Microbiology, 2001, 3, 137-143.	1.8	74
31	Different portions of the maize root system host Burkholderia cepacia populations with different degrees of genetic polymorphism. Environmental Microbiology, 2000, 2, 111-118.	1.8	19
32	Bias Caused by Using Different Isolation Media for Assessing the Genetic Diversity of a Natural Microbial Population. Microbial Ecology, 2000, 40, 169-176.	1.4	69
33	Efficacy of Burkholderia cepacia MCI 7 in disease suppression and growth promotion of maize. Biology and Fertility of Soils, 2000, 31, 225-231.	2.3	54
34	Soil Type and Maize Cultivar Affect the Genetic Diversity of Maize Root–Associated Burkholderia cepacia Populations. Microbial Ecology, 1999, 38, 273-284.	1.4	131
35	Characterization of a free-living maize-rhizosphere population of Burkholderia cepacia: effect of seed treatment on disease suppression and growth promotion of maize. FEMS Microbiology Ecology, 1998, 27, 225-237.	1.3	117
36	Inoculation of Burkholderia cepacia, Pseudomonas fluorescens and Enterobacter sp. on Sorghum bicolor: Root colonization and plant growth promotion of dual strain inocula. Soil Biology and Biochemistry, 1998, 30, 81-87.	4.2	70

3

#	Article	IF	CITATION
37	Influence of plant development, cultivar and soil type on microbial colonization of maize roots. Applied Soil Ecology, 1998, 8, 11-18.	2.1	98
38	Perturbation of maize rhizosphere microflora following seed bacterization with Burkholderia cepacia MCI 7. FEMS Microbiology Ecology, 1997, 23, 183-193.	1.3	7
39	Molecular characterization of rhizosphere and clinical isolates of Burkholderia cepacia. Research in Microbiology, 1995, 146, 531-542.	1.0	32
40	Influence of growth supplements on lactic acid production in whey ultrafiltrate by Lactobacillus helveticus. Applied Microbiology and Biotechnology, 1992, 36, 461.	1.7	65