

Hyeon-Seok Eom

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2626273/publications.pdf>

Version: 2024-02-01

74
papers

854
citations

623574

14
h-index

552653

26
g-index

74
all docs

74
docs citations

74
times ranked

1583
citing authors

#	ARTICLE	IF	CITATIONS
1	Nilotinib combined with multiagent chemotherapy for newly diagnosed Philadelphia-positive acute lymphoblastic leukemia. <i>Blood</i> , 2015, 126, 746-756.	0.6	160
2	Elevated Red Blood Cell Distribution Width as a Simple Prognostic Factor in Patients with Symptomatic Multiple Myeloma. <i>BioMed Research International</i> , 2014, 2014, 1-8.	0.9	62
3	Nationwide Statistical Analysis of Lymphoid Malignancies in Korea. <i>Cancer Research and Treatment</i> , 2018, 50, 222-238.	1.3	47
4	Nationwide statistical analysis of myeloid malignancies in Korea: incidence and survival rate from 1999 to 2012. <i>Blood Research</i> , 2015, 50, 204.	0.5	44
5	Multiple myeloma in Korea: past, present, and future perspectives. Experience of the Korean Multiple Myeloma Working Party. <i>International Journal of Hematology</i> , 2010, 92, 52-57.	0.7	41
6	Long-term follow-up of imatinib plus combination chemotherapy in patients with newly diagnosed Philadelphia chromosome-positive acute lymphoblastic leukemia. <i>American Journal of Hematology</i> , 2015, 90, 1013-1020.	2.0	37
7	Comprehensive evaluation of the revised international staging system in multiple myeloma patients treated with novel agents as a primary therapy. <i>American Journal of Hematology</i> , 2017, 92, 1280-1286.	2.0	34
8	Guillain-Barré syndrome after vaccination against COVID-19. <i>Lancet Neurology</i> , The, 2022, 21, 117-119.	4.9	27
9	Allogeneic Stem Cell Transplantation for Patients with Natural Killer/T Cell Lymphoid Malignancy: A Multicenter Analysis Comparing Upfront and Salvage Transplantation. <i>Biology of Blood and Marrow Transplantation</i> , 2018, 24, 2471-2478.	2.0	24
10	Post-Transplantation Natural Killer Cell Count: A Predictor of Acute Graft-Versus-Host Disease and Survival Outcomes After Allogeneic Hematopoietic Stem Cell Transplantation. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2016, 16, 527-535.e2.	0.2	23
11	Sequential High-Dose Dexamethasone and Response Adapted PAD (Bortezomib, Adriamycin, Ifosfamide, and Gemtuzumab) by High-Dose Chemotherapy with Autologous Stem Cell Transplantation for Newly Diagnosed Multiple Myeloma; Open-Labelled, Multicenter Phase 2 Study (KMM-94 Study)-Interim Analysis. <i>Blood</i> , 2011, 118, 2044-2044.	0.6	19
12	Expression of Myeloid Antigen in Neoplastic Plasma Cells Is Related to Adverse Prognosis in Patients with Multiple Myeloma. <i>BioMed Research International</i> , 2014, 2014, 1-8.	0.9	18
13	Prognostic role of the neutrophil-to-lymphocyte ratio in patients with primary central nervous system lymphoma. <i>Oncotarget</i> , 2017, 8, 74975-74986.	0.8	17
14	Phase I Clinical Trial of 4-1BB-based Adoptive T-Cell Therapy for Epstein-Barr Virus (EBV)-positive Tumors. <i>Journal of Immunotherapy</i> , 2016, 39, 140-148.	1.2	16
15	Prognostic Value of Platelet Count in Patients with Peripheral T Cell Lymphoma. <i>Acta Haematologica</i> , 2019, 141, 176-186.	0.7	16
16	Adverse Prognostic Impact of Bone Marrow Microvessel Density in Multiple Myeloma. <i>Annals of Laboratory Medicine</i> , 2015, 35, 563-569.	1.2	15
17	Highly elevated serum lactate dehydrogenase is associated with central nervous system relapse in patients with diffuse large B-cell lymphoma: Results of a multicenter prospective cohort study. <i>Oncotarget</i> , 2016, 7, 72033-72043.	0.8	14
18	Bortezomib, thalidomide, dexamethasone induction therapy followed by melphalan, prednisolone, thalidomide consolidation therapy as a first line of treatment for patients with multiple myeloma who are non-transplant candidates: results of the Korean Multiple Myeloma Working Party (KMMWP). <i>Annals of Hematology</i> , 2010, 89, 489-497.	0.8	12

#	ARTICLE	IF	CITATIONS
19	A Case of Disseminated Infection with Skin Manifestation due to Non- <i>neoformans</i> and Non- <i>gattii</i> <i>Cryptococcus</i> in a Patient with Refractory Acute Myeloid Leukemia. <i>Infection and Chemotherapy</i> , 2017, 49, 142.	1.0	12
20	Improved prognostic stratification using NCCN- and GELTAMO-international prognostic index in patients with diffuse large B-cell lymphoma. <i>Oncotarget</i> , 2017, 8, 92171-92182.	0.8	12
21	Zanubrutinib monotherapy in relapsed/refractory indolent non-Hodgkin lymphoma. <i>Blood Advances</i> , 2022, 6, 3472-3479.	2.5	12
22	Clinical features and outcomes of Hodgkin's lymphoma in Korea: Consortium for Improving Survival of Lymphoma (CISL). <i>Annals of Hematology</i> , 2012, 91, 223-233.	0.8	11
23	Efficacy of eculizumab in paroxysmal nocturnal hemoglobinuria patients with or without aplastic anemia: prospective study of a Korean PNH cohort. <i>Blood Research</i> , 2017, 52, 207.	0.5	11
24	R-CHOP chemoimmunotherapy followed by autologous transplantation for the treatment of diffuse large B-cell lymphoma. <i>Blood Research</i> , 2014, 49, 107.	0.5	10
25	Prognostic Implications of Monosomies in Patients With Multiple Myeloma. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2017, 17, 159-164.e2.	0.2	10
26	Phase 2 Study of an Intravenous Busulfan and Melphalan Conditioning Regimen for Autologous Stem Cell Transplantation in Patients with Multiple Myeloma (KMM150). <i>Biology of Blood and Marrow Transplantation</i> , 2018, 24, 923-929.	2.0	10
27	A multicenter phase II trial of etoposide, methylprednisolone, high-dose cytarabine, and oxaliplatin for patients with primary refractory/relapsed aggressive non-Hodgkin's lymphoma. <i>Cancer Chemotherapy and Pharmacology</i> , 2009, 64, 27-33.	1.1	9
28	The different roles of molecular classification according to upfront autologous stem cell transplantation in advanced-stage diffuse large B cell lymphoma patients with elevated serum lactate dehydrogenase. <i>Annals of Hematology</i> , 2016, 95, 1491-1501.	0.8	9
29	Radiation therapy for gastric mucosa-associated lymphoid tissue lymphoma: dose-volumetric analysis and its clinical implications. <i>Radiation Oncology Journal</i> , 2016, 34, 193-201.	0.7	8
30	Adverse prognostic impact of vascular endothelial growth factor gene polymorphisms in patients with diffuse large B-cell lymphoma. <i>Leukemia and Lymphoma</i> , 2017, 58, 2677-2682.	0.6	7
31	Sequential High-Dose Dexamethasone and Response Adapted PAD (Bortezomib, Adriamycin,) Tj ETQq1 1 0.784314 rgBT /Overlock 10 High-Dose Therapy with ASCT for Newly Diagnosed Multiple Myeloma; Open-Labeled, Multicenter Phase 2 Study (KMM-94 Study)-Interim Analysis. <i>Blood</i> , 2012, 120, 3106-3106.	0.6	7
32	Low-dose cisplatin converts the tumor microenvironment into a permissive state for HSVtk-induced antitumor immunity in HPV16-related tonsillar carcinoma. <i>Cancer Letters</i> , 2015, 356, 743-750.	3.2	6
33	Bendamustine in heavily pre-treated multiple myeloma patients: Results of a retrospective analysis from the Korean Multiple Myeloma Working Party. <i>Blood Research</i> , 2016, 51, 193.	0.5	6
34	Selection of a mobilization regimen for multiple myeloma based on the response to induction therapy: granulocyte-colony stimulating factor (G-CSF) alone versus high-dose cyclophosphamide plus G-CSF. <i>Leukemia and Lymphoma</i> , 2016, 57, 1389-1397.	0.6	6
35	High level of pre-treatment C-reactive protein to albumin ratio predicts inferior prognosis in diffuse large B-cell lymphoma. <i>Scientific Reports</i> , 2021, 11, 2674.	1.6	6
36	Recent Clinical Update of Acute Myeloid Leukemia: Focus on Epigenetic Therapies. <i>Journal of Korean Medical Science</i> , 2021, 36, e85.	1.1	6

#	ARTICLE	IF	CITATIONS
37	Efficacy and toxicity of the combination chemotherapy of thalidomide, alkylating agent, and steroid for relapsed/refractory myeloma patients: a report from the Korean Multiple Myeloma Working Party (KMMWP) retrospective study. <i>Cancer Medicine</i> , 2017, 6, 100-108.	1.3	5
38	Phase I study of CKD-581, a pan-histone deacetylase inhibitor, in patients with lymphoma or multiple myeloma refractory to standard therapy. <i>Investigational New Drugs</i> , 2018, 36, 877-885.	1.2	5
39	Autologous stem cell transplantation in elderly patients with multiple myeloma in Korea: the KMM1807 study. <i>International Journal of Hematology</i> , 2020, 112, 84-95.	0.7	5
40	Use of lenalidomide in the management of relapsed or refractory multiple myeloma: expert recommendations in Korea. <i>Blood Research</i> , 2015, 50, 7.	0.5	4
41	A phase II study of oxaliplatin and prednisone for patients with relapsed or refractory marginal zone lymphoma: Consortium for Improving Survival of Lymphoma trial. <i>Leukemia and Lymphoma</i> , 2016, 57, 1406-1412.	0.6	4
42	Analysis of laboratory parameters for optimal autologous peripheral blood stem cell collection from lymphoma and myeloma patients. <i>Journal of Clinical Apheresis</i> , 2021, 36, 135-142.	0.7	4
43	Comparison of spectra optia and amicus cell separators for autologous peripheral blood stem cell collection. <i>Journal of Clinical Apheresis</i> , 2021, 36, 28-33.	0.7	4
44	Real-world outcomes of ibrutinib therapy in Korean patients with relapsed or refractory mantle cell lymphoma: a multicenter, retrospective analysis. <i>Cancer Communications</i> , 2021, 41, 275-278.	3.7	4
45	Sequential VAD (Vincristine, Adriamycin, Dexamethasone) and VTD (VELCADE®, Thalidomide,) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Transplantation and Maintenance Treatment with VELCADE for Newly Diagnosed Multiple Myeloma: Interim Results of Phase II Trial.. <i>Blood</i> , 2007, 110, 952-952.	0.6	4
46	Nilotinib Combined With Multi-Agent Chemotherapy For Adult Patients With Newly Diagnosed Philadelphia Chromosome-Positive Acute Lymphoblastic Leukemia: Final Results Of Prospective Multicenter Phase 2 Study. <i>Blood</i> , 2013, 122, 55-55.	0.6	4
47	Pegfilgrastim Prophylaxis is Effective in the Prevention of Febrile Neutropenia and Reduces Mortality in Patients Aged ≥75 Years with Diffuse Large B-Cell Lymphoma Treated with R-CHOP: A Prospective Cohort Study. <i>Cancer Research and Treatment</i> , 2021, , .	1.3	4
48	Prediction of survival by applying current prognostic models in diffuse large B-cell lymphoma treated with R-CHOP followed by autologous transplantation. <i>Blood Research</i> , 2015, 50, 160.	0.5	3
49	A phase II study of etoposide, methylprednisolone, high-dose cytarabine, and oxaliplatin (ESHAOx) for patients with refractory or relapsed Hodgkin's lymphoma. <i>Annals of Hematology</i> , 2020, 99, 255-264.	0.8	3
50	Analysis of the Efficacy of Thalidomide Plus Dexamethasone-Based Regimens in Patients With Relapsed/Refractory Multiple Myeloma Who Received Prior Chemotherapy, Including Bortezomib and Lenalidomide: KMM-166 Study. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2020, 20, e97-e104.	0.2	2
51	Current Use of Total Body Irradiation in Haploidentical Allogeneic Hematopoietic Stem Cell Transplantation. <i>Journal of Korean Medical Science</i> , 2021, 36, e55.	1.1	2
52	The Efficacy of Bortezomib-CHOP In Patients with Advanced Stage T or NK/T Cell Lymphomas: The Results of Multicenter Phase II Study.. <i>Blood</i> , 2010, 116, 1791-1791.	0.6	2
53	Phase I Study of mTOR Inhibitor Everolimus Plus CHOP in Patients with Advanced, Aggressive T-Cell lymphomas. <i>Blood</i> , 2011, 118, 1642-1642.	0.6	2
54	A Retrospective Study to Evaluate the Survival Rates in R-CHOP Chemotherapy Followed by Autologous Stem Cell Transplantation for the Treatment of Diffuse Large B Cell Lymphoma. <i>Blood</i> , 2012, 120, 4517-4517.	0.6	2

#	ARTICLE	IF	CITATIONS
55	The prognostic impact of inflammatory factors in patients with multiple myeloma treated with thalidomide in Korea. Korean Journal of Internal Medicine, 2015, 30, 675-683.	0.7	2
56	Influence of creatinine levels on survival in patients with veno-occlusive disease treated with defibrotide. Korean Journal of Internal Medicine, 2022, 37, 179-189.	0.7	2
57	Outcomes in Refractory Diffuse Large B-Cell Lymphoma: Results from Two Prospective Korean Cohorts. Cancer Research and Treatment, 2023, 55, 325-333.	1.3	2
58	Impact of Renal Impairment (RI) on Outcomes after Treatment (Tx) with Lenalidomide and Low-Dose Dexamethasone (Rd) in Newly Diagnosed Multiple Myeloma (NDMM) Patients (Pts): First Trial Results. Blood, 2014, 124, 2112-2112.	0.6	1
59	Rapid Sequential Gain of <i>ABL1</i> Kinase Domain Mutations with a Complex Karyotype in the Progression of Chronic Myelogenous Leukemia. Annals of Laboratory Medicine, 2014, 34, 399-401.	1.2	0
60	Development of an Automated Image Analyzer for Microvessel Density Measurement in Bone Marrow Biopsies. Annals of Laboratory Medicine, 2020, 40, 312-316.	1.2	0
61	Limited benefits of thalidomide and dexamethasone maintenance after autologous stem cell transplantation in newly diagnosed multiple myeloma patients: a prospective phase II multi-center study in Korea. Current Problems in Cancer, 2021, 46, 100786.	1.0	0
62	A Multicenter Phase II Trial of Etoposide, Methylprednisolone, High-Dose Cytarabine, and Oxaliplatin (ESHAOx) for Patients with Refractory/Relapsed Aggressive Non-Hodgkin's Lymphoma.. Blood, 2007, 110, 3446-3446.	0.6	0
63	Fluorescent in Situ Hybridization Analysis at Bone Marrow Biopsy Section in Multiple Myeloma.. Blood, 2009, 114, 4898-4898.	0.6	0
64	Concurrent Chemo-Radiotherapy Followed by VIDL (Etoposide, Ifosfamide, Dexamethasone,) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 387 1 Blood, 2010, 116, 1765-1765.	0.6	0
65	Prognostic Significance of Immune Thrombocytopenic Purpura In Patients with Malignancy.. Blood, 2010, 116, 1428-1428.	0.6	0
66	Effects of Prophylactic Defibrotide for Veno-Occlusive Disease in Hematopoietic Stem Cell Transplantation. Blood, 2011, 118, 4517-4517.	0.6	0
67	Response Assessment Using Early and End-of-Treatment 18F-FLT PET Can Predict Outcome in Patients with Non-Hodgkin Lymphoma,. Blood, 2011, 118, 3679-3679.	0.6	0
68	Matched-Pair Analysis Comparing Outcomes of Second Autologous Stem Cell Transplantation and Chemotherapy As a Salvage Therapy in Patients with Multiple Myeloma Who Relapsed After Front-Line Autologous Stem Cell Transplantation. Blood, 2011, 118, 1990-1990.	0.6	0
69	Bortezomib Induction Followed by ASCT in Patients with Multiple Myeloma: Achievement of Response After Induction and Achieving CR Post-ASCT Are Both Important Prognostic Markers. Blood, 2011, 118, 1866-1866.	0.6	0
70	Early Response to Bortezomib Combined Chemotherapy Can Be Helpful for Prediction of Progression Free Survival in Patients with Multiple Myeloma Who Were Ineligible for Stem Cell Transplantation. Blood, 2011, 118, 5138-5138.	0.6	0
71	Adverse Prognostic Impact of Vascular Endothelial Growth Factor Gene Polymorphism in Patients with Non-Hodgkin Lymphoma.. Blood, 2012, 120, 2674-2674.	0.6	0
72	Hematopoietic Progenitor Cell Counts Is a Useful Indicator To Determine Appropriate Time For Peripheral Blood Stem Cell Collection. Blood, 2013, 122, 2037-2037.	0.6	0

#	ARTICLE	IF	CITATIONS
73	A Case of Disseminated Infection with Skin Manifestation due to Non- <i>neofomans</i> and Non- <i>gattii</i> <i>Cryptococcus</i> in a Patient with Refractory Acute Myeloid Leukemia. <i>Infection and Chemotherapy</i> , 2017, 49, 142.	1.0	0
74	Characteristics and clinical outcome of high-risk multiple myeloma patients in Korea (KMM 1805). <i>International Journal of Hematology</i> , 2022, , 1.	0.7	0