

List of Publications by Citations

Source: <https://exaly.com/author-pdf/2628320/andrew-green-publications-by-citations.pdf>
Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

335 papers	21,256 citations	60 h-index	138 g-index
351 ext. papers	25,385 ext. citations	6.8 avg, IF	6.38 L-index

#	Paper	IF	Citations
335	The genomic and transcriptomic architecture of 2,000 breast tumours reveals novel subgroups. <i>Nature</i> , 2012 , 486, 346-52	50.4	3479
334	Differential oestrogen receptor binding is associated with clinical outcome in breast cancer. <i>Nature</i> , 2012 , 481, 389-93	50.4	1011
333	Prognostic markers in triple-negative breast cancer. <i>Cancer</i> , 2007 , 109, 25-32	6.4	963
332	Tumor-infiltrating CD8+ lymphocytes predict clinical outcome in breast cancer. <i>Journal of Clinical Oncology</i> , 2011 , 29, 1949-55	2.2	961
331	The somatic mutation profiles of 2,433 breast cancers refines their genomic and transcriptomic landscapes. <i>Nature Communications</i> , 2016 , 7, 11479	17.4	779
330	MicroRNA expression profiling of human breast cancer identifies new markers of tumor subtype. <i>Genome Biology</i> , 2007 , 8, R214	18.3	742
329	Subtyping of breast cancer by immunohistochemistry to investigate a relationship between subtype and short and long term survival: a collaborative analysis of data for 10,159 cases from 12 studies. <i>PLoS Medicine</i> , 2010 , 7, e1000279	11.6	616
328	Prognostic value of a combined estrogen receptor, progesterone receptor, Ki-67, and human epidermal growth factor receptor 2 immunohistochemical score and comparison with the Genomic Health recurrence score in early breast cancer. <i>Journal of Clinical Oncology</i> , 2011 , 29, 4273-8	2.2	549
327	Triple-negative breast cancer: distinguishing between basal and nonbasal subtypes. <i>Clinical Cancer Research</i> , 2009 , 15, 2302-10	12.9	371
326	Global histone modifications in breast cancer correlate with tumor phenotypes, prognostic factors, and patient outcome. <i>Cancer Research</i> , 2009 , 69, 3802-9	10.1	340
325	Beta-Blocker Drug Therapy Reduces Secondary Cancer Formation in Breast Cancer and Improves Cancer Specific Survival. <i>Oncotarget</i> , 2010 , 1, 628-638	3.3	332
324	The shaping and functional consequences of the microRNA landscape in breast cancer. <i>Nature</i> , 2013 , 497, 378-82	50.4	321
323	Expression of mucins (MUC1, MUC2, MUC3, MUC4, MUC5AC and MUC6) and their prognostic significance in human breast cancer. <i>Modern Pathology</i> , 2005 , 18, 1295-304	9.8	257
322	High-resolution aCGH and expression profiling identifies a novel genomic subtype of ER negative breast cancer. <i>Genome Biology</i> , 2007 , 8, R215	18.3	230
321	BCL2 in breast cancer: a favourable prognostic marker across molecular subtypes and independent of adjuvant therapy received. <i>British Journal of Cancer</i> , 2010 , 103, 668-75	8.7	218
320	Biologic and clinical characteristics of breast cancer with single hormone receptor positive phenotype. <i>Journal of Clinical Oncology</i> , 2007 , 25, 4772-8	2.2	213
319	FGFR1 amplification in breast carcinomas: a chromogenic in situ hybridisation analysis. <i>Breast Cancer Research</i> , 2007 , 9, R23	8.3	211

318	A gene-expression signature to predict survival in breast cancer across independent data sets. <i>Oncogene</i> , 2007 , 26, 1507-16	9.2	191
317	Nuclear and cytoplasmic expression of ERbeta1, ERbeta2, and ERbeta5 identifies distinct prognostic outcome for breast cancer patients. <i>Clinical Cancer Research</i> , 2008 , 14, 5228-35	12.9	187
316	The prognostic significance of B lymphocytes in invasive carcinoma of the breast. <i>Breast Cancer Research and Treatment</i> , 2012 , 132, 545-53	4.4	181
315	Invasive lobular carcinoma of the breast: response to hormonal therapy and outcomes. <i>European Journal of Cancer</i> , 2008 , 44, 73-83	7.5	164
314	Basal phenotype identifies a poor prognostic subgroup of breast cancer of clinical importance. <i>European Journal of Cancer</i> , 2006 , 42, 3149-56	7.5	164
313	Transferrin receptor (CD71) is a marker of poor prognosis in breast cancer and can predict response to tamoxifen. <i>Breast Cancer Research and Treatment</i> , 2010 , 119, 283-93	4.4	155
312	Prognostic significance of vascular endothelial cell growth factors -A, -C and -D in breast cancer and their relationship with angio- and lymphangiogenesis. <i>British Journal of Cancer</i> , 2007 , 96, 1092-100	8.7	153
311	Improved methods of detection of lymphovascular invasion demonstrate that it is the predominant method of vascular invasion in breast cancer and has important clinical consequences. <i>American Journal of Surgical Pathology</i> , 2007 , 31, 1825-33	6.7	142
310	Breast carcinoma with basal differentiation: a proposal for pathology definition based on basal cytokeratin expression. <i>Histopathology</i> , 2007 , 50, 434-8	7.3	132
309	PREDICT Plus: development and validation of a prognostic model for early breast cancer that includes HER2. <i>British Journal of Cancer</i> , 2012 , 107, 800-7	8.7	130
308	An evaluation of the clinical significance of FOXP3+ infiltrating cells in human breast cancer. <i>Breast Cancer Research and Treatment</i> , 2011 , 127, 99-108	4.4	129
307	Caveolin 1 and Caveolin 2 are associated with breast cancer basal-like and triple-negative immunophenotype. <i>British Journal of Cancer</i> , 2008 , 99, 327-34	8.7	122
306	Expression of BRCA1 protein in breast cancer and its prognostic significance. <i>Human Pathology</i> , 2008 , 39, 857-65	3.7	115
305	Expression of the stress-related MHC class I chain-related protein MICA is an indicator of good prognosis in colorectal cancer patients. <i>International Journal of Cancer</i> , 2006 , 118, 1445-52	7.5	112
304	Clinical and biological significance of E-cadherin protein expression in invasive lobular carcinoma of the breast. <i>American Journal of Surgical Pathology</i> , 2010 , 34, 1472-9	6.7	110
303	CCND1 amplification and cyclin D1 expression in breast cancer and their relation with proteomic subgroups and patient outcome. <i>Breast Cancer Research and Treatment</i> , 2008 , 109, 325-35	4.4	110
302	Dysregulated expression of Fau and MELK is associated with poor prognosis in breast cancer. <i>Breast Cancer Research</i> , 2009 , 11, R60	8.3	109
301	Dynamics of breast-cancer relapse reveal late-recurring ER-positive genomic subgroups. <i>Nature</i> , 2019 , 567, 399-404	50.4	108

300	Expression of cytokine messenger RNA in normal and neoplastic human breast tissue: identification of interleukin-8 as a potential regulatory factor in breast tumours. <i>International Journal of Cancer</i> , 1997 , 72, 937-41	7.5	105
299	Kinome screening for regulators of the estrogen receptor identifies LMTK3 as a new therapeutic target in breast cancer. <i>Nature Medicine</i> , 2011 , 17, 715-9	50.5	101
298	An updated PREDICT breast cancer prognostication and treatment benefit prediction model with independent validation. <i>Breast Cancer Research</i> , 2017 , 19, 58	8.3	100
297	Adenomyosis--a result of disordered stromal differentiation. <i>American Journal of Pathology</i> , 2001 , 159, 623-30	5.8	99
296	Histologic grading is an independent prognostic factor in invasive lobular carcinoma of the breast. <i>Breast Cancer Research and Treatment</i> , 2008 , 111, 121-7	4.4	96
295	Prognostic value of proliferation assay in the luminal, HER2-positive, and triple-negative biologic classes of breast cancer. <i>Breast Cancer Research</i> , 2012 , 14, R3	8.3	94
294	Therapeutic targeting of integrin $\alpha 5$ in breast cancer. <i>Journal of the National Cancer Institute</i> , 2014 , 106,	9.7	89
293	Caspase-3 and caspase-8 expression in breast cancer: caspase-3 is associated with survival. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2017 , 22, 357-368	5.4	85
292	Lymphatic and blood vessels in basal and triple-negative breast cancers: characteristics and prognostic significance. <i>Modern Pathology</i> , 2011 , 24, 774-85	9.8	84
291	NapA protects <i>Helicobacter pylori</i> from oxidative stress damage, and its production is influenced by the ferric uptake regulator. <i>Journal of Medical Microbiology</i> , 2003 , 52, 461-469	3.2	84
290	MIB1/Ki-67 labelling index can classify grade 2 breast cancer into two clinically distinct subgroups. <i>Breast Cancer Research and Treatment</i> , 2011 , 127, 591-9	4.4	82
289	A CD44 ⁺ /CD24 ⁺ phenotype is a poor prognostic marker in early invasive breast cancer. <i>Breast Cancer Research and Treatment</i> , 2012 , 133, 979-95	4.4	79
288	Alpha- and beta-adrenergic receptor (AR) protein expression is associated with poor clinical outcome in breast cancer: an immunohistochemical study. <i>Breast Cancer Research and Treatment</i> , 2011 , 130, 457-63	4.4	70
287	Loss of Dicer expression is associated with breast cancer progression and recurrence. <i>Breast Cancer Research and Treatment</i> , 2012 , 135, 403-13	4.4	69
286	Tumor size is an unreliable predictor of prognosis in basal-like breast cancers and does not correlate closely with lymph node status. <i>Breast Cancer Research and Treatment</i> , 2009 , 117, 199-204	4.4	69
285	Chromosome 16 tumor-suppressor genes in breast cancer. <i>Genes Chromosomes and Cancer</i> , 2006 , 45, 527-35	5	69
284	Nottingham Prognostic Index Plus (NPI+): a modern clinical decision making tool in breast cancer. <i>British Journal of Cancer</i> , 2014 , 110, 1688-97	8.7	68
283	The collagen receptor Endo180 (CD280) is expressed on basal-like breast tumor cells and promotes tumor growth in vivo. <i>Cancer Research</i> , 2007 , 67, 10230-40	10.1	68

282	Determination of HER2 amplification in primary breast cancer using dual-colour chromogenic in situ hybridization is comparable to fluorescence in situ hybridization: a European multicentre study involving 168 specimens. <i>Histopathology</i> , 2010 , 56, 472-80	7.3	67
281	The prognostic significance of PELP1 expression in invasive breast cancer with emphasis on the ER-positive luminal-like subtype. <i>Breast Cancer Research and Treatment</i> , 2010 , 120, 603-12	4.4	66
280	SPAG5 as a prognostic biomarker and chemotherapy sensitivity predictor in breast cancer: a retrospective, integrated genomic, transcriptomic, and protein analysis. <i>Lancet Oncology</i> , 2016 , 17, 1004-1018	21.7	65
279	MYC functions are specific in biological subtypes of breast cancer and confers resistance to endocrine therapy in luminal tumours. <i>British Journal of Cancer</i> , 2016 , 114, 917-28	8.7	64
278	Objective assessment of lymphatic and blood vascular invasion in lymph node-negative breast carcinoma: findings from a large case series with long-term follow-up. <i>Journal of Pathology</i> , 2011 , 223, 358-65	9.4	63
277	PIK3CA expression in invasive breast cancer: a biomarker of poor prognosis. <i>Breast Cancer Research and Treatment</i> , 2010 , 122, 45-53	4.4	63
276	Clinical and biological significance of glucocorticoid receptor (GR) expression in breast cancer. <i>Breast Cancer Research and Treatment</i> , 2015 , 150, 335-46	4.4	59
275	Prognostic significance of androgen receptor expression in invasive breast cancer: transcriptomic and protein expression analysis. <i>Breast Cancer Research and Treatment</i> , 2016 , 159, 215-27	4.4	59
274	FOXO3a nuclear localisation is associated with good prognosis in luminal-like breast cancer. <i>Breast Cancer Research and Treatment</i> , 2011 , 129, 11-21	4.4	59
273	Genomic gain of 5p15 leads to over-expression of Misu (NSUN2) in breast cancer. <i>Cancer Letters</i> , 2010 , 289, 71-80	9.9	59
272	The amino acid transporter SLC7A5 confers a poor prognosis in the highly proliferative breast cancer subtypes and is a key therapeutic target in luminal B tumours. <i>Breast Cancer Research</i> , 2018 , 20, 21	8.3	58
271	Untangling the ATR-CHEK1 network for prognostication, prediction and therapeutic target validation in breast cancer. <i>Molecular Oncology</i> , 2015 , 9, 569-85	7.9	57
270	Notch-1-PTEN-ERK1/2 signaling axis promotes HER2+ breast cancer cell proliferation and stem cell survival. <i>Oncogene</i> , 2018 , 37, 4489-4504	9.2	57
269	Targeting BRCA1-BER deficient breast cancer by ATM or DNA-PKcs blockade either alone or in combination with cisplatin for personalized therapy. <i>Molecular Oncology</i> , 2015 , 9, 204-17	7.9	55
268	Immune Infiltration in Invasive Lobular Breast Cancer. <i>Journal of the National Cancer Institute</i> , 2018 , 110, 768-776	9.7	55
267	Activity and gene expression of 17beta-hydroxysteroid dehydrogenase type I in primary cultures of epithelial and stromal cells derived from normal and tumorous human breast tissue: the role of IL-8. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 1998 , 67, 267-74	5.1	55
266	Molecular classification of breast cancer: what the pathologist needs to know. <i>Pathology</i> , 2017 , 49, 111-119	11.9	54
265	Understanding the HER family in breast cancer: interaction with ligands, dimerization and treatments. <i>Histopathology</i> , 2010 , 56, 560-72	7.3	52

264	Guidelines and considerations for conducting experiments using tissue microarrays. <i>Histopathology</i> , 2013 , 62, 827-39	7.3	50
263	Biology of primary breast cancer in older women treated by surgery: with correlation with long-term clinical outcome and comparison with their younger counterparts. <i>British Journal of Cancer</i> , 2013 , 108, 1042-51	8.7	49
262	p53 status identifies two subgroups of triple-negative breast cancers with distinct biological features. <i>Japanese Journal of Clinical Oncology</i> , 2011 , 41, 172-9	2.8	49
261	Calpain system protein expression in basal-like and triple-negative invasive breast cancer. <i>Annals of Oncology</i> , 2012 , 23, 2289-2296	10.3	49
260	The biological, clinical and prognostic implications of p53 transcriptional pathways in breast cancers. <i>Journal of Pathology</i> , 2010 , 220, 419-34	9.4	49
259	The prognostic significance of steroid receptor co-regulators in breast cancer: co-repressor NCOR2/SMRT is an independent indicator of poor outcome. <i>Breast Cancer Research and Treatment</i> , 2008 , 110, 427-37	4.4	49
258	Biological and clinical significance of PARP1 protein expression in breast cancer. <i>Breast Cancer Research and Treatment</i> , 2015 , 149, 353-62	4.4	48
257	Sonographic correlations with the new molecular classification of invasive breast cancer. <i>European Radiology</i> , 2009 , 19, 2342-8	8	48
256	A validated gene expression profile for detecting clinical outcome in breast cancer using artificial neural networks. <i>Breast Cancer Research and Treatment</i> , 2010 , 120, 83-93	4.4	47
255	MYC regulation of glutamine-proline regulatory axis is key in luminal B breast cancer. <i>British Journal of Cancer</i> , 2018 , 118, 258-265	8.7	47
254	Clinicopathological significance of KU70/KU80, a key DNA damage repair protein in breast cancer. <i>Breast Cancer Research and Treatment</i> , 2013 , 139, 301-10	4.4	46
253	IL6/STAT3 Signaling Hijacks Estrogen Receptor Enhancers to Drive Breast Cancer Metastasis. <i>Cancer Cell</i> , 2020 , 38, 412-423.e9	24.3	46
252	Clinicopathologic and molecular significance of phospho-Akt expression in early invasive breast cancer. <i>Breast Cancer Research and Treatment</i> , 2011 , 127, 407-16	4.4	45
251	Epithelial mesenchymal transition in early invasive breast cancer: an immunohistochemical and reverse phase protein array study. <i>Breast Cancer Research and Treatment</i> , 2014 , 145, 339-48	4.4	44
250	A whole-genome massively parallel sequencing analysis of BRCA1 mutant oestrogen receptor-negative and -positive breast cancers. <i>Journal of Pathology</i> , 2012 , 227, 29-41	9.4	44
249	ELF5 Drives Lung Metastasis in Luminal Breast Cancer through Recruitment of Gr1+ CD11b+ Myeloid-Derived Suppressor Cells. <i>PLoS Biology</i> , 2015 , 13, e1002330	9.7	44
248	KPNA2 is a nuclear export protein that contributes to aberrant localisation of key proteins and poor prognosis of breast cancer. <i>British Journal of Cancer</i> , 2015 , 112, 1929-37	8.7	43
247	The proteins FABP7 and OATP2 are associated with the basal phenotype and patient outcome in human breast cancer. <i>Breast Cancer Research and Treatment</i> , 2010 , 121, 41-51	4.4	43

246	Neonatal tamoxifen treatment of mice leads to adenomyosis but not uterine cancer. <i>Experimental and Toxicologic Pathology</i> , 2005 , 56, 255-63		43
245	Small molecule inhibition of group I p21-activated kinases in breast cancer induces apoptosis and potentiates the activity of microtubule stabilizing agents. <i>Breast Cancer Research</i> , 2015 , 17, 59	8.3	42
244	Loss of expression of chromosome 16q genes DPEP1 and CTCF in lobular carcinoma in situ of the breast. <i>Breast Cancer Research and Treatment</i> , 2009 , 113, 59-66	4.4	42
243	A methodology to identify consensus classes from clustering algorithms applied to immunohistochemical data from breast cancer patients. <i>Computers in Biology and Medicine</i> , 2010 , 40, 318-30	7	42
242	Clinical Impact of Tumor DNA Repair Expression and T-cell Infiltration in Breast Cancers. <i>Cancer Immunology Research</i> , 2017 , 5, 292-299	12.5	40
241	IL-6 and IL-10 are associated with good prognosis in early stage invasive breast cancer patients. <i>Cancer Immunology, Immunotherapy</i> , 2018 , 67, 537-549	7.4	40
240	Prognostic significance of tumor-infiltrating lymphocytes in ductal carcinoma in situ of the breast. <i>Modern Pathology</i> , 2018 , 31, 1226-1236	9.8	40
239	Involvement of metformin and AMPK in the radioresponse and prognosis of luminal versus basal-like breast cancer treated with radiotherapy. <i>Oncotarget</i> , 2014 , 5, 12936-49	3.3	39
238	The role of glutaminase in cancer. <i>Histopathology</i> , 2020 , 76, 498-508	7.3	39
237	C-Met in invasive breast cancer: is there a relationship with the basal-like subtype?. <i>Cancer</i> , 2014 , 120, 163-71	6.4	38
236	Molecular characteristics and prognostic features of breast cancer in Nigerian compared with UK women. <i>Breast Cancer Research and Treatment</i> , 2012 , 135, 555-69	4.4	38
235	Heterogeneity of tumour-infiltrating lymphocytes in breast cancer and its prognostic significance. <i>Histopathology</i> , 2018 , 73, 887-896	7.3	38
234	CTEN (C-terminal tensin-like), a novel oncogene overexpressed in invasive breast carcinoma of poor prognosis. <i>Breast Cancer Research and Treatment</i> , 2011 , 126, 47-54	4.4	37
233	Investigating AP-2 and YY1 protein expression as a cause of high HER2 gene transcription in breast cancers with discordant HER2 gene amplification. <i>Breast Cancer Research</i> , 2009 , 11, R90	8.3	37
232	Evaluation of CDK12 Protein Expression as a Potential Novel Biomarker for DNA Damage Response-Targeted Therapies in Breast Cancer. <i>Molecular Cancer Therapeutics</i> , 2018 , 17, 306-315	6.1	37
231	Prognostic significance of tumour infiltrating B lymphocytes in breast ductal carcinoma in situ. <i>Histopathology</i> , 2017 , 71, 258-268	7.3	36
230	Breast carcinoma with basal phenotype: mammographic findings. <i>American Journal of Roentgenology</i> , 2008 , 191, 346-51	5.4	36
229	A tumor DNA complex aberration index is an independent predictor of survival in breast and ovarian cancer. <i>Molecular Oncology</i> , 2015 , 9, 115-27	7.9	35

228	Long-term clinical outcome of oestrogen receptor-positive operable primary breast cancer in older women: a large series from a single centre. <i>British Journal of Cancer</i> , 2011 , 104, 1393-400	8.7	35
227	Identification of key clinical phenotypes of breast cancer using a reduced panel of protein biomarkers. <i>British Journal of Cancer</i> , 2013 , 109, 1886-94	8.7	34
226	Lymph-node metastases in invasive lobular carcinoma are different from those in ductal carcinoma of the breast. <i>Journal of Clinical Pathology</i> , 2011 , 64, 995-1000	3.9	34
225	Proposal for a modified grading system based on mitotic index and Bcl2 provides objective determination of clinical outcome for patients with breast cancer. <i>Journal of Pathology</i> , 2010 , 222, 388-994	9.4	34
224	Ki67 expression in invasive breast cancer: the use of tissue microarrays compared with whole tissue sections. <i>Breast Cancer Research and Treatment</i> , 2017 , 164, 341-348	4.4	33
223	Inclusion of KI67 significantly improves performance of the PREDICT prognostication and prediction model for early breast cancer. <i>BMC Cancer</i> , 2014 , 14, 908	4.8	33
222	RERG (Ras-like, oestrogen-regulated, growth-inhibitor) expression in breast cancer: a marker of ER-positive luminal-like subtype. <i>Breast Cancer Research and Treatment</i> , 2011 , 128, 315-26	4.4	33
221	Encapsulated papillary carcinoma of the breast: a study of invasion associated markers. <i>Journal of Clinical Pathology</i> , 2012 , 65, 710-4	3.9	33
220	CD8(+) T lymphocytes infiltrating breast cancer: A promising new prognostic marker?. <i>Oncolmmunology</i> , 2012 , 1, 364-365	7.2	33
219	Overexpression of the cancer stem cell marker CD133 confers a poor prognosis in invasive breast cancer. <i>Breast Cancer Research and Treatment</i> , 2019 , 174, 387-399	4.4	33
218	Altered glutamine metabolism in breast cancer; subtype dependencies and alternative adaptations. <i>Histopathology</i> , 2018 , 72, 183-190	7.3	33
217	Transcriptomic and Protein Expression Analysis Reveals Clinicopathological Significance of Bloom Syndrome Helicase (BLM) in Breast Cancer. <i>Molecular Cancer Therapeutics</i> , 2015 , 14, 1057-65	6.1	32
216	Histological grading of breast cancer on needle core biopsy: the role of immunohistochemical assessment of proliferation. <i>Histopathology</i> , 2010 , 57, 212-9	7.3	32
215	DNA damage response markers are differentially expressed in BRCA-mutated breast cancers. <i>Breast Cancer Research and Treatment</i> , 2015 , 150, 81-90	4.4	30
214	Poly(adenosine diphosphate-ribose) polymerase expression in BRCA-proficient ovarian high-grade serous carcinoma; association with patient survival. <i>Human Pathology</i> , 2013 , 44, 1638-47	3.7	30
213	Clinicopathological significance of ATM-Chk2 expression in sporadic breast cancers: a comprehensive analysis in large cohorts. <i>Neoplasia</i> , 2014 , 16, 982-91	6.4	30
212	Molecular Aspects and Future Perspectives of Cytokine-Based Anti-cancer Immunotherapy. <i>Frontiers in Cell and Developmental Biology</i> , 2020 , 8, 402	5.7	29
211	RECQL4 helicase has oncogenic potential in sporadic breast cancers. <i>Journal of Pathology</i> , 2016 , 238, 495-501	9.4	29

210	LMTK3 expression in breast cancer: association with tumor phenotype and clinical outcome. <i>Breast Cancer Research and Treatment</i> , 2012 , 132, 537-44	4.4	29
209	Calpain-1 expression is associated with relapse-free survival in breast cancer patients treated with trastuzumab following adjuvant chemotherapy. <i>International Journal of Cancer</i> , 2011 , 129, 1773-80	7.5	29
208	Growth fraction as a predictor of response to chemotherapy in node-negative breast cancer. <i>International Journal of Cancer</i> , 2010 , 126, 1761-9	7.5	29
207	Checkpoint kinase1 (CHK1) is an important biomarker in breast cancer having a role in chemotherapy response. <i>British Journal of Cancer</i> , 2015 , 112, 901-11	8.7	28
206	Characteristics of basal cytokeratin expression in breast cancer. <i>Breast Cancer Research and Treatment</i> , 2013 , 139, 23-37	4.4	28
205	The cadherin switch in ovarian high-grade serous carcinoma is associated with disease progression. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2011 , 459, 21-9	5.1	28
204	The pioneer factor PBX1 is a novel driver of metastatic progression in ER α -positive breast cancer. <i>Oncotarget</i> , 2015 , 6, 21878-91	3.3	28
203	Clinical and biological significance of RAD51 expression in breast cancer: a key DNA damage response protein. <i>Breast Cancer Research and Treatment</i> , 2016 , 159, 41-53	4.4	28
202	Prognostic Role of Androgen Receptor in Triple Negative Breast Cancer: A Multi-Institutional Study. <i>Cancers</i> , 2019 , 11,	6.6	27
201	Microcephalin is a new novel prognostic indicator in breast cancer associated with BRCA1 inactivation. <i>Breast Cancer Research and Treatment</i> , 2011 , 127, 639-48	4.4	27
200	Chk1 phosphorylated at serine345 is a predictor of early local recurrence and radio-resistance in breast cancer. <i>Molecular Oncology</i> , 2016 , 10, 213-23	7.9	26
199	SUMOylation proteins in breast cancer. <i>Breast Cancer Research and Treatment</i> , 2014 , 144, 519-30	4.4	26
198	HER2/HER3 heterodimers and p21 expression are capable of predicting adjuvant trastuzumab response in HER2+ breast cancer. <i>Breast Cancer Research and Treatment</i> , 2014 , 145, 33-44	4.4	26
197	The kinase LMTK3 promotes invasion in breast cancer through GRB2-mediated induction of integrin α 5 β 1. <i>Science Signaling</i> , 2014 , 7, ra58	8.8	26
196	Clinicopathological and prognostic significance of RECQL5 helicase expression in breast cancers. <i>Carcinogenesis</i> , 2016 , 37, 63-71	4.6	25
195	Expression of Lamin A/C in early-stage breast cancer and its prognostic value. <i>Breast Cancer Research and Treatment</i> , 2019 , 174, 661-668	4.4	25
194	Low expression of G protein-coupled oestrogen receptor 1 (GPER) is associated with adverse survival of breast cancer patients. <i>Oncotarget</i> , 2018 , 9, 25946-25956	3.3	25
193	Inhibition of HER2 Increases JAGGED1-dependent Breast Cancer Stem Cells: Role for Membrane JAGGED1. <i>Clinical Cancer Research</i> , 2018 , 24, 4566-4578	12.9	24

192	Calpastatin is associated with lymphovascular invasion in breast cancer. <i>Breast</i> , 2011 , 20, 413-8	3.6	24
191	Impact of tissue sampling on accuracy of Ki67 immunohistochemistry evaluation in breast cancer. <i>Diagnostic Pathology</i> , 2016 , 11, 82	3	24
190	Nottingham Prognostic Index Plus: Validation of a clinical decision making tool in breast cancer in an independent series. <i>Journal of Pathology: Clinical Research</i> , 2016 , 2, 32-40	5.3	24
189	The multifunctional solute carrier 3A2 (SLC3A2) confers a poor prognosis in the highly proliferative breast cancer subtypes. <i>British Journal of Cancer</i> , 2018 , 118, 1115-1122	8.7	23
188	Further evidence that E-cadherin is not a tumour suppressor gene in invasive ductal carcinoma of the breast: an immunohistochemical study. <i>Histopathology</i> , 2013 , 62, 695-701	7.3	23
187	Biological and clinical implications of nicastrin expression in invasive breast cancer. <i>Breast Cancer Research and Treatment</i> , 2011 , 125, 43-53	4.4	23
186	Prognostic significance of cathepsin V (CTSV/CTSL2) in breast ductal carcinoma in situ. <i>Journal of Clinical Pathology</i> , 2020 , 73, 76-82	3.9	23
185	Bimodality of intratumor Ki67 expression is an independent prognostic factor of overall survival in patients with invasive breast carcinoma. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2016 , 468, 493-502	5.1	22
184	The prognostic significance of STAT3 in invasive breast cancer: analysis of protein and mRNA expressions in large cohorts. <i>Breast Cancer Research and Treatment</i> , 2016 , 156, 9-20	4.4	22
183	A whole slide image-based machine learning approach to predict ductal carcinoma in situ (DCIS) recurrence risk. <i>Breast Cancer Research</i> , 2019 , 21, 83	8.3	22
182	The mammographic correlations of a new immunohistochemical classification of invasive breast cancer. <i>Clinical Radiology</i> , 2008 , 63, 1228-35	2.9	22
181	Cluster-based visualisation with scatter matrices. <i>Pattern Recognition Letters</i> , 2008 , 29, 1814-1823	4.7	22
180	AKT and 14-3-3 regulate Notch4 nuclear localization. <i>Scientific Reports</i> , 2015 , 5, 8782	4.9	21
179	Thioredoxin-interacting protein is an independent risk stratifier for breast ductal carcinoma in situ. <i>Modern Pathology</i> , 2018 , 31, 1807-1815	9.8	21
178	Basal phenotype: a powerful prognostic factor in small screen-detected invasive breast cancer with long-term follow-up. <i>Journal of Medical Screening</i> , 2007 , 14, 210-4	1.4	21
177	Current trials to reduce surgical intervention in ductal carcinoma in situ of the breast: Critical review. <i>Breast</i> , 2017 , 35, 151-156	3.6	21
176	Prolyl-4-hydroxylase β subunit 2 (P4HA2) expression is a predictor of poor outcome in breast ductal carcinoma in situ (DCIS). <i>British Journal of Cancer</i> , 2018 , 119, 1518-1526	8.7	21
175	Markers of progression in early-stage invasive breast cancer: a predictive immunohistochemical panel algorithm for distant recurrence risk stratification. <i>Breast Cancer Research and Treatment</i> , 2015 , 151, 325-33	4.4	20

174	Breast cancer histologic grading using digital microscopy: concordance and outcome association. <i>Journal of Clinical Pathology</i> , 2018 , 71, 680-686	3.9	20
173	Biological characteristics and clinical outcome of triple negative primary breast cancer in older women - comparison with their younger counterparts. <i>PLoS ONE</i> , 2014 , 9, e100573	3.7	20
172	TOMM34 expression in early invasive breast cancer: a biomarker associated with poor outcome. <i>Breast Cancer Research and Treatment</i> , 2012 , 136, 419-27	4.4	20
171	Fatty acid binding protein 7 expression and its sub-cellular localization in breast cancer. <i>Breast Cancer Research and Treatment</i> , 2012 , 134, 519-29	4.4	20
170	EpCAM expression is an indicator of recurrence in basal-like breast cancer. <i>Breast Cancer Research and Treatment</i> , 2012 , 133, 575-82	4.4	20
169	Long-term (37 years) clinical outcome of older women with early operable primary breast cancer managed in a dedicated clinic. <i>Annals of Oncology</i> , 2012 , 23, 1465-71	10.3	20
168	Connexin 43 is an independent predictor of patient outcome in breast cancer patients. <i>Breast Cancer Research and Treatment</i> , 2019 , 174, 93-102	4.4	20
167	Is there a role for base excision repair in estrogen/estrogen receptor-driven breast cancers?. <i>Antioxidants and Redox Signaling</i> , 2014 , 21, 2262-8	8.4	19
166	Overexpression of Specific CD44 Isoforms Is Associated with Aggressive Cell Features in Acquired Endocrine Resistance. <i>Frontiers in Oncology</i> , 2016 , 6, 145	5.3	19
165	Prognostic stratification of oestrogen receptor-positive HER2-negative lymph node-negative class of breast cancer. <i>Histopathology</i> , 2017 , 70, 622-631	7.3	18
164	Phenotypic characterisation of breast cancer: the role of CDC42. <i>Breast Cancer Research and Treatment</i> , 2017 , 164, 317-325	4.4	18
163	Combining clustering and classification ensembles: A novel pipeline to identify breast cancer profiles. <i>Artificial Intelligence in Medicine</i> , 2019 , 97, 27-37	7.4	18
162	A quantifier-based fuzzy classification system for breast cancer patients. <i>Artificial Intelligence in Medicine</i> , 2013 , 58, 175-84	7.4	18
161	Anti-nicestrin monoclonal antibodies elicit pleiotropic anti-tumour pharmacological effects in invasive breast cancer cells. <i>Breast Cancer Research and Treatment</i> , 2014 , 148, 455-62	4.4	18
160	A review of the biological and clinical characteristics of luminal-like oestrogen receptor-positive breast cancer. <i>Histopathology</i> , 2012 , 60, 854-63	7.3	18
159	The prognostic value of the tumor-stroma ratio is most discriminative in patients with grade III or triple-negative breast cancer. <i>International Journal of Cancer</i> , 2020 , 146, 2296-2304	7.5	18
158	Targeting PARP1 in XRCC1-Deficient Sporadic Invasive Breast Cancer or Preinvasive Ductal Carcinoma Induces Synthetic Lethality and Chemoprevention. <i>Cancer Research</i> , 2018 , 78, 6818-6827	10.1	18
157	PKC δ Attenuates Jagged-1-Mediated Notch Signaling in ErbB-2-Positive Breast Cancer to Reverse Trastuzumab Resistance. <i>Clinical Cancer Research</i> , 2016 , 22, 175-86	12.9	17

156	CDC20 expression in oestrogen receptor positive breast cancer predicts poor prognosis and lack of response to endocrine therapy. <i>Breast Cancer Research and Treatment</i> , 2019 , 178, 535-544	4.4	17
155	Prognostic and biological significance of peroxisome proliferator-activated receptor-gamma in luminal breast cancer. <i>Breast Cancer Research and Treatment</i> , 2015 , 150, 511-22	4.4	17
154	The microRNA maturation regulator Drosha is an independent predictor of outcome in breast cancer patients. <i>Breast Cancer Research and Treatment</i> , 2013 , 137, 139-53	4.4	17
153	A comparative study of cytokine gene transcripts in normal and malignant breast tissue and primary cell cultures derived from the same tissue samples. <i>International Journal of Cancer</i> , 1996 , 66, 551-6	7.5	17
152	DNA repair prognostic index modelling reveals an essential role for base excision repair in influencing clinical outcomes in ER negative and triple negative breast cancers. <i>Oncotarget</i> , 2015 , 6, 21964-78	3.2	17
151	Nottingham prognostic index plus (NPI+) predicts risk of distant metastases in primary breast cancer. <i>Breast Cancer Research and Treatment</i> , 2016 , 157, 65-75	4.4	17
150	The molecular mechanisms underlying reduced E-cadherin expression in invasive ductal carcinoma of the breast: high throughput analysis of large cohorts. <i>Modern Pathology</i> , 2019 , 32, 967-976	9.8	17
149	An End-to-End Deep Learning Histochemical Scoring System for Breast Cancer TMA. <i>IEEE Transactions on Medical Imaging</i> , 2019 , 38, 617-628	11.7	17
148	Clinicopathological and prognostic significance of mitogen-activated protein kinases (MAPK) in breast cancers. <i>Breast Cancer Research and Treatment</i> , 2016 , 159, 457-67	4.4	16
147	Overexpression of a novel cell cycle regulator ecdysoneless in breast cancer: a marker of poor prognosis in HER2/neu-overexpressing breast cancer patients. <i>Breast Cancer Research and Treatment</i> , 2012 , 134, 171-80	4.4	16
146	Glutamate dehydrogenase (GLUD1) expression in breast cancer. <i>Breast Cancer Research and Treatment</i> , 2019 , 174, 79-91	4.4	16
145	The clinical and biological significance of HER2 over-expression in breast ductal carcinoma in situ: a large study from a single institution. <i>British Journal of Cancer</i> , 2019 , 120, 1075-1082	8.7	15
144	The prognostic significance of immune microenvironment in breast ductal carcinoma in situ. <i>British Journal of Cancer</i> , 2020 , 122, 1496-1506	8.7	15
143	Impact of breast cancer grade discordance on prediction of outcome. <i>Histopathology</i> , 2018 , 73, 904-915	7.3	15
142	Lack of expression of the proteins GMPR2 and PPAR α are associated with the basal phenotype and patient outcome in breast cancer. <i>Breast Cancer Research and Treatment</i> , 2013 , 137, 127-37	4.4	15
141	The combined expression of solute carriers is associated with a poor prognosis in highly proliferative ER+ breast cancer. <i>Breast Cancer Research and Treatment</i> , 2019 , 175, 27-38	4.4	14
140	Rare case of type I hypersensitivity reaction to sodium hypochlorite solution in a healthcare setting. <i>BMJ Case Reports</i> , 2016 , 2016,	0.9	14
139	Characterisation of HER heterodimers in breast cancer using in situ proximity ligation assay. <i>Breast Cancer Research and Treatment</i> , 2014 , 144, 273-85	4.4	14

138	Low calpain-9 is associated with adverse disease-specific survival following endocrine therapy in breast cancer. <i>BMC Cancer</i> , 2014 , 14, 995	4.8	14
137	Assessing agreement between multiple raters with missing rating information, applied to breast cancer tumour grading. <i>PLoS ONE</i> , 2008 , 3, e2925	3.7	14
136	Prognostic significance of KN motif and ankyrin repeat domains 1 (KANK1) in invasive breast cancer. <i>Breast Cancer Research and Treatment</i> , 2020 , 179, 349-357	4.4	14
135	HAGE in Triple-Negative Breast Cancer Is a Novel Prognostic, Predictive, and Actionable Biomarker: A Transcriptomic and Protein Expression Analysis. <i>Clinical Cancer Research</i> , 2016 , 22, 905-14	12.9	13
134	Collagen (XI) alpha-1 chain is an independent prognostic factor in breast ductal carcinoma in situ. <i>Modern Pathology</i> , 2019 , 32, 1460-1472	9.8	13
133	A novel prognostic two-gene signature for triple negative breast cancer. <i>Modern Pathology</i> , 2020 , 33, 2208-2220	9.8	13
132	The genetic architecture of breast papillary lesions as a predictor of progression to carcinoma. <i>Npj Breast Cancer</i> , 2020 , 6, 9	7.8	13
131	Clinical utility of reverse phase protein array for molecular classification of breast cancer. <i>Breast Cancer Research and Treatment</i> , 2016 , 155, 25-35	4.4	13
130	ERK1/2 is related to oestrogen receptor and predicts outcome in hormone-treated breast cancer. <i>Breast Cancer Research and Treatment</i> , 2014 , 147, 25-37	4.4	13
129	Cytoplasmic localization of alteration/deficiency in activation 3 (ADA3) predicts poor clinical outcome in breast cancer patients. <i>Breast Cancer Research and Treatment</i> , 2013 , 137, 721-31	4.4	13
128	Ki67/SATB1 ratio is an independent prognostic factor of overall survival in patients with early hormone receptor-positive invasive ductal breast carcinoma. <i>Oncotarget</i> , 2015 , 6, 41134-45	3.3	13
127	PIK3C expression by fibroblasts promotes triple-negative breast cancer progression. <i>Journal of Clinical Investigation</i> , 2020 , 130, 3188-3204	15.9	13
126	Enhanced glutamine uptake influences composition of immune cell infiltrates in breast cancer. <i>British Journal of Cancer</i> , 2020 , 122, 94-101	8.7	13
125	Increased expression of glutamine transporter SNAT2/SLC38A2 promotes glutamine dependence and oxidative stress resistance, and is associated with worse prognosis in triple-negative breast cancer. <i>British Journal of Cancer</i> , 2021 , 124, 494-505	8.7	13
124	Invasion in breast lesions: the role of the epithelial-stroma barrier. <i>Histopathology</i> , 2018 , 72, 1075-1083	7.3	13
123	A key genomic subtype associated with lymphovascular invasion in invasive breast cancer. <i>British Journal of Cancer</i> , 2019 , 120, 1129-1136	8.7	12
122	Atypical ductal hyperplasia is a multipotent precursor of breast carcinoma. <i>Journal of Pathology</i> , 2019 , 248, 326-338	9.4	12
121	Co-Expression Effect of SLC7A5/SLC3A2 to Predict Response to Endocrine Therapy in Oestrogen-Receptor-Positive Breast Cancer. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	12

120	Current issues with luminal subtype classification in terms of prediction of benefit from endocrine therapy in early breast cancer. <i>Histopathology</i> , 2018 , 73, 545-558	7.3	12
119	BQ323636.1, a Novel Splice Variant to 2, as a Predictor for Tamoxifen-Resistant Breast Cancer. <i>Clinical Cancer Research</i> , 2018 , 24, 3681-3691	12.9	12
118	Construction of tissue microarrays from core needle biopsies - a systematic literature review. <i>Histopathology</i> , 2016 , 68, 323-32	7.3	12
117	The role of PIP5K1 β /pAKT and targeted inhibition of growth of subtypes of breast cancer using PIP5K1 β inhibitor. <i>Oncogene</i> , 2019 , 38, 375-389	9.2	12
116	Clinicopathological and Functional Significance of RECQL1 Helicase in Sporadic Breast Cancers. <i>Molecular Cancer Therapeutics</i> , 2017 , 16, 239-250	6.1	12
115	Poor-prognosis estrogen receptor-positive breast cancer identified by histopathologic subclassification. <i>Clinical Cancer Research</i> , 2008 , 14, 6625-33	12.9	12
114	Elevated MMP9 expression in breast cancer is a predictor of shorter patient survival. <i>Breast Cancer Research and Treatment</i> , 2020 , 182, 267-282	4.4	12
113	Attention by Selection: A Deep Selective Attention Approach to Breast Cancer Classification. <i>IEEE Transactions on Medical Imaging</i> , 2020 , 39, 1930-1941	11.7	12
112	Kinesin family member-18A (KIF18A) is a predictive biomarker of poor benefit from endocrine therapy in early ER+ breast cancer. <i>Breast Cancer Research and Treatment</i> , 2019 , 173, 93-102	4.4	12
111	The prognostic significance of lysosomal protective protein (cathepsin A) in breast ductal carcinoma in situ. <i>Histopathology</i> , 2019 , 74, 1025-1035	7.3	11
110	KSR1 regulates BRCA1 degradation and inhibits breast cancer growth. <i>Oncogene</i> , 2015 , 34, 2103-14	9.2	11
109	Combined HER3-EGFR score in triple-negative breast cancer provides prognostic and predictive significance superior to individual biomarkers. <i>Scientific Reports</i> , 2020 , 10, 3009	4.9	11
108	Breast conservation in ductal carcinoma in situ (DCIS): what defines optimal margins?. <i>Histopathology</i> , 2017 , 70, 681-692	7.3	11
107	Cell Proliferation (KI-67) Expression Is Associated with Poorer Prognosis in Nigerian Compared to British Breast Cancer Women. <i>ISRN Oncology</i> , 2013 , 2013, 675051		11
106	Influence of E-cadherin expression on the mammographic appearance of invasive nonlobular breast carcinoma detected at screening. <i>Radiology</i> , 2009 , 253, 51-5	20.5	11
105	Targetable ERBB2 mutation status is an independent marker of adverse prognosis in estrogen receptor positive, ERBB2 non-amplified primary lobular breast carcinoma: a retrospective in silico analysis of public datasets. <i>Breast Cancer Research</i> , 2020 , 22, 85	8.3	11
104	The prognostic significance of interferon-stimulated gene 15 (ISG15) in invasive breast cancer. <i>Breast Cancer Research and Treatment</i> , 2021 , 185, 293-305	4.4	11
103	The mammalian target of rapamycin complex 1 (mTORC1) in breast cancer: the impact of oestrogen receptor and HER2 pathways. <i>Breast Cancer Research and Treatment</i> , 2015 , 150, 91-103	4.4	10

102	The prognostic significance of ALDH1A1 expression in early invasive breast cancer. <i>Histopathology</i> , 2020 , 77, 437-448	7.3	10
101	Targeting ataxia telangiectasia-mutated- and Rad3-related kinase (ATR) in PTEN-deficient breast cancers for personalized therapy. <i>Breast Cancer Research and Treatment</i> , 2018 , 169, 277-286	4.4	10
100	Adverse prognostic and predictive significance of low DNA-dependent protein kinase catalytic subunit (DNA-PKcs) expression in early-stage breast cancers. <i>Breast Cancer Research and Treatment</i> , 2014 , 146, 309-20	4.4	10
99	Prognostic and biological significance of proliferation and HER2 expression in the luminal class of breast cancer. <i>Breast Cancer Research and Treatment</i> , 2014 , 145, 317-30	4.4	10
98	Rho-GTPase activating-protein 18: a biomarker associated with good prognosis in invasive breast cancer. <i>British Journal of Cancer</i> , 2017 , 117, 1176-1184	8.7	10
97	Impact of intratumoural heterogeneity on the assessment of Ki67 expression in breast cancer. <i>Breast Cancer Research and Treatment</i> , 2016 , 158, 287-95	4.4	10
96	SPAG5: An Emerging Oncogene. <i>Trends in Cancer</i> , 2020 , 6, 543-547	12.5	9
95	Clinical and biological roles of Kelch-like family member 7 in breast cancer: a marker of poor prognosis. <i>Breast Cancer Research and Treatment</i> , 2018 , 170, 525-533	4.4	9
94	High nuclear MSK1 is associated with longer survival in breast cancer patients. <i>Journal of Cancer Research and Clinical Oncology</i> , 2018 , 144, 509-517	4.9	9
93	Prognostic significance of lymphatic invasion in lymph node-positive breast carcinoma: findings from a large case series with long-term follow-up using immunohistochemical endothelial marker. <i>Modern Pathology</i> , 2014 , 27, 1568-77	9.8	9
92	Further evidence to support bimodality of oestrogen receptor expression in breast cancer. <i>Histopathology</i> , 2017 , 70, 456-465	7.3	9
91	PIAS1 expression in relation to clinicopathological, tumour factors and survival in indigenous black breast cancer women. <i>Journal of Clinical Pathology</i> , 2014 , 67, 301-6	3.9	9
90	Biological profile of oestrogen receptor positive primary breast cancers in the elderly and response to primary endocrine therapy. <i>Critical Reviews in Oncology/Hematology</i> , 2009 , 72, 76-82	7	9
89	The expression of ERalpha, ERbeta and PR in lobular carcinoma in situ of the breast determined using laser microdissection and real-time PCR. <i>Histopathology</i> , 2009 , 54, 419-27	7.3	9
88	FKBP1: a marker of good prognosis in breast cancer. <i>Oncotarget</i> , 2015 , 6, 12209-23	3.3	9
87	Legumain is an independent predictor for invasive recurrence in breast ductal carcinoma in situ. <i>Modern Pathology</i> , 2019 , 32, 639-649	9.8	9
86	Activated zinc transporter ZIP7 as an indicator of anti-hormone resistance in breast cancer. <i>Metallomics</i> , 2019 , 11, 1579-1592	4.5	8
85	Molecular Complexity of Lymphovascular Invasion: The Role of Cell Migration in Breast Cancer as a Prototype. <i>Pathobiology</i> , 2020 , 87, 218-231	3.6	8

84	Checkpoint Kinase 1 Expression Predicts Poor Prognosis in Nigerian Breast Cancer Patients. <i>Molecular Diagnosis and Therapy</i> , 2018 , 22, 79-90	4.5	8
83	The nucleolar-related protein Dyskerin pseudouridine synthase 1 (DKC1) predicts poor prognosis in breast cancer. <i>British Journal of Cancer</i> , 2020 , 123, 1543-1552	8.7	8
82	The prognostic significance of wild-type isocitrate dehydrogenase 2 (IDH2) in breast cancer. <i>Breast Cancer Research and Treatment</i> , 2020 , 179, 79-90	4.4	8
81	Retinoid X receptor gamma (RXRG) is an independent prognostic biomarker in ER-positive invasive breast cancer. <i>British Journal of Cancer</i> , 2019 , 121, 776-785	8.7	7
80	Liver Kinase B1-A Potential Therapeutic Target in Hormone-Sensitive Breast Cancer in Older Women. <i>Cancers</i> , 2019 , 11,	6.6	7
79	Saccharomyces cerevisiae-like 1 (SEC14L1) is a prognostic factor in breast cancer associated with lymphovascular invasion. <i>Modern Pathology</i> , 2018 , 31, 1675-1682	9.8	7
78	Human epidermal growth receptor-2 overexpressing early operable primary breast cancers in older (≥70 years) women: biology and clinical outcome in comparison with younger (. <i>Annals of Oncology</i> , 2014 , 25, 837-842	10.3	7
77	Novel immunohistochemistry-based signatures to predict metastatic site of triple-negative breast cancers. <i>British Journal of Cancer</i> , 2017 , 117, 826-834	8.7	7
76	Clinical importance of estrogen receptor beta isoforms in breast cancer. <i>Journal of Clinical Oncology</i> , 2008 , 26, 5825; author reply 5825-6	2.2	7
75	Prognostic significance of nucleolar assessment in invasive breast cancer. <i>Histopathology</i> , 2020 , 76, 671-684	9.4	7
74	Clinicopathological significance of ataxia telangiectasia-mutated (ATM) kinase and ataxia telangiectasia-mutated and Rad3-related (ATR) kinase in MYC overexpressed breast cancers. <i>Breast Cancer Research and Treatment</i> , 2019 , 175, 105-115	4.4	7
73	Geometric characteristics of collagen have independent prognostic significance in breast ductal carcinoma in situ: an image analysis study. <i>Modern Pathology</i> , 2019 , 32, 1473-1485	9.8	6
72	Myxovirus resistance 1 (MX1) is an independent predictor of poor outcome in invasive breast cancer. <i>Breast Cancer Research and Treatment</i> , 2020 , 181, 541-551	4.4	6
71	Mediator complex (MED) 7: a biomarker associated with good prognosis in invasive breast cancer, especially ER+ luminal subtypes. <i>British Journal of Cancer</i> , 2018 , 118, 1142-1151	8.7	6
70	ERCC1 Is a Predictor of Anthracycline Resistance and Taxane Sensitivity in Early Stage or Locally Advanced Breast Cancers. <i>Cancers</i> , 2019 , 11,	6.6	6
69	Interleukin-3: a putative protective factor against breast cancer which is secreted by male but not female breast fibroblasts. <i>International Journal of Cancer</i> , 1995 , 61, 416-9	7.5	6
68	Intratumoural Cytochrome P450 Expression in Breast Cancer: Impact on Standard of Care Treatment and New Efforts to Develop Tumour-Selective Therapies. <i>Biomedicines</i> , 2021 , 9,	4.8	6
67	Nucleolar protein 10 (NOP10) predicts poor prognosis in invasive breast cancer. <i>Breast Cancer Research and Treatment</i> , 2021 , 185, 615-627	4.4	6

66	DNA methylation landscapes of 1538 breast cancers reveal a replication-linked clock, epigenomic instability and cis-regulation. <i>Nature Communications</i> , 2021 , 12, 5406	17.4	6
65	Chemokine (C-C motif) receptor 7 (CCR7) associates with the tumour immune microenvironment but not progression in invasive breast carcinoma. <i>Journal of Pathology: Clinical Research</i> , 2017 , 3, 105-114	5.3	5
64	PPFIA1 expression associates with poor response to endocrine treatment in luminal breast cancer. <i>BMC Cancer</i> , 2020 , 20, 425	4.8	5
63	The prognostic significance of BMI1 expression in invasive breast cancer is dependent on its molecular subtypes. <i>Breast Cancer Research and Treatment</i> , 2020 , 182, 581-589	4.4	5
62	SHON is a novel estrogen-regulated oncogene in mammary carcinoma that predicts patient response to endocrine therapy. <i>Cancer Research</i> , 2013 , 73, 6951-62	10.1	5
61	Clinicopathological significance of lipocalin 2 nuclear expression in invasive breast cancer. <i>Breast Cancer Research and Treatment</i> , 2020 , 179, 557-564	4.4	5
60	Dopamine and cAMP-regulated phosphoprotein 32 kDa (DARPP-32) and survival in breast cancer: a retrospective analysis of protein and mRNA expression. <i>Scientific Reports</i> , 2019 , 9, 16987	4.9	5
59	Molecular profiling of breast cancer in Nigerian women identifies an altered p53 pathway as a major mechanism underlying its poor prognosis compared with British counterpart. <i>Malaysian Journal of Pathology</i> , 2014 , 36, 3-17	1.7	5
58	Utility of ankyrin 3 as a prognostic marker in androgen-receptor-positive breast cancer. <i>Breast Cancer Research and Treatment</i> , 2019 , 176, 63-73	4.4	4
57	The TRAIL system is over-expressed in breast cancer and FLIP a marker of good prognosis. <i>Journal of Cancer Research and Clinical Oncology</i> , 2015 , 141, 505-14	4.9	4
56	Retrospective assessment of cyclin-dependent kinase 5 mRNA and protein expression and its association with patient survival in breast cancer. <i>Journal of Cellular and Molecular Medicine</i> , 2020 , 24, 6263-6271	5.6	4
55	Visual histological assessment of morphological features reflects the underlying molecular profile in invasive breast cancer: a morphomolecular study. <i>Histopathology</i> , 2020 , 77, 631-645	7.3	4
54	A Quantitative Centrosomal Amplification Score Predicts Local Recurrence of Ductal Carcinoma. <i>Clinical Cancer Research</i> , 2020 , 26, 2898-2907	12.9	4
53	Clinicopathological and prognostic significance of Ras association and pleckstrin homology domains 1 (RAPH1) in breast cancer. <i>Breast Cancer Research and Treatment</i> , 2018 , 172, 61-68	4.4	4
52	Expression of angiogenic chemokines in ovarian clear cell carcinoma. <i>Journal of Obstetrics and Gynaecology Research</i> , 2013 , 39, 297-304	1.9	4
51	The intra-tumoural stroma in patients with breast cancer increases with age. <i>Breast Cancer Research and Treatment</i> , 2020 , 179, 37-45	4.4	4
50	Clinicopathological and molecular characteristics of Ku 70/80 expression in Nigerian breast cancer and its potential therapeutic implications. <i>Pathology Research and Practice</i> , 2017 , 213, 27-33	3.4	3
49	Integrated Analysis of Key Differentially Expressed Genes Identifies DBN1 as a Predictive Marker of Response to Endocrine Therapy in Luminal Breast Cancer. <i>Cancers</i> , 2020 , 12,	6.6	3

48	The solute carrier SLC7A8 is a marker of favourable prognosis in ER-positive low proliferative invasive breast cancer. <i>Breast Cancer Research and Treatment</i> , 2020 , 181, 1-12	4.4	3
47	Surgical management of ductal carcinoma in situ of the breast: A large retrospective study from a single institution. <i>Breast Journal</i> , 2019 , 25, 1143-1153	1.2	3
46	Detection and quantification of microRNAs in laser-microdissected formalin-fixed paraffin-embedded breast cancer tissues. <i>Methods in Molecular Biology</i> , 2011 , 755, 119-42	1.4	3
45	Abstract B046: Therapeutic targeting of integrin $\alpha 5 \beta 1$ in high-risk breast cancer 2013 ,		3
44	XRCC1 deficient triple negative breast cancers are sensitive to ATR, ATM and Wee1 inhibitor either alone or in combination with olaparib. <i>Therapeutic Advances in Medical Oncology</i> , 2020 , 12, 17588359209742013	5.4	3
43	The ITIM-Containing Receptor: Leukocyte-Associated Immunoglobulin-Like Receptor-1 (LAIR-1) Modulates Immune Response and Confers Poor Prognosis in Invasive Breast Carcinoma. <i>Cancers</i> , 2020 , 13,	6.6	3
42	The clinical significance of oestrogen receptor expression in breast ductal carcinoma in situ. <i>British Journal of Cancer</i> , 2020 , 123, 1513-1520	8.7	3
41	RANK signaling increases after anti-HER2 therapy contributing to the emergence of resistance in HER2-positive breast cancer. <i>Breast Cancer Research</i> , 2021 , 23, 42	8.3	3
40	The role of ALDH1A1 in contributing to breast tumour aggressiveness: A study conducted in an African population. <i>Annals of Diagnostic Pathology</i> , 2021 , 51, 151696	2.2	3
39	Werner Syndrome Protein Expression in Breast Cancer. <i>Clinical Breast Cancer</i> , 2021 , 21, 57-73.e7	3	3
38	Co-expression of nuclear P38 and hormone receptors is prognostic of good long-term clinical outcome in primary breast cancer and is linked to upregulation of DNA repair. <i>BMC Cancer</i> , 2018 , 18, 1027	4.8	3
37	Treatment strategies and survival outcomes in older women with breast cancer: A comparative study between the FOCUS cohort and Nottingham cohort. <i>Journal of Geriatric Oncology</i> , 2018 , 9, 635-641	3.6	3
36	The Biological and Clinical Significance of Glutaminase in Luminal Breast Cancer. <i>Cancers</i> , 2021 , 13,	6.6	3
35	Cytoplasmic Cyclin E Is an Independent Marker of Aggressive Tumor Biology and Breast Cancer-Specific Mortality in Women over 70 Years of Age. <i>Cancers</i> , 2020 , 12,	6.6	2
34	Machine learning-based prediction of breast cancer growth rate in vivo. <i>British Journal of Cancer</i> , 2019 , 121, 497-504	8.7	2
33	Comparison of the PharmDx immunohistochemical system with standard methods for assessing estrogen and progesterone receptors in invasive carcinoma of the breast. <i>Applied Immunohistochemistry and Molecular Morphology</i> , 2013 , 21, 90-3	1.9	2
32	Biology of Oestrogen-Receptor Positive Primary Breast Cancer in Older Women with Utilisation of Core Needle Biopsy Samples and Correlation with Clinical Outcome. <i>Cancers</i> , 2020 , 12,	6.6	2
31	Association of Sperm-Associated Antigen 5 and Treatment Response in Patients With Estrogen Receptor-Positive Breast Cancer. <i>JAMA Network Open</i> , 2020 , 3, e209486	10.4	2

30	SHON expression predicts response and relapse risk of breast cancer patients after anthracycline-based combination chemotherapy or tamoxifen treatment. <i>British Journal of Cancer</i> , 2019 , 120, 728-745	8.7	2
29	A review of the racial heterogeneity of breast cancer stem cells. <i>Gene</i> , 2021 , 796-797, 145805	3.8	2
28	Standardization of the tumor-stroma ratio scoring method for breast cancer research.. <i>Breast Cancer Research and Treatment</i> , 2022 , 193, 545	4.4	2
27	Ubiquitin-conjugating enzyme 2C (UBE2C) is a poor prognostic biomarker in invasive breast cancer.. <i>Breast Cancer Research and Treatment</i> , 2022 , 192, 529	4.4	1
26	Oestrogen receptor negative early operable primary breast cancer in older women-Biological characteristics and long-term clinical outcome. <i>PLoS ONE</i> , 2017 , 12, e0188528	3.7	1
25	Untangling the clinicopathological significance of MRE11-RAD50-NBS1 complex in sporadic breast cancers. <i>Npj Breast Cancer</i> , 2021 , 7, 143	7.8	1
24	HAGE (DDX43) protein expression as an independent biomarker of poor clinical outcome of breast cancer (BC) and potential as a therapeutic target for ER-negative BC.. <i>Journal of Clinical Oncology</i> , 2012 , 30, 1013-1013	2.2	1
23	Breast cancer stem cells: A fallow research ground in Africa. <i>Pathology Research and Practice</i> , 2020 , 216, 153118	3.4	1
22	Genome-Wide Analysis Unveils DNA Helicase RECQ1 as a Regulator of Estrogen Response Pathway in Breast Cancer Cells. <i>Molecular and Cellular Biology</i> , 2021 , 41,	4.8	1
21	PP1, PKA and DARPP-32 in breast cancer: A retrospective assessment of protein and mRNA expression. <i>Journal of Cellular and Molecular Medicine</i> , 2021 , 25, 5015-5024	5.6	1
20	SLC1A5 co-expression with TALDO1 associates with endocrine therapy failure in estrogen receptor-positive breast cancer. <i>Breast Cancer Research and Treatment</i> , 2021 , 189, 317-331	4.4	1
19	Oestrogen-regulated protein SLC39A6: a biomarker of good prognosis in luminal breast cancer. <i>Breast Cancer Research and Treatment</i> , 2021 , 189, 621-630	4.4	1
18	Epigenome erosion and SOX10 drive neural crest phenotypic mimicry in triple-negative breast cancer.. <i>Npj Breast Cancer</i> , 2022 , 8, 57	7.8	1
17	Aurora Kinase A Is an Independent Predictor of Invasive Recurrence in Breast Ductal Carcinoma in situ.. <i>Pathobiology</i> , 2022 , 1-11	3.6	1
16	Differing pattern of biological characteristics of early operable primary breast cancer according to age.. <i>Journal of Clinical Oncology</i> , 2016 , 34, e23272-e23272	2.2	0
15	Prognostic significance of receptor expression discordance between primary and recurrent breast cancers: a meta-analysis. <i>Breast Cancer Research and Treatment</i> , 2021 , 1	4.4	0
14	A multi-institutional study of racial differences in androgen receptor status among triple-negative breast cancers.. <i>Journal of Clinical Oncology</i> , 2016 , 34, 1089-1089	2.2	0
13	The prognostic significance of Flap Endonuclease 1 (FEN1) in breast ductal carcinoma in situ. <i>Breast Cancer Research and Treatment</i> , 2021 , 188, 53-63	4.4	0

- 12 The combination phenotype of B-cell specific Moloney murine leukaemia virus integration site 1 (BMI1) and CD44+/CD24^{low} associates with poor clinicopathological features in African patients with breast cancer. *Gene Reports*, **2022**, 26, 101475 1.4
- 11 Predicting chemotherapy response in invasive breast cancer.. *Journal of Clinical Oncology*, **2014**, 32, 1084-1084 1.4
- 10 Intracellular intermolecular relationships of LKB1 in breast cancer.. *Journal of Clinical Oncology*, **2014**, 32, 554-554 2.2
- 9 Mechanistic and clinical analysis of Sperm associated antigen 5 (SPAG5) as a novel prognostic, predictive, actionable gene in Breast Cancer (BC).. *Journal of Clinical Oncology*, **2015**, 33, 1040-1040 2.2
- 8 Potential of a novel technique for constructing tissue microarrays from core needle biopsy as seen in older women with primary breast cancer.. *Journal of Clinical Oncology*, **2016**, 34, e23283-e23283 2.2
- 7 Multi-institutional study of triple negative breast cancer stratification by a metric that quantifies cell cycling kinetics.. *Journal of Clinical Oncology*, **2016**, 34, 1091-1091 2.2
- 6 A combined HER3-EGFR score in triple-negative breast cancer: racial differences.. *Journal of Clinical Oncology*, **2016**, 34, e12560-e12560 2.2
- 5 A study of sperm-associated antigen 5 (SPAG5) in predicting response to anthracycline (ATC)/platinum chemotherapies (CT) in breast (BC) and ovarian cancers (OVC).. *Journal of Clinical Oncology*, **2012**, 30, 1098-1098 2.2
- 4 Targeting XRCC1 (X-ray repair cross-complementing gene 1) deficiency in tumors for personalized cancer therapy.. *Journal of Clinical Oncology*, **2012**, 30, 1014-1014 2.2
- 3 Human epidermal growth receptor-2 overexpressing early operable primary breast cancers in older (≥70 years) women: Biology and clinical outcome in comparison with younger (. *Journal of Clinical Oncology*, **2013**, 31, e11608-e11608 2.2
- 2 The frequency and clinical significance of DNA polymerase beta (POLβ) expression in breast ductal carcinoma in situ (DCIS). *Breast Cancer Research and Treatment*, **2021**, 190, 39-51 4.4
- 1 Upregulation of Cyclin B2 (C) in breast cancer contributes to the development of lymphovascular invasion.. *American Journal of Cancer Research*, **2022**, 12, 469-489 4.4