

Alexandre S C Reis

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2629618/publications.pdf>

Version: 2024-02-01

12
papers

1,009
citations

1306789

7
h-index

1473754

9
g-index

12
all docs

12
docs citations

12
times ranked

759
citing authors

#	ARTICLE	IF	CITATIONS
1	Enhanced Detection of Open-angle Glaucoma with an Anatomically Accurate Optical Coherence Tomography-Derived Neuroretinal Rim Parameter. <i>Ophthalmology</i> , 2013, 120, 535-543.	2.5	323
2	Optic Disc Margin Anatomy in Patients with Glaucoma and Normal Controls with Spectral Domain Optical Coherence Tomography. <i>Ophthalmology</i> , 2012, 119, 738-747.	2.5	239
3	Influence of Clinically Invisible, but Optical Coherence Tomography Detected, Optic Disc Margin Anatomy on Neuroretinal Rim Evaluation. , 2012, 53, 1852.		231
4	Laminar Displacement and Prelaminar Tissue Thickness Change after Glaucoma Surgery Imaged with Optical Coherence Tomography. , 2012, 53, 5819.		100
5	Automated Segmentation of Optic Nerve Head Structures With Optical Coherence Tomography. , 2014, 55, 1161.		32
6	Intra- and interobserver reproducibility of Bruch's membrane opening minimum rim width measurements with spectral domain optical coherence tomography. <i>Acta Ophthalmologica</i> , 2017, 95, e548-e555.	0.6	28
7	Rates of Change in the Visual Field and Optic Disc in Patients with Distinct Patterns of Glaucomatous Optic Disc Damage. <i>Ophthalmology</i> , 2012, 119, 294-303.	2.5	19
8	Bruch's membrane opening minimum rim width and retinal nerve fiber layer thickness in a Brazilian population of healthy subjects. <i>PLoS ONE</i> , 2018, 13, e0206887.	1.1	12
9	Evaluation of contrast sensitivity in patients with advanced glaucoma: comparison of two tests. <i>British Journal of Ophthalmology</i> , 2020, 104, 1418-1422.	2.1	9
10	Interocular Asymmetry of Minimum Rim Width and Retinal Nerve Fiber Layer Thickness in Healthy Brazilian Individuals. <i>Journal of Glaucoma</i> , 2018, 27, 1136-1141.	0.8	7
11	Influence of Bruch's Membrane Opening Area in Diagnosing Glaucoma With Neuroretinal Parameters From Optical Coherence Tomography. <i>American Journal of Ophthalmology</i> , 2019, 208, 94-102.	1.7	5
12	Qualitative evaluation of neuroretinal rim and retinal nerve fibre layer on optical coherence tomography to detect glaucomatous damage. <i>British Journal of Ophthalmology</i> , 2020, 104, 980-984.	2.1	4