

Ronald P Stolk

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/2642623/ronald-p-stolk-publications-by-year.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

236
papers

20,242
citations

68
h-index

137
g-index

243
ext. papers

23,654
ext. citations

7.2
avg, IF

6.07
L-index

#	Paper	IF	Citations
236	Effect of metabolic genetic variants on long-term disease comorbidity in patients with type 2 diabetes. <i>Scientific Reports</i> , 2021 , 11, 2794	4.9	
235	Later achievement of infant motor milestones is related to lower levels of physical activity during childhood: the GECKO Drenthe cohort. <i>BMC Pediatrics</i> , 2019 , 19, 388	2.6	0
234	Combining informal care and paid work: The use of work arrangements by working adult-child caregivers in the Netherlands. <i>Health and Social Care in the Community</i> , 2018 , 26, e122-e131	2.6	13
233	Long-term Air Pollution Exposure, Genome-wide DNA Methylation and Lung Function in the LifeLines Cohort Study. <i>Environmental Health Perspectives</i> , 2018 , 126, 027004	8.4	50
232	Ambient air pollution, traffic noise and adult asthma prevalence: a BioSHaRE approach. <i>European Respiratory Journal</i> , 2017 , 49,	13.6	44
231	Missing heritability: is the gap closing? An analysis of 32 complex traits in the Lifelines Cohort Study. <i>European Journal of Human Genetics</i> , 2017 , 25, 877-885	5.3	43
230	Long-term exposure to road traffic noise, ambient air pollution, and cardiovascular risk factors in the HUNT and lifelines cohorts. <i>European Heart Journal</i> , 2017 , 38, 2290-2296	9.5	80
229	Plasma N-terminal Prohomocysteinemia and Risk of Incident Cardiovascular Disease and All-Cause Mortality in a Prospective Observational Cohort: the PREVEND Study. <i>Clinical Chemistry</i> , 2017 , 63, 278-287	5.5	3
228	Changing the obesogenic environment to improve cardiometabolic health in residential patients with a severe mental illness: cluster randomised controlled trial. <i>British Journal of Psychiatry</i> , 2017 , 211, 296-303	5.4	13
227	Maelstrom Research guidelines for rigorous retrospective data harmonization. <i>International Journal of Epidemiology</i> , 2017 , 46, 103-105	7.8	62
226	Factors of physical activity among Chinese children and adolescents: a systematic review. <i>International Journal of Behavioral Nutrition and Physical Activity</i> , 2017 , 14, 36	8.4	59
225	The relationship between social functioning and subjective memory complaints in older persons: a population-based longitudinal cohort study. <i>International Journal of Geriatric Psychiatry</i> , 2017 , 32, 1059-1071	3.9	20
224	The LifeLines Cohort Study: Prevalence and treatment of cardiovascular disease and risk factors. <i>International Journal of Cardiology</i> , 2017 , 228, 495-500	3.2	48
223	A principal component meta-analysis on multiple anthropometric traits identifies novel loci for body shape. <i>Nature Communications</i> , 2016 , 7, 13357	17.4	46
222	52 Genetic Loci Influencing Myocardial Mass. <i>Journal of the American College of Cardiology</i> , 2016 , 68, 1435-1448	15.1	76
221	Neighborhood income and major depressive disorder in a large Dutch population: results from the LifeLines Cohort study. <i>BMC Public Health</i> , 2016 , 16, 773	4.1	16
220	Dutch healthcare professionals inadequately perceived if three- and four-year-old preschool children were overweight. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2016 , 105, 1198-203	3.1	1

219	Social relationships and cognitive decline: a systematic review and meta-analysis of longitudinal cohort studies. <i>International Journal of Epidemiology</i> , 2016 , 45, 1169-1206	7.8	95
218	Effect of diagnosis with a chronic disease on physical activity behavior in middle-aged women. <i>Preventive Medicine</i> , 2016 , 83, 56-62	4.3	14
217	The association of air pollution and depressed mood in 70,928 individuals from four European cohorts. <i>International Journal of Hygiene and Environmental Health</i> , 2016 , 219, 212-9	6.9	101
216	Eosinophil Count Is a Common Factor for Complex Metabolic and Pulmonary Traits and Diseases: The LifeLines Cohort Study. <i>PLoS ONE</i> , 2016 , 11, e0168480	3.7	18
215	Survey nonresponse among informal caregivers: effects on the presence and magnitude of associations with caregiver burden and satisfaction. <i>BMC Public Health</i> , 2016 , 16, 480	4.1	17
214	Association of Depressive and Anxiety Disorders With Diagnosed Versus Undiagnosed Diabetes: An Epidemiological Study of 90,686 Participants. <i>Psychosomatic Medicine</i> , 2016 , 78, 233-41	3.7	37
213	The LifeLines Cohort Study: a resource providing new opportunities for environmental epidemiology. <i>Archives of Public Health</i> , 2016 , 74, 32	2.6	8
212	Glucocorticoid receptor haplotype and metabolic syndrome: the Lifelines cohort study. <i>European Journal of Endocrinology</i> , 2016 , 175, 645-651	6.5	16
211	Air pollution exposure is associated with restrictive ventilatory patterns. <i>European Respiratory Journal</i> , 2016 , 48, 1221-1224	13.6	14
210	Social relationships and risk of dementia: A systematic review and meta-analysis of longitudinal cohort studies. <i>Ageing Research Reviews</i> , 2015 , 22, 39-57	12	432
209	Low levels of vitamin D are associated with multimorbidity: results from the LifeLines Cohort Study. <i>Annals of Medicine</i> , 2015 , 47, 474-81	1.5	24
208	An activity stimulation programme during a child's first year reduces some indicators of adiposity at the age of two-and-a-half. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2015 , 104, 414-21	3.1	18
207	Bilirubin as a potential causal factor in type 2 diabetes risk: a Mendelian randomization study. <i>Diabetes</i> , 2015 , 64, 1459-69	0.9	68
206	Genome-wide meta-analysis identifies six novel loci associated with habitual coffee consumption. <i>Molecular Psychiatry</i> , 2015 , 20, 647-656	15.1	167
205	Cohort profile: LifeLines DEEP, a prospective, general population cohort study in the northern Netherlands: study design and baseline characteristics. <i>BMJ Open</i> , 2015 , 5, e006772	3	136
204	Change in HbA1c levels between the age of 8 years and the age of 12 years in Dutch children without diabetes: the PIAMA birth cohort study. <i>PLoS ONE</i> , 2015 , 10, e0119615	3.7	4
203	The Influence of Age and Sex on Genetic Associations with Adult Body Size and Shape: A Large-Scale Genome-Wide Interaction Study. <i>PLoS Genetics</i> , 2015 , 11, e1005378	6	220
202	Representativeness of the LifeLines Cohort Study. <i>PLoS ONE</i> , 2015 , 10, e0137203	3.7	145

201	Noise and somatic symptoms: A role for personality traits?. <i>International Journal of Hygiene and Environmental Health</i> , 2015 , 218, 543-9	6.9	11
200	New genetic loci link adipose and insulin biology to body fat distribution. <i>Nature</i> , 2015 , 518, 187-196	50.4	920
199	Genetic studies of body mass index yield new insights for obesity biology. <i>Nature</i> , 2015 , 518, 197-206	50.4	2687
198	Cohort Profile: LifeLines, a three-generation cohort study and biobank. <i>International Journal of Epidemiology</i> , 2015 , 44, 1172-80	7.8	364
197	Determinants of Weight Gain during the First Two Years of Life--The GECKO Drenthe Birth Cohort. <i>PLoS ONE</i> , 2015 , 10, e0133326	3.7	19
196	(Un)Healthy in the City: Respiratory, Cardiometabolic and Mental Health Associated with Urbanity. <i>PLoS ONE</i> , 2015 , 10, e0143910	3.7	21
195	Associations of life events during pregnancy with longitudinal change in symptoms of antenatal anxiety and depression. <i>Midwifery</i> , 2014 , 30, 526-31	2.8	30
194	The prevalence of metabolic syndrome and metabolically healthy obesity in Europe: a collaborative analysis of ten large cohort studies. <i>BMC Endocrine Disorders</i> , 2014 , 14, 9	3.3	311
193	Gene-age interactions in blood pressure regulation: a large-scale investigation with the CHARGE, Global BPgen, and ICBP Consortia. <i>American Journal of Human Genetics</i> , 2014 , 95, 24-38	11	80
192	Defining the role of common variation in the genomic and biological architecture of adult human height. <i>Nature Genetics</i> , 2014 , 46, 1173-86	36.3	1339
191	Circulating peroxiredoxin 4 and type 2 diabetes risk: the Prevention of Renal and Vascular Endstage Disease (PREVEND) study. <i>Diabetologia</i> , 2014 , 57, 1842-9	10.3	15
190	Association of vitamin D status with arterial blood pressure and hypertension risk: a mendelian randomisation study. <i>Lancet Diabetes and Endocrinology</i> , 2014 , 2, 719-29	18.1	250
189	Identification of relevant biomarkers for type 2 diabetes. <i>Lancet Diabetes and Endocrinology</i> , 2014 , 2, 106-7	18.1	5
188	Seasonal influenza vaccine effectiveness against influenza in 2012-2013: a hospital-based case-control study in Lithuania. <i>Vaccine</i> , 2014 , 32, 857-63	4.1	12
187	Causal inference algorithms can be useful in life course epidemiology. <i>Journal of Clinical Epidemiology</i> , 2014 , 67, 190-8	5.7	10
186	Comment on Vassy et al. polygenic type 2 diabetes prediction at the limit of common variant detection. <i>Diabetes</i> 2014;63:2172-2182. <i>Diabetes</i> , 2014 , 63, e11-2	0.9	1
185	A multidisciplinary intervention programme has positive effects on quality of life in overweight and obese preschool children. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2014 , 103, 962-7	3.1	6
184	Predictive accuracy of Edinburgh postnatal depression scale assessment during pregnancy for the risk of developing postpartum depressive symptoms: a prospective cohort study. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2014 , 121, 1604-10	3.7	35

183	DataSHIELD: taking the analysis to the data, not the data to the analysis. <i>International Journal of Epidemiology</i> , 2014 , 43, 1929-44	7.8	116
182	Gene-centric meta-analysis in 87,736 individuals of European ancestry identifies multiple blood-pressure-related loci. <i>American Journal of Human Genetics</i> , 2014 , 94, 349-60	11	131
181	Three-year follow-up of 3-year-old to 5-year-old children after participation in a multidisciplinary or a usual-care obesity treatment program. <i>Clinical Nutrition</i> , 2014 , 33, 1095-100	5.9	14
180	Pleiotropic genes for metabolic syndrome and inflammation. <i>Molecular Genetics and Metabolism</i> , 2014 , 112, 317-38	3.7	81
179	Effect of obesity intervention programs on adipokines, insulin resistance, lipid profile, and low-grade inflammation in 3- to 5-y-old children. <i>Pediatric Research</i> , 2014 , 75, 352-7	3.2	12
178	The role of fitness in the association between fatness and cardiometabolic risk from childhood to adolescence. <i>Pediatric Diabetes</i> , 2013 , 14, 57-65	3.6	32
177	Growth during infancy and childhood, and adiposity at age 16 years: ages 2 to 7 years are pivotal. <i>Journal of Pediatrics</i> , 2013 , 162, 287-92.e2	3.6	22
176	Role of HDL cholesterol and estimates of HDL particle composition in future development of type 2 diabetes in the general population: the PREVEND study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013 , 98, E1352-9	5.6	72
175	Loci influencing blood pressure identified using a cardiovascular gene-centric array. <i>Human Molecular Genetics</i> , 2013 , 22, 1663-78	5.6	119
174	Infant movement opportunities are related to early growth--GECKO Drenthe cohort. <i>Early Human Development</i> , 2013 , 89, 457-61	2.2	13
173	Insulin resistance and cardiovascular risk factors in 3- to 5-year-old overweight or obese children. <i>Hormone Research in Paediatrics</i> , 2013 , 80, 201-6	3.3	29
172	Data harmonization and federated analysis of population-based studies: the BioSHaRE project. <i>Emerging Themes in Epidemiology</i> , 2013 , 10, 12	3.9	80
171	Contamination rates of three urine-sampling methods to assess bacteriuria in pregnant women. <i>Obstetrics and Gynecology</i> , 2013 , 121, 299-305	4.9	18
170	Ultrasound estimates of visceral and subcutaneous-abdominal adipose tissues in infancy. <i>Journal of Obesity</i> , 2013 , 2013, 951954	3.7	22
169	Parental history of type 2 diabetes and cardiometabolic biomarkers in offspring. <i>European Journal of Clinical Investigation</i> , 2012 , 42, 974-82	4.6	6
168	External validation of the KORA S4/F4 prediction models for the risk of developing type 2 diabetes in older adults: the PREVEND study. <i>European Journal of Epidemiology</i> , 2012 , 27, 47-52	12.1	14
167	Motorized transportation, social status, and adiposity: the China Health and Nutrition Survey. <i>American Journal of Preventive Medicine</i> , 2012 , 43, 1-10	6.1	61
166	Meta-analyses identify 13 loci associated with age at menopause and highlight DNA repair and immune pathways. <i>Nature Genetics</i> , 2012 , 44, 260-8	36.3	243

165	Large-scale gene-centric meta-analysis across 32 studies identifies multiple lipid loci. <i>American Journal of Human Genetics</i> , 2012 , 91, 823-38	11	189
164	Comparison of frequency of periprocedural myocardial infarction in patients with and without diabetes mellitus to those with previously unknown but elevated glycated hemoglobin levels (from the TWENTE Trial). <i>American Journal of Cardiology</i> , 2012 , 110, 1561-7	3	20
163	Differential effects of comorbidity on antihypertensive and glucose-regulating treatment in diabetes mellitus--a cohort study. <i>PLoS ONE</i> , 2012 , 7, e38707	3.7	23
162	Liver function tests and risk prediction of incident type 2 diabetes: evaluation in two independent cohorts. <i>PLoS ONE</i> , 2012 , 7, e51496	3.7	25
161	Sex differences in the association between plasma copeptin and incident type 2 diabetes: the Prevention of Renal and Vascular Endstage Disease (PREVEND) study. <i>Diabetologia</i> , 2012 , 55, 1963-70	10.3	62
160	Peroxiredoxin 4, a novel circulating biomarker for oxidative stress and the risk of incident cardiovascular disease and all-cause mortality. <i>Journal of the American Heart Association</i> , 2012 , 1, e002956	6	33
159	Results of a multidisciplinary treatment program in 3-year-old to 5-year-old overweight or obese children: a randomized controlled clinical trial. <i>JAMA Pediatrics</i> , 2012 , 166, 1109-15		52
158	Low HDL cholesterol and the risk of diabetic nephropathy and retinopathy: results of the ADVANCE study. <i>Diabetes Care</i> , 2012 , 35, 2201-6	14.6	76
157	PS4 - 22. HDL-cholesterol, Apolipoprotein A-I/A-II, and HDL-cholesterol particle composition for the risk of developing type 2 diabetes in the community: the PREVEND Study. <i>Nederlands Tijdschrift Voor Diabetologie</i> , 2012 , 10, 113-113	0	
156	PS4 - 23. Bilirubin and risk of type 2 diabetes: a mendelian randomization approach. <i>Nederlands Tijdschrift Voor Diabetologie</i> , 2012 , 10, 113-114	0	
155	Prediction models for risk of developing type 2 diabetes: systematic literature search and independent external validation study. <i>BMJ, The</i> , 2012 , 345, e5900	5.9	195
154	Is directly measured physical activity related to adiposity in preschool children?. <i>Pediatric Obesity</i> , 2011 , 6, 389-400		19
153	Genome-wide association study identifies loci influencing concentrations of liver enzymes in plasma. <i>Nature Genetics</i> , 2011 , 43, 1131-8	36.3	415
152	Validation of ultrasound estimates of visceral fat in black South African adolescents. <i>Obesity</i> , 2011 , 19, 1892-7	8	28
151	Vitamin D deficiency: universal risk factor for multifactorial diseases?. <i>Current Drug Targets</i> , 2011 , 12, 97-106	3	24
150	Maternal and paternal transmission of type 2 diabetes: influence of diet, lifestyle and adiposity. <i>Journal of Internal Medicine</i> , 2011 , 270, 388-96	10.8	28
149	Medication adherence affects treatment modifications in patients with type 2 diabetes. <i>Clinical Therapeutics</i> , 2011 , 33, 121-34	3.5	24
148	Plasma procalcitonin and risk of type 2 diabetes in the general population. <i>Diabetologia</i> , 2011 , 54, 2463-50.3	50.3	22

147	PRegnancy Outcomes after a Maternity Intervention for Stressful EmotionS (PROMISES): study protocol for a randomised controlled trial. <i>Trials</i> , 2011 , 12, 157	2.8	15
146	Is rigorous retrospective harmonization possible? Application of the DataSHaPER approach across 53 large studies. <i>International Journal of Epidemiology</i> , 2011 , 40, 1314-28	7.8	65
145	PS9 - 48. External validation of the KORA S4/F4 prediction models for the risk of developing type 2 diabetes in older adults: the PREVEND Study. <i>Nederlands Tijdschrift Voor Diabetologie</i> , 2011 , 9, 124-124 ^o		
144	Low physical activity and risk of cardiovascular and all-cause mortality in renal transplant recipients. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2011 , 6, 898-905	6.9	94
143	HbA1c in nondiabetic Dutch infants aged 8-12 months: the GECKO-Drenthe birth cohort study. <i>Diabetes Care</i> , 2011 , 34, 403-5	14.6	6
142	Parental factors affecting the weights of the placenta and the offspring. <i>Journal of Perinatal Medicine</i> , 2011 , 39, 27-34	2.7	27
141	Development and validation of a general population renal risk score. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2011 , 6, 1731-8	6.9	56
140	Genome-wide association study identifies six new loci influencing pulse pressure and mean arterial pressure. <i>Nature Genetics</i> , 2011 , 43, 1005-11	36.3	338
139	Ultrasound measurements of visceral and subcutaneous abdominal thickness to predict abdominal adiposity among older men and women. <i>Obesity</i> , 2010 , 18, 625-31	8	85
138	Association between alcohol consumption and diabetic retinopathy and visual acuity-the AdRem Study. <i>Diabetic Medicine</i> , 2010 , 27, 1130-7	3.5	26
137	Physical activity, adiposity, and diabetes risk in middle-aged and older Chinese population: the Guangzhou Biobank Cohort Study. <i>Diabetes Care</i> , 2010 , 33, 2342-8	14.6	34
136	Influence of common variants near INSIG2, in FTO, and near MC4R genes on overweight and the metabolic profile in adolescence: the TRAILS (TRacking AdolescentsIndividual Lives Survey) Study. <i>American Journal of Clinical Nutrition</i> , 2010 , 91, 321-8	7	41
135	Association between birth weight and visceral fat in adults. <i>American Journal of Clinical Nutrition</i> , 2010 , 92, 347-52	7	75
134	Changes in renal risk factors versus renal function outcome during follow-up in a population-based cohort study. <i>Nephrology Dialysis Transplantation</i> , 2010 , 25, 1846-53	4.3	9
133	Cardiometabolic treatment decisions in patients with type 2 diabetes: the role of repeated measurements and medication burden. <i>BMJ Quality and Safety</i> , 2010 , 19, 411-5	5.4	7
132	Plasma procalcitonin is associated with obesity, insulin resistance, and the metabolic syndrome. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2010 , 95, E26-31	5.6	40
131	Escherichia coli bacteriuria in female adults is associated with the development of hypertension. <i>International Journal of Infectious Diseases</i> , 2010 , 14, e304-7	10.5	6
130	How do parents of 4- to 5-year-old children perceive the weight of their children?. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2010 , 99, 263-7	3.1	36

129	The analysis of longitudinal quality of life measures with informative drop-out: a pattern mixture approach. <i>Quality of Life Research</i> , 2010 , 19, 137-48	3.7	18
128	Does physical activity modify the risk of obesity for type 2 diabetes: a review of epidemiological data. <i>European Journal of Epidemiology</i> , 2010 , 25, 5-12	12.1	51
127	The bioelectrical impedance vector migration in healthy infants. <i>Clinical Nutrition</i> , 2010 , 29, 222-6	5.9	12
126	Comparison of methods to assess body fat in non-obese six to seven-year-old children. <i>Clinical Nutrition</i> , 2010 , 29, 317-22	5.9	17
125	Markers of the hepatic component of the metabolic syndrome as predictors of mortality in renal transplant recipients. <i>American Journal of Transplantation</i> , 2010 , 10, 106-14	8.7	22
124	The LIFESTYLE study: costs and effects of a structured lifestyle program in overweight and obese subfertile women to reduce the need for fertility treatment and improve reproductive outcome. A randomised controlled trial. <i>BMC Women's Health</i> , 2010 , 10, 22	2.9	47
123	Cost-effectiveness of a screening strategy for Q fever among pregnant women in risk areas: a clustered randomized controlled trial. <i>BMC Women's Health</i> , 2010 , 10, 32	2.9	10
122	MRI-determined fat content of human liver, pancreas and kidney. <i>World Journal of Gastroenterology</i> , 2010 , 16, 1993-8	5.6	59
121	Clinical inertia in general practice: widespread and related to the outcome of diabetes care. <i>Family Practice</i> , 2009 , 26, 428-36	1.9	66
120	High protein intake associates with cardiovascular events but not with loss of renal function. <i>Journal of the American Society of Nephrology: JASN</i> , 2009 , 20, 1797-804	12.7	59
119	Cochrane review: Interventions for treating obesity in children. <i>Evidence-Based Child Health: A Cochrane Review Journal</i> , 2009 , 4, 1571-1729		50
118	Acute lymphoblastic leukemia and obesity: increased energy intake or decreased physical activity?. <i>Supportive Care in Cancer</i> , 2009 , 17, 103-6	3.9	48
117	Obesity and impaired renal function: potential for lifestyle intervention?. <i>European Journal of Epidemiology</i> , 2009 , 24, 275-80	12.1	12
116	Bivariate genetic modelling of the response to an oral glucose tolerance challenge: a gene x environment interaction approach. <i>Diabetologia</i> , 2009 , 52, 1048-55	10.3	4
115	Effects of blood pressure lowering and intensive glucose control on the incidence and progression of retinopathy in patients with type 2 diabetes mellitus: a randomised controlled trial. <i>Diabetologia</i> , 2009 , 52, 2027-36	10.3	129
114	Comparison of various measures for assessing medication refill adherence using prescription data. <i>Pharmacoepidemiology and Drug Safety</i> , 2009 , 18, 159-65	2.6	57
113	Refill adherence and polypharmacy among patients with type 2 diabetes in general practice. <i>Pharmacoepidemiology and Drug Safety</i> , 2009 , 18, 983-91	2.6	44
112	Measuring abdominal adiposity in 6 to 7-year-old children. <i>European Journal of Clinical Nutrition</i> , 2009 , 63, 835-41	5.2	42

111	Ultrasonography to quantify hepatic fat content: validation by 1H magnetic resonance spectroscopy. <i>Obesity</i> , 2009 , 17, 2239-44	8	35
110	Non-alcoholic fatty liver disease is associated with cardiovascular disease risk markers. <i>Obesity Reviews</i> , 2009 , 10, 412-9	10.6	59
109	HbA(1c) levels in non-diabetic Dutch children aged 8-9 years: the PIAMA birth cohort study. <i>Diabetic Medicine</i> , 2009 , 26, 122-7	3.5	10
108	Which factors are important in adults Suptake of a (pre)pandemic influenza vaccine?. <i>Vaccine</i> , 2009 , 28, 207-27	4.1	42
107	Interventions for treating obesity in children. <i>The Cochrane Library</i> , 2009 , CD001872	5.2	624
106	Randomised controlled trial of intensive multifactorial treatment for cardiovascular risk in patients with screen-detected type 2 diabetes: 1-year data from the ADDITION Netherlands study. <i>British Journal of General Practice</i> , 2009 , 59, 43-8	1.6	59
105	Interventions for treating obesity in children. <i>Sao Paulo Medical Journal</i> , 2009 , 127, 321-321	1.6	13
104	Do characteristics of practices and general practitioners influence the yield of diabetes screening in primary care? The ADDITION Netherlands study. <i>Scandinavian Journal of Primary Health Care</i> , 2008 , 26, 160-5	2.7	13
103	Cohort Profile: the GECKO Drenthe study, overweight programming during early childhood. <i>International Journal of Epidemiology</i> , 2008 , 37, 486-9	7.8	41
102	Influence of elevated cardiometabolic risk factor levels on treatment changes in type 2 diabetes. <i>Diabetes Care</i> , 2008 , 31, 501-3	14.6	14
101	Overall quality of diabetes care in a defined geographic region: different sides of the same story. <i>British Journal of General Practice</i> , 2008 , 58, 339-45	1.6	15
100	Differences in the pattern of antibiotic prescription profile and recurrence rate for possible urinary tract infections in women with and without diabetes. <i>Diabetes Care</i> , 2008 , 31, 1380-5	14.6	35
99	Retinal vascular lesions in patients of Caucasian and Asian origin with type 2 diabetes: baseline results from the ADVANCE Retinal Measurements (AdRem) study. <i>Diabetes Care</i> , 2008 , 31, 708-13	14.6	37
98	Gender differences in predictors of the decline of renal function in the general population. <i>Kidney International</i> , 2008 , 74, 505-12	9.9	87
97	Homocysteine, S-adenosylmethionine and S-adenosylhomocysteine are associated with retinal microvascular abnormalities: the Hoorn Study. <i>Clinical Science</i> , 2008 , 114, 479-87	6.5	20
96	Association between depressive symptoms in childhood and adolescence and overweight in later life: review of the recent literature. <i>JAMA Pediatrics</i> , 2008 , 162, 981-8		52
95	Predictors for starting depot administration of risperidone in chronic users of antipsychotics. <i>Journal of Clinical Psychopharmacology</i> , 2008 , 28, 625-30	1.7	8
94	Universal risk factors for multifactorial diseases: LifeLines: a three-generation population-based study. <i>European Journal of Epidemiology</i> , 2008 , 23, 67-74	12.1	226

93	Stepwise screening for diabetes identifies people with high but modifiable coronary heart disease risk. The ADDITION study. <i>Diabetologia</i> , 2008 , 51, 1127-34	10.3	76
92	Three years follow-up of screen-detected diabetic and non-diabetic subjects: who is better off? The ADDITION Netherlands study. <i>BMC Family Practice</i> , 2008 , 9, 67	2.6	8
91	Screen detected subjects with type 2 diabetes and impaired glucose tolerance have more adverse cardiovascular risk than subjects with impaired fasting glucose especially when they are obese: the ADDITION Netherlands study. <i>Primary Care Diabetes</i> , 2007 , 1, 69-74	2.4	9
90	Shared and delegated systems are not quick remedies for improving diabetes care: a systematic review. <i>Primary Care Diabetes</i> , 2007 , 1, 59-68	2.4	14
89	High dietary glycemic load and glycemic index increase risk of cardiovascular disease among middle-aged women: a population-based follow-up study. <i>Journal of the American College of Cardiology</i> , 2007 , 50, 14-21	15.1	125
88	Accident proneness, does it exist? A review and meta-analysis. <i>Accident Analysis and Prevention</i> , 2007 , 39, 556-64	6.1	72
87	Abdominal fat and risk of coronary heart disease in patients with peripheral arterial disease. <i>Obesity</i> , 2007 , 15, 1623-30	8	5
86	Low yield of population-based screening for Type 2 diabetes in the Netherlands: the ADDITION Netherlands study. <i>Family Practice</i> , 2007 , 24, 555-61	1.9	42
85	Common genetic variations in CCK, leptin, and leptin receptor genes are associated with specific human eating patterns. <i>Diabetes</i> , 2007 , 56, 276-80	0.9	65
84	Screening for diabetes. <i>BMJ, The</i> , 2007 , 335, 457-8	5.9	4
83	Association between Escherichia coli bacteriuria and renal function in women: long-term follow-up. <i>Archives of Internal Medicine</i> , 2007 , 167, 253-7		7
82	Rationale and design of the AdRem study: evaluating the effects of blood pressure lowering and intensive glucose control on vascular retinal disorders in patients with type 2 diabetes mellitus. <i>Contemporary Clinical Trials</i> , 2007 , 28, 6-17	2.3	19
81	The threshold for diagnosing impaired fasting glucose: a position statement by the European Diabetes Epidemiology Group. <i>Diabetologia</i> , 2006 , 49, 822-7	10.3	87
80	Determinants of daily insulin use in Type 1 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2006 , 20, 356-60	3.2	11
79	Asymptomatic bacteriuria in women with diabetes mellitus: effect on renal function after 6 years of follow-up. <i>Archives of Internal Medicine</i> , 2006 , 166, 2222-7		42
78	Hyperglycemia and diabetes in patients with schizophrenia or schizoaffective disorders. <i>Diabetes Care</i> , 2006 , 29, 786-91	14.6	83
77	Is a single definition of the metabolic syndrome appropriate?--A comparative study of the USA and Asia. <i>Atherosclerosis</i> , 2006 , 184, 225-32	3.1	50
76	Lower-extremity arterial calcification as a correlate of coronary artery calcification. <i>Metabolism: Clinical and Experimental</i> , 2006 , 55, 1689-96	12.7	29

75	Are retinal microvascular abnormalities associated with large artery endothelial dysfunction and intima-media thickness? The Hoorn Study. <i>Clinical Science</i> , 2006 , 110, 597-604	6.5	56
74	Noninferiority testing in cost-minimization studies: Practical issues concerning power analysis. <i>International Journal of Technology Assessment in Health Care</i> , 2006 , 22, 261-6	1.8	5
73	The effect of moderate alcohol consumption on fat distribution and adipocytokines. <i>Obesity</i> , 2006 , 14, 60-6	8	72
72	High cumulative insulin exposure: a risk factor of atherosclerosis in type 1 diabetes?. <i>Atherosclerosis</i> , 2005 , 181, 185-92	3.1	26
71	Serial echocardiographic assessment of left ventricular mass. <i>Journal of Hypertension</i> , 2005 , 23, 460-1; author reply 461-2	1.9	1
70	Insulin treatment and cardiovascular disease; friend or foe? A point of view. <i>Diabetic Medicine</i> , 2005 , 22, 118-26	3.5	44
69	Fat distribution is strongly associated with plasma glucose levels and diabetes in Thai adults-the InterASIA study. <i>Diabetologia</i> , 2005 , 48, 657-60	10.3	25
68	Alcohol consumption and risk of type 2 diabetes among older women. <i>Diabetes Care</i> , 2005 , 28, 2933-8	14.6	60
67	Long-term effects of self-management education for patients with Type 2 diabetes taking maximal oral hypoglycaemic therapy: a randomized trial in primary care. <i>Diabetic Medicine</i> , 2004 , 21, 491-6	3.5	73
66	Initiation of glucose-lowering therapy in Type 2 diabetes mellitus patients in general practice. <i>Diabetic Medicine</i> , 2004 , 21, 896-900	3.5	10
65	History taking and leukocyturia predict the presence of asymptomatic bacteriuria in women with diabetes mellitus. <i>European Journal of Epidemiology</i> , 2004 , 19, 1021-7	12.1	4
64	Antihypertensive drug therapy and the risk of lower extremity amputations in pharmacologically treated type 2 diabetes patients. <i>Pharmacoepidemiology and Drug Safety</i> , 2004 , 13, 139-46	2.6	23
63	Patient characteristics do not predict poor glycaemic control in type 2 diabetes patients treated in primary care. <i>European Journal of Epidemiology</i> , 2004 , 19, 541-5	12.1	71
62	alpha-Tocopherol levels in plasma in new-onset, insulin-dependent diabetes mellitus. <i>European Journal of Internal Medicine</i> , 2004 , 15, 371-374	3.9	4
61	Insulin monotherapy versus combinations of insulin with oral hypoglycaemic agents in patients with type 2 diabetes mellitus. <i>The Cochrane Library</i> , 2004 , CD003418	5.2	48
60	Antipsychotic drugs may worsen metabolic control in type 2 diabetes mellitus. <i>Journal of Clinical Psychiatry</i> , 2004 , 65, 674-8	4.6	18
59	The prevalence and management of diabetes in Thai adults: the international collaborative study of cardiovascular disease in Asia. <i>Diabetes Care</i> , 2003 , 26, 2758-63	14.6	88
58	Atorvastatin dose-dependently decreases hepatic lipase activity in type 2 diabetes: effect of sex and the LIPC promoter variant. <i>Diabetes Care</i> , 2003 , 26, 427-32	14.6	44

57	Ultrasound measurements of intraabdominal fat estimate the metabolic syndrome better than do measurements of waist circumference. <i>American Journal of Clinical Nutrition</i> , 2003 , 77, 857-60	7	83
56	Quality of life of patients with type I diabetes mellitus. <i>Quality of Life Research</i> , 2003 , 12, 1089-97	3.7	50
55	Improving glycaemic control in patients with Type 2 diabetes mellitus without insulin therapy. <i>Diabetic Medicine</i> , 2003 , 20, 540-4	3.5	19
54	Refill compliance in type 2 diabetes mellitus: a predictor of switching to insulin therapy?. <i>Pharmacoepidemiology and Drug Safety</i> , 2003 , 12, 121-7	2.6	33
53	Common cholesteryl ester transfer protein gene polymorphisms and the effect of atorvastatin therapy in type 2 diabetes. <i>Diabetes Care</i> , 2003 , 26, 1216-23	14.6	98
52	Quality of recording of data from patients with type 2 diabetes is not a valid indicator of quality of care. A cross-sectional study. <i>Family Practice</i> , 2003 , 20, 173-7	1.9	46
51	Use of fluorquinolones is associated with a reduced risk of coronary heart disease in diabetes mellitus type 2 patients. <i>European Heart Journal</i> , 2002 , 23, 1575-9	9.5	5
50	Primary and secondary prevention in cardiovascular disease: an old-fashioned concept?. <i>Journal of Internal Medicine</i> , 2002 , 251, 301-6	10.8	3
49	Angiographic distribution of lower extremity atherosclerosis in patients with and without diabetes. <i>Diabetic Medicine</i> , 2002 , 19, 366-70	3.5	72
48	Resource consumption and costs in Dutch patients with type 2 diabetes mellitus. Results from 29 general practices. <i>Diabetic Medicine</i> , 2002 , 19, 246-53	3.5	33
47	Aggressive lipid lowering does not improve endothelial function in type 2 diabetes: the Diabetes Atorvastatin Lipid Intervention (DALI) Study: a randomized, double-blind, placebo-controlled trial. <i>Diabetes Care</i> , 2002 , 25, 1211-6	14.6	86
46	Thyroid dysfunction during pregnancy and in the first postpartum year in women with diabetes mellitus type 1. <i>European Journal of Endocrinology</i> , 2002 , 147, 443-51	6.5	23
45	Tissue factor pathway inhibitor and other endothelium-dependent hemostatic factors in elderly individuals with normal or impaired glucose tolerance and type 2 diabetes. <i>Diabetes Care</i> , 2002 , 25, 1340-5	14.6	47
44	Health-related quality of life and treatment satisfaction in Dutch patients with type 2 diabetes. <i>Diabetes Care</i> , 2002 , 25, 458-63	14.6	354
43	Cardiovascular risk factors and diseases precede oral hypoglycaemic therapy in patients with type 2 diabetes mellitus. <i>Journal of Clinical Epidemiology</i> , 2002 , 55, 345-9	5.7	4
42	Study rationale and design of ADVANCE: action in diabetes and vascular disease--preterax and diamicon MR controlled evaluation. <i>Diabetologia</i> , 2001 , 44, 1118-20	10.3	148
41	Virulence factors of Escherichia coli isolated from urine of diabetic women with asymptomatic bacteriuria: correlation with clinical characteristics. <i>Antonie Van Leeuwenhoek</i> , 2001 , 80, 119-27	2.1	19
40	Validity and reproducibility of ultrasonography for the measurement of intra-abdominal adipose tissue. <i>International Journal of Obesity</i> , 2001 , 25, 1346-51	5.5	184

39	Ethnic differences in C-peptide levels and anti-GAD antibodies in South African patients with diabetic ketoacidosis. <i>QJM - Monthly Journal of the Association of Physicians</i> , 2001 , 94, 39-43	2.7	12
38	Cardiovascular drug use and hospitalizations attributable to type 2 diabetes. <i>Diabetes Care</i> , 2001 , 24, 1428-32	14.6	9
37	Consequences of asymptomatic bacteriuria in women with diabetes mellitus. <i>Archives of Internal Medicine</i> , 2001 , 161, 1421-7		74
36	Experts' Opinions on the profile of optimal care for patients with diabetes mellitus type 2 in the Netherlands. <i>Netherlands Journal of Medicine</i> , 2001 , 58, 225-31	0.5	3
35	Asymptomatic bacteriuria can be considered a diabetic complication in women with diabetes mellitus. <i>Advances in Experimental Medicine and Biology</i> , 2000 , 485, 309-14	3.6	20
34	Education integrated into structured general practice care for Type 2 diabetic patients results in sustained improvement of disease knowledge and self-care. <i>Diabetic Medicine</i> , 2000 , 17, 190-7	3.5	40
33	Cytokine secretion is impaired in women with diabetes mellitus. <i>European Journal of Clinical Investigation</i> , 2000 , 30, 995-1001	4.6	89
32	Management of type 2 diabetes: a challenge for patient and physician. <i>Patient Education and Counseling</i> , 2000 , 40, 187-94	3.1	66
31	Physical activity in elderly subjects with impaired glucose tolerance and newly diagnosed diabetes mellitus. <i>American Journal of Epidemiology</i> , 1999 , 149, 219-27	3.8	40
30	Prevalence of variants in candidate genes for type 2 diabetes mellitus in The Netherlands: the Rotterdam study and the Hoorn study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1999 , 84, 1002-8	5.6	42
29	Incidence and determinants of mortality and cardiovascular events in diabetes mellitus: a meta-analysis. <i>Vascular Medicine</i> , 1999 , 4, 67-75	3.3	35
28	Variants in the sulphonylurea receptor gene: association of the exon 16-3t variant with Type II diabetes mellitus in Dutch Caucasians. <i>Diabetologia</i> , 1999 , 42, 617-20	10.3	70
27	Insulin resistance syndrome and left ventricular mass in an elderly population (The Rotterdam Study). <i>American Journal of Cardiology</i> , 1999 , 84, 233-6, A9	3	24
26	Performance of a predictive model to identify undiagnosed diabetes in a health care setting. <i>Diabetes Care</i> , 1999 , 22, 213-9	14.6	141
25	Discontinuation of prophylaxis for <i>Pneumocystis carinii</i> pneumonia in HIV-1-infected patients treated with highly active antiretroviral therapy. <i>Lancet, The</i> , 1999 , 353, 201-3	40	167
24	Prevalence of deafness is increased in patients with type 2 diabetes in The Netherlands. <i>Diabetes Care</i> , 1999 , 22, 180	14.6	4
23	Prevalence of Variants in Candidate Genes for Type 2 Diabetes Mellitus in The Netherlands: The Rotterdam Study and the Hoorn Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1999 , 84, 1002-1006	5.6	37
22	Serum free IGF-I, total IGF-I, IGFBP-1 and IGFBP-3 levels in an elderly population: relation to age and sex steroid levels. <i>Clinical Endocrinology</i> , 1998 , 48, 471-8	3.4	69

21	The acute effect of dexamethasone on plasma leptin concentrations and the relationships between fasting leptin, the IGF-I/IGFBP system, dehydroepiandrosterone, androstenedione and testosterone in an elderly population. <i>Clinical Endocrinology</i> , 1998 , 48, 621-6	3.4	18
20	Lack of associations between serum leptin, a polymorphism in the gene for the beta 3-adrenergic receptor and glucose tolerance in the Dutch population. <i>Clinical Endocrinology</i> , 1998 , 49, 229-34	3.4	20
19	Interperson variability but intraperson stability of baseline plasma cortisol concentrations, and its relation to feedback sensitivity of the hypothalamo-pituitary-adrenal axis to a low dose of dexamethasone in elderly individuals. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1998 , 83, 47-54	5.6	231
18	A polymorphism in the glucocorticoid receptor gene may be associated with an increased sensitivity to glucocorticoids in vivo. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1998 , 83, 144-51	5.6	365
17	Serum total IGF-I, free IGF-I, and IGFB-1 levels in an elderly population: relation to cardiovascular risk factors and disease. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 1998 , 18, 277-82	9.4	201
16	Gender-specific relationship between serum free and total IGF-I and bone mineral density in elderly men and women. <i>European Journal of Endocrinology</i> , 1998 , 138, 627-32	6.5	46
15	Effects of nitrendipine and enalapril on left ventricular mass in patients with non-insulin-dependent diabetes mellitus and hypertension. <i>Journal of Hypertension</i> , 1998 , 16, 689-96	1.9	19
14	Serum free and total insulin-like growth factor-I, insulin-like growth factor binding protein-1 and insulin-like growth factor binding protein-3 Levels in healthy elderly individuals. Relation to self-reported quality of health and disability. <i>Gerontology</i> , 1998 , 44, 277-80	5.5	27
13	A Polymorphism in the Glucocorticoid Receptor Gene May Be Associated with an Increased Sensitivity to Glucocorticoids in Vivo. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1998 , 83, 144-151	5.6	343
12	Interperson Variability but Intraperson Stability of Baseline Plasma Cortisol Concentrations, and Its Relation to Feedback Sensitivity of the Hypothalamo-Pituitary-Adrenal Axis to a Low Dose of Dexamethasone in Elderly Individuals. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1998 , 83, 47-54	5.6	152
11	Insulin and cognitive function in an elderly population. The Rotterdam Study. <i>Diabetes Care</i> , 1997 , 20, 792-5	14.6	171
10	Lack of association between five polymorphisms in the human glucocorticoid receptor gene and glucocorticoid resistance. <i>Human Genetics</i> , 1997 , 99, 663-8	6.3	190
9	Gender differences in the associations between cortisol and insulin in healthy subjects. <i>Journal of Endocrinology</i> , 1996 , 149, 313-8	4.7	56
8	Plasma fibrinogen in NIDDM: the Rotterdam study. <i>Diabetes Care</i> , 1996 , 19, 157-9	14.6	35
7	Hyperinsulinemia and bone mineral density in an elderly population: The Rotterdam Study. <i>Bone</i> , 1996 , 18, 545-9	4.7	138
6	Insulin, hypertension and antihypertensive drugs in elderly patients: the Rotterdam Study. <i>Journal of Hypertension</i> , 1996 , 14, 237-42	1.9	6
5	Association of diabetes mellitus and dementia: the Rotterdam Study. <i>Diabetologia</i> , 1996 , 39, 1392-7	10.3	502
4	Bone density in non-insulin-dependent diabetes mellitus. The Rotterdam Study. <i>Annals of Internal Medicine</i> , 1995 , 122, 409-14	8	221

- 3 Why use the oral glucose tolerance test?. *Diabetes Care*, **1995**, 18, 1045-9 14.6 47
- 2 Retinopathy, glucose, and insulin in an elderly population. The Rotterdam Study. *Diabetes*, **1995**, 44, 11-159 26
- 1 High blood pressure and the incidence of non-insulin dependent diabetes mellitus: findings in a 11.5 year follow-up study in The Netherlands. *European Journal of Epidemiology*, **1993**, 9, 134-9 12.1 23