

Markus Barth

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/2643602/markus-barth-publications-by-year.pdf>

Version: 2024-04-29

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

109
papers

4,273
citations

38
h-index

62
g-index

126
ext. papers

5,087
ext. citations

5.5
avg, IF

5.49
L-index

#	Paper	IF	Citations
109	QSMxT: Robust masking and artifact reduction for quantitative susceptibility mapping. <i>Magnetic Resonance in Medicine</i> , 2021 ,	4.4	1
108	Improving FLAIR SAR efficiency at 7T by adaptive tailoring of adiabatic pulse power through deep learning estimation. <i>Magnetic Resonance in Medicine</i> , 2021 , 85, 2462-2476	4.4	1
107	Open-access quantitative MRI data of the spinal cord and reproducibility across participants, sites and manufacturers. <i>Scientific Data</i> , 2021 , 8, 219	8.2	6
106	7-Tesla Functional Cardiovascular MR Using Vectorcardiographic Triggering-Overcoming the Magnetohydrodynamic Effect. <i>Tomography</i> , 2021 , 7, 323-332	3.1	
105	Generic acquisition protocol for quantitative MRI of the spinal cord. <i>Nature Protocols</i> , 2021 , 16, 4611-4638	13.8	11
104	Improved susceptibility weighted imaging at ultra-high field using bipolar multi-echo acquisition and optimized image processing: CLEAR-SWI. <i>NeuroImage</i> , 2021 , 237, 118175	7.9	6
103	Influence of 7T GRE-MRI Signal Compartment Model Choice on Tissue Parameters. <i>Frontiers in Neuroscience</i> , 2020 , 14, 271	5.1	2
102	Bayesian population receptive field modeling in human somatosensory cortex. <i>NeuroImage</i> , 2020 , 208, 116465	7.9	19
101	Towards Optimising MRI Characterisation of Tissue (TOMCAT) Dataset including all Longitudinal Automatic Segmentation of Hippocampal Subfields (LASHiS) data. <i>Data in Brief</i> , 2020 , 32, 106043	1.2	
100	Field strength influences on gradient recalled echo MRI signal compartment frequency shifts. <i>Magnetic Resonance Imaging</i> , 2020 , 70, 98-107	3.3	0
99	New acquisition techniques and their prospects for the achievable resolution of fMRI. <i>Progress in Neurobiology</i> , 2020 , 207, 101936	10.9	7
98	Longitudinal Automatic Segmentation of Hippocampal Subfields (LASHiS) using multi-contrast MRI. <i>NeuroImage</i> , 2020 , 218, 116798	7.9	4
97	Non-linear realignment improves hippocampus subfield segmentation reliability. <i>NeuroImage</i> , 2019 , 203, 116206	7.9	6
96	7T GRE-MRI signal compartments are sensitive to dysplastic tissue in focal epilepsy. <i>Magnetic Resonance Imaging</i> , 2019 , 61, 1-8	3.3	7
95	DeepQSM - using deep learning to solve the dipole inversion for quantitative susceptibility mapping. <i>NeuroImage</i> , 2019 , 195, 373-383	7.9	48
94	A numerical and experimental study of RF shimming in the presence of hip prostheses using adaptive SAR at 3 T. <i>Magnetic Resonance in Medicine</i> , 2019 , 81, 3826-3839	4.4	4
93	SHARQnet - Sophisticated harmonic artifact reduction in quantitative susceptibility mapping using a deep convolutional neural network. <i>Zeitschrift Fur Medizinische Physik</i> , 2019 , 29, 139-149	7.6	9

92	Adaptive SAR mass-averaging framework to improve predictions of local RF heating near a hip implant for parallel transmit at 7 T. <i>Magnetic Resonance in Medicine</i> , 2019 , 81, 615-627	4.4	11
91	Patient with ALS with a novel TBK1 mutation, widespread brain involvement, behaviour changes and metabolic dysfunction. <i>Journal of Neurology, Neurosurgery and Psychiatry</i> , 2019 , 90, 952-954	5.5	4
90	PEC-GRAPPA reconstruction of simultaneous multislice EPI with slice-dependent 2D Nyquist ghost correction. <i>Magnetic Resonance in Medicine</i> , 2019 , 81, 1924-1934	4.4	7
89	Cardiac Magnetic Resonance Imaging at 7 Tesla. <i>Journal of Visualized Experiments</i> , 2019 ,	1.6	6
88	Robust SENSE reconstruction of simultaneous multislice EPI with low-rank enhanced coil sensitivity calibration and slice-dependent 2D Nyquist ghost correction. <i>Magnetic Resonance in Medicine</i> , 2018 , 80, 1376-1390	4.4	10
87	A method for the dynamic correction of B-related distortions in single-echo EPI at 7T. <i>NeuroImage</i> , 2018 , 168, 321-331	7.9	36
86	The challenge of bias-free coil combination for quantitative susceptibility mapping at ultra-high field. <i>Magnetic Resonance in Medicine</i> , 2018 , 79, 97-107	4.4	12
85	Serial correlations in single-subject fMRI with sub-second TR. <i>NeuroImage</i> , 2018 , 166, 152-166	7.9	38
84	Using multi-echo simultaneous multi-slice (SMS) EPI to improve functional MRI of the subcortical nuclei of the basal ganglia at ultra-high field (7T). <i>NeuroImage</i> , 2018 , 172, 886-895	7.9	15
83	Assessment of microstructural signal compartments across the corpus callosum using multi-echo gradient recalled echo at 7T. <i>NeuroImage</i> , 2018 , 182, 407-416	7.9	15
82	Cued reactivation during slow-wave sleep induces brain connectivity changes related to memory stabilization. <i>Scientific Reports</i> , 2018 , 8, 16958	4.9	13
81	Echo time-dependent quantitative susceptibility mapping contains information on tissue properties. <i>Magnetic Resonance in Medicine</i> , 2017 , 77, 1946-1958	4.4	39
80	A time-efficient acquisition protocol for multipurpose diffusion-weighted microstructural imaging at 7 Tesla. <i>Magnetic Resonance in Medicine</i> , 2017 , 78, 2170-2184	4.4	16
79	Diffusion parameter mapping with the combined intravoxel incoherent motion and kurtosis model using artificial neural networks at 3T. <i>NMR in Biomedicine</i> , 2017 , 30, e3833	4.4	30
78	Measuring the effects of attention to individual fingertips in somatosensory cortex using ultra-high field (7T) fMRI. <i>NeuroImage</i> , 2017 , 161, 179-187	7.9	25
77	Accelerated mapping of magnetic susceptibility using 3D planes-on-a-paddlewheel (POP) EPI at ultra-high field strength. <i>NMR in Biomedicine</i> , 2017 , 30, e3620	4.4	4
76	Combining phase images from array coils using a short echo time reference scan (COMPOSER). <i>Magnetic Resonance in Medicine</i> , 2017 , 77, 318-327	4.4	37
75	A cortical vascular model for examining the specificity of the laminar BOLD signal. <i>NeuroImage</i> , 2016 , 132, 491-498	7.9	90

74	ECG Triggering in Ultra-High Field Cardiovascular MRI. <i>Tomography</i> , 2016 , 2, 167-174	3.1	11
73	Selective channel combination of MRI signal phase. <i>Magnetic Resonance in Medicine</i> , 2016 , 76, 1469-1477	4.4	7
72	Pulsed arterial spin labelling at ultra-high field with a B 1 (+) -optimised adiabatic labelling pulse. <i>Magnetic Resonance Materials in Physics, Biology, and Medicine</i> , 2016 , 29, 463-73	2.8	10
71	Correcting dynamic distortions in 7T echo planar imaging using a jittered echo time sequence. <i>Magnetic Resonance in Medicine</i> , 2016 , 76, 1388-1399	4.4	16
70	Simultaneous multislice (SMS) imaging techniques. <i>Magnetic Resonance in Medicine</i> , 2016 , 75, 63-81	4.4	280
69	Diffusion tensor characteristics of gyrencephaly using high resolution diffusion MRI in vivo at 7T. <i>NeuroImage</i> , 2015 , 109, 378-87	7.9	39
68	Improved sensitivity and specificity for resting state and task fMRI with multiband multi-echo EPI compared to multi-echo EPI at 7 T. <i>NeuroImage</i> , 2015 , 119, 352-61	7.9	43
67	Fast quantitative susceptibility mapping using 3D EPI and total generalized variation. <i>NeuroImage</i> , 2015 , 111, 622-30	7.9	112
66	Online decoding of object-based attention using real-time fMRI. <i>European Journal of Neuroscience</i> , 2014 , 39, 319-29	3.5	11
65	BOLD fMRI signal characteristics of S1- and S2-SSFP at 7 Tesla. <i>Frontiers in Neuroscience</i> , 2014 , 8, 49	5.1	19
64	A study-specific fMRI normalization approach that operates directly on high resolution functional EPI data at 7 Tesla. <i>NeuroImage</i> , 2014 , 100, 710-4	7.9	16
63	Application of PINS radiofrequency pulses to reduce power deposition in RARE/turbo spin echo imaging of the human head. <i>Magnetic Resonance in Medicine</i> , 2014 , 71, 44-9	4.4	36
62	Whole brain, high resolution multiband spin-echo EPI fMRI at 7 T: a comparison with gradient-echo EPI using a color-word Stroop task. <i>NeuroImage</i> , 2014 , 97, 142-50	7.9	38
61	Single-shot echo-planar imaging with Nyquist ghost compensation: interleaved dual echo with acceleration (IDEA) echo-planar imaging (EPI). <i>Magnetic Resonance in Medicine</i> , 2013 , 69, 37-47	4.4	16
60	Simultaneous multislice inversion contrast imaging using power independent of the number of slices (PINS) and delays alternating with nutation for tailored excitation (DANTE) radio frequency pulses. <i>Magnetic Resonance in Medicine</i> , 2013 , 69, 1670-6	4.4	13
59	Linear reconstruction of perceived images from human brain activity. <i>NeuroImage</i> , 2013 , 83, 951-61	7.9	71
58	Generalized INverse imaging (GIN): ultrafast fMRI with physiological noise correction. <i>Magnetic Resonance in Medicine</i> , 2013 , 70, 962-71	4.4	27
57	Layer-specific diffusion weighted imaging in human primary visual cortex in vitro. <i>Cortex</i> , 2013 , 49, 2569-88	3.8	40

56	Electrophysiological correlation patterns of resting state networks in single subjects: a combined EEG-fMRI study. <i>Brain Topography</i> , 2013 , 26, 98-109	4.3	26
55	Modeling and suppression of respiration induced B0-fluctuations in non-balanced steady-state free precession sequences at 7 Tesla. <i>Magnetic Resonance Materials in Physics, Biology, and Medicine</i> , 2013 , 26, 377-87	2.8	2
54	An Investigation of RSN Frequency Spectra Using Ultra-Fast Generalized Inverse Imaging. <i>Frontiers in Human Neuroscience</i> , 2013 , 7, 156	3.3	26
53	The Quest for EEG Power Band Correlation with ICA Derived fMRI Resting State Networks. <i>Frontiers in Human Neuroscience</i> , 2013 , 7, 315	3.3	14
52	Structure Tensor Informed Fiber Tractography (STIFT) by combining gradient echo MRI and diffusion weighted imaging. <i>NeuroImage</i> , 2012 , 59, 3941-54	7.9	15
51	Whole brain, high resolution spin-echo resting state fMRI using PINS multiplexing at 7 T. <i>NeuroImage</i> , 2012 , 62, 1939-46	7.9	53
50	Sleep supports selective retention of associative memories based on relevance for future utilization. <i>PLoS ONE</i> , 2012 , 7, e43426	3.7	79
49	Reference-free unwarping of EPI data using dynamic off-resonance correction with multiecho acquisition (DOCMA). <i>Magnetic Resonance in Medicine</i> , 2012 , 68, 1247-54	4.4	25
48	Memory stabilization with targeted reactivation during human slow-wave sleep. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012 , 109, 10575-80	11.5	96
47	Multi-echo fMRI of the cortical laminae in humans at 7 T. <i>NeuroImage</i> , 2011 , 56, 1276-85	7.9	117
46	Functional connectivity during light sleep is correlated with memory performance for face-location associations. <i>NeuroImage</i> , 2011 , 57, 262-270	7.9	41
45	7 tesla MRI of microbleeds and white matter lesions as seen in vascular dementia. <i>Journal of Magnetic Resonance Imaging</i> , 2011 , 33, 782-91	5.6	65
44	Power Independent of Number of Slices (PINS) radiofrequency pulses for low-power simultaneous multislice excitation. <i>Magnetic Resonance in Medicine</i> , 2011 , 66, 1234-40	4.4	99
43	Advances in High-Field BOLD fMRI. <i>Materials</i> , 2011 , 4, 1941-1955	3.5	16
42	Susceptibility weighted magnetic resonance imaging of cerebral cavernous malformations: prospects, drawbacks, and first experience at ultra-high field strength (7-Tesla) magnetic resonance imaging. <i>Neurosurgical Focus</i> , 2010 , 29, E5	4.2	32
41	Three dimensional echo-planar imaging at 7 Tesla. <i>NeuroImage</i> , 2010 , 51, 261-6	7.9	197
40	Layer-specific BOLD activation in human V1. <i>Human Brain Mapping</i> , 2010 , 31, 1297-304	5.9	146
39	A population-specific symmetric phase model to automatically analyze susceptibility-weighted imaging (SWI) phase shifts and phase symmetry in the human brain. <i>Journal of Magnetic Resonance Imaging</i> , 2010 , 31, 215-20	5.6	9

38	Filtered deconvolution of a simulated and an in vivo phase model of the human brain. <i>Journal of Magnetic Resonance Imaging</i> , 2010 , 32, 289-97	5.6	9
37	T2-weighted 3D fMRI using S2-SSFP at 7 tesla. <i>Magnetic Resonance in Medicine</i> , 2010 , 63, 1015-20	4.4	34
36	Phase unwrapping of MR images using Phi UN--a fast and robust region growing algorithm. <i>Medical Image Analysis</i> , 2009 , 13, 257-68	15.4	76
35	Improved elimination of phase effects from background field inhomogeneities for susceptibility weighted imaging at high magnetic field strengths. <i>Magnetic Resonance Imaging</i> , 2008 , 26, 1145-51	3.3	35
34	MR venography of the human brain using susceptibility weighted imaging at very high field strength. <i>Magnetic Resonance Materials in Physics, Biology, and Medicine</i> , 2008 , 21, 149-58	2.8	99
33	T1 mapping of the entire lung parenchyma: Influence of respiratory phase and correlation to lung function test results in patients with diffuse lung disease. <i>Magnetic Resonance in Medicine</i> , 2008 , 59, 96-101	4.4	47
32	Very high-resolution three-dimensional functional MRI of the human visual cortex with elimination of large venous vessels. <i>NMR in Biomedicine</i> , 2007 , 20, 477-84	4.4	32
31	Contrast-to-noise ratio (CNR) as a quality parameter in fMRI. <i>Journal of Magnetic Resonance Imaging</i> , 2007 , 25, 1263-70	5.6	54
30	Comparison of fMRI coregistration results between human experts and software solutions in patients and healthy subjects. <i>European Radiology</i> , 2007 , 17, 1634-43	8	15
29	Contrast enhanced susceptibility weighted imaging (CE-SWI) of the mouse brain using ultrasmall superparamagnetic ironoxide particles (USPIO). <i>Zeitschrift Fur Medizinische Physik</i> , 2006 , 16, 269-74	7.6	10
28	Contrast-enhanced, high-resolution, susceptibility-weighted magnetic resonance imaging of the brain: dose-dependent optimization at 3 tesla and 1.5 tesla in healthy volunteers. <i>Investigative Radiology</i> , 2006 , 41, 249-55	10.1	39
27	Influence of fMRI smoothing procedures on replicability of fine scale motor localization. <i>NeuroImage</i> , 2005 , 24, 323-31	7.9	50
26	Evaluation of preoperative high magnetic field motor functional MRI (3 Tesla) in glioma patients by navigated electrocortical stimulation and postoperative outcome. <i>Journal of Neurology, Neurosurgery and Psychiatry</i> , 2005 , 76, 1152-7	5.5	112
25	fMRI reveals functional cortex in a case of inconclusive Wada testing. <i>Clinical Neurology and Neurosurgery</i> , 2005 , 107, 147-51	2	14
24	T1 mapping of the entire lung parenchyma: Influence of the respiratory phase in healthy individuals. <i>Journal of Magnetic Resonance Imaging</i> , 2005 , 21, 759-64	5.6	42
23	Noninvasive assessment of vascular architecture and function during modulated blood oxygenation using susceptibility weighted magnetic resonance imaging. <i>Magnetic Resonance in Medicine</i> , 2005 , 54, 87-95	4.4	118
22	Magnetic susceptibility-weighted MR phase imaging of the human brain. <i>American Journal of Neuroradiology</i> , 2005 , 26, 736-42	4.4	177
21	A quantitative comparison of functional MRI cluster analysis. <i>Artificial Intelligence in Medicine</i> , 2004 , 31, 57-71	7.4	68

20	Robust field map generation using a triple-echo acquisition. <i>Journal of Magnetic Resonance Imaging</i> , 2004 , 20, 730-4	5.6	50
19	Wavelet-based multifractal analysis of fMRI time series. <i>NeuroImage</i> , 2004 , 22, 1195-202	7.9	79
18	High-resolution three-dimensional contrast-enhanced blood oxygenation level-dependent magnetic resonance venography of brain tumors at 3 Tesla: first clinical experience and comparison with 1.5 Tesla. <i>Investigative Radiology</i> , 2003 , 38, 409-14	10.1	47
17	MR contrast agent at high-field MRI (3 Tesla). <i>Topics in Magnetic Resonance Imaging</i> , 2003 , 14, 365-75	2.3	46
16	High-resolution three-dimensional contrast-enhanced blood oxygenation level-dependent magnetic resonance venography of brain tumors at 3 Tesla: First clinical experience and comparison with 1.5 Tesla. <i>Investigative Radiology</i> , 2003 , 38, 409-414	10.1	17
15	Fuzzy cluster analysis of high-field functional MRI data. <i>Artificial Intelligence in Medicine</i> , 2003 , 29, 203-237	7.4	30
14	Automated unwrapping of MR phase images applied to BOLD MR-venography at 3 Tesla. <i>Journal of Magnetic Resonance Imaging</i> , 2003 , 18, 175-80	5.6	89
13	Scaling laws and persistence in human brain activity. <i>Physica A: Statistical Mechanics and Its Applications</i> , 2003 , 326, 511-521	3.3	37
12	Comparison of multi-echo spiral and echo planar imaging in functional MRI. <i>Magnetic Resonance Imaging</i> , 2002 , 20, 359-64	3.3	8
11	Magnetic resonance imaging contrast enhancement of brain tumors at 3 tesla versus 1.5 tesla. <i>Investigative Radiology</i> , 2002 , 37, 114-9	10.1	95
10	Characterization of BOLD activation in multi-echo fMRI data using fuzzy cluster analysis and a comparison with quantitative modeling. <i>NMR in Biomedicine</i> , 2001 , 14, 484-9	4.4	9
9	High-resolution MR venography at 3.0 Tesla. <i>Journal of Computer Assisted Tomography</i> , 2000 , 24, 949-57	2.2	173
8	High-resolution, multiple gradient-echo functional MRI at 1.5 T. <i>Magnetic Resonance Imaging</i> , 1999 , 17, 321-9	3.3	47
7	Functional MRI of the human motor cortex using single-shot, multiple gradient-echo spiral imaging. <i>Magnetic Resonance Imaging</i> , 1999 , 17, 1239-43	3.3	33
6	Explorative signal processing in functional MR imaging. <i>International Journal of Imaging Systems and Technology</i> , 1999 , 10, 166-176	2.5	22
5	Quantification of signal changes in gradient recalled echo FMRI. <i>Magnetic Resonance Imaging</i> , 1997 , 15, 753-62	3.3	10
4	Modulation of signal changes in gradient-recalled echo functional MRI with increasing echo time correlate with model calculations. <i>Magnetic Resonance Imaging</i> , 1997 , 15, 745-52	3.3	18
3	Functional Susceptibility Weighted Magnetic Resonance Imaging	561-575	

2 Automatic Vein Segmentation and Lesion Detection: From SWI-MIPs to MR Venograms 6056-618

1 QSMxT: Robust Masking and Artefact Reduction for Quantitative Susceptibility Mapping

1