
Martin Klein

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/264807/publications.pdf Version: 2024-02-01



| # | Article | IF | CITATIONS |
|----|---|------|-----------|
| 1 | Lomustine and Bevacizumab in Progressive Glioblastoma. New England Journal of Medicine, 2017, 377, 1954-1963. | 13.9 | 670 |
| 2 | Cognitive deficits in adult patients with brain tumours. Lancet Neurology, The, 2004, 3, 159-168. | 4.9 | 641 |
| 3 | Cognitive and radiological effects of radiotherapy in patients with low-grade glioma: long-term follow-up. Lancet Neurology, The, 2009, 8, 810-818. | 4.9 | 598 |
| 4 | Epilepsy in low-grade gliomas: The impact on cognitive function and quality of life. Annals of Neurology, 2003, 54, 514-520. | 2.8 | 358 |
| 5 | Cognitive Rehabilitation in Patients With Gliomas: A Randomized, Controlled Trial. Journal of Clinical Oncology, 2009, 27, 3712-3722. | 0.8 | 294 |
| 6 | Disturbed functional connectivity in brain tumour patients: Evaluation by graph analysis of synchronization matrices. Clinical Neurophysiology, 2006, 117, 2039-2049. | 0.7 | 257 |
| 7 | Precuneus atrophy in early-onset Alzheimer's disease: a morphometric structural MRI study. Neuroradiology, 2007, 49, 967-976. | 1.1 | 251 |
| 8 | Neurobehavioral Status and Health-Related Quality of Life in Newly Diagnosed High-Grade Glioma Patients. Journal of Clinical Oncology, 2001, 19, 4037-4047. | 0.8 | 232 |
| 9 | Patients With Alzheimer Disease With Multiple Microbleeds. Stroke, 2009, 40, 3455-3460. | 1.0 | 202 |
| 10 | Symptoms and problems in the end-of-life phase of high-grade glioma patients. Neuro-Oncology, 2010, 12, 1162-1166. | 0.6 | 171 |
| 11 | How do brain tumors alter functional connectivity? A magnetoencephalography study. Annals of Neurology, 2006, 59, 128-138. | 2.8 | 164 |
| 12 | Compromised Health-Related Quality of Life in Patients With Low-Grade Glioma. Journal of Clinical Oncology, 2011, 29, 4430-4435. | 0.8 | 160 |
| 13 | The course of neurocognitive functioning in high-grade glioma patients1. Neuro-Oncology, 2007, 9, 53-62. | 0.6 | 153 |
| 14 | Epilepsy is related to theta band brain connectivity and network topology in brain tumor patients. BMC Neuroscience, 2010, 11, 103. | 0.8 | 145 |
| 15 | Cognition and resective surgery for diffuse infiltrative glioma: an overview. Journal of Neuro-Oncology, 2012, 108, 309-318. | 1.4 | 129 |
| 16 | Synchronized brain activity and neurocognitive function in patients with low-grade glioma: A magnetoencephalography study. Neuro-Oncology, 2008, 10, 734-744. | 0.6 | 119 |
| 17 | The Effectiveness of Cognitive Rehabilitation for Attention Deficits in Focal Seizures: A Randomized Controlled Study. Epilepsia, 2002, 43, 587-595. | 2.6 | 117 |
| 18 | Disturbed functional brain networks and neurocognitive function in low-grade glioma patients: a graph theoretical analysis of resting-state MEG. Nonlinear Biomedical Physics, 2009, 3, 9. | 1.5 | 116 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Resting-State Brain Networks in Type 1 Diabetic Patients With and Without Microangiopathy and Their Relation to Cognitive Functions and Disease Variables. Diabetes, 2012, 61, 1814-1821. | 0.3 | 109 |
| 20 | The effect of modafinil on fatigue, cognitive functioning, and mood in primary brain tumor patients: a multicenter randomized controlled trial. Neuro-Oncology, 2013, 15, 1420-1428. | 0.6 | 109 |
| 21 | Differential effect of surgery and radiotherapy on neurocognitive functioning and health-related quality of life in WHO grade I meningioma patients. Journal of Neuro-Oncology, 2007, 84, 271-278. | 1.4 | 105 |
| 22 | Large and fast human pyramidal neurons associate with intelligence. ELife, 2018, 7, . | 2.8 | 103 |
| 23 | Decision-making in the end-of-life phase of high-grade glioma patients. European Journal of Cancer, 2012, 48, 226-232. | 1.3 | 102 |
| 24 | Tumour and surgery effects on cognitive functioning in high-grade glioma patients. Acta Neurochirurgica, 2014, 156, 1451-1459. | 0.9 | 100 |
| 25 | MEG Network Differences between Low- and High-Grade Glioma Related to Epilepsy and Cognition. PLoS ONE, 2012, 7, e50122. | 1.1 | 100 |
| 26 | Health-related quality of life in patients with high-risk low-grade glioma (EORTC 22033-26033): a randomised, open-label, phase 3 intergroup study. Lancet Oncology, The, 2016, 17, 1533-1542. | 5.1 | 97 |
| 27 | Enhancing quality of life and mastery of informal caregivers of high-grade glioma patients: a randomized controlled trial. Journal of Neuro-Oncology, 2013, 111, 303-311. | 1.4 | 96 |
| 28 | Long-Term Persisting Cognitive Sequelae of Traumatic Brain Injury and the Effect of Age. Journal of Nervous and Mental Disease, 1996, 184, 459-467. | 0.5 | 89 |
| 29 | Neurocognitive functioning and health-related quality of life in patients treated with stereotactic radiotherapy for brain metastases: a prospective study. Neuro-Oncology, 2015, 18, 435-444. | 0.6 | 89 |
| 30 | Effect of test duration on age-related differences in stroop interference. Journal of Clinical and Experimental Neuropsychology, 1997, 19, 77-82. | 0.8 | 88 |
| 31 | CSF α-Synuclein Does Not Discriminate Dementia with Lewy Bodies from Alzheimer's Disease. Journal of Alzheimer's Disease, 2010, 22, 87-95. | 1.2 | 87 |
| 32 | Long-term Impact of Cognitive Deficits and Epilepsy on Quality of Life in Patients With Low-Grade Meningiomas. Neurosurgery, 2011, 69, 72-79. | 0.6 | 86 |
| 33 | Health-related quality of life and cognitive functioning in long-term anaplastic oligodendroglioma and oligoastrocytoma survivors. Journal of Neuro-Oncology, 2014, 116, 161-168. | 1.4 | 83 |
| 34 | Survey on current cognitive practices within the European Low-Grade Glioma Network: towards a European assessment protocol. Acta Neurochirurgica, 2017, 159, 1167-1178. | 0.9 | 80 |
| 35 | Bevacizumab and temozolomide in patients with first recurrence of WHO grade II and III glioma, without 1p/19q co-deletion (TAVAREC): a randomised controlled phase 2 EORTC trial. Lancet Oncology, The, 2018, 19, 1170-1179. | 5.1 | 80 |
| 36 | Fatigue in low-grade glioma. Journal of Neuro-Oncology, 2009, 92, 73-78. | 1.4 | 77 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Psychiatric symptoms in glioma patients: from diagnosis to management. Neuropsychiatric Disease and Treatment, 2015, 11, 1413. | 1.0 | 72 |
| 38 | Cognitive functioning in glioblastoma patients during radiotherapy and temozolomide treatment: initial findings. Journal of Neuro-Oncology, 2010, 97, 89-94. | 1.4 | 71 |
| 39 | Symptom management and quality of life in glioma patients. CNS Oncology, 2014, 3, 37-47. | 1.2 | 71 |
| 40 | Health-related quality of life of long-term high-grade glioma survivors. Neuro-Oncology, 2009, 11, 51-58. | 0.6 | 70 |
| 41 | Functional Brain Connectivity and Neurocognitive Functioning in Patients With Long-Standing Type 1 Diabetes With and Without Microvascular Complications. Diabetes, 2009, 58, 2335-2343. | 0.3 | 67 |
| 42 | Cognitive Impairment in Alzheimer's Disease Is Modified by APOE Genotype. Dementia and Geriatric Cognitive Disorders, 2007, 24, 98-103. | 0.7 | 66 |
| 43 | Neurocognitive functioning in adult WHO grade II gliomas: impact of old and new treatment modalities. Neuro-Oncology, 2012, 14, iv17-iv24. | 0.6 | 65 |
| 44 | Treatment-related changes in functional connectivity in brain tumor patients: A magnetoencephalography study. Experimental Neurology, 2008, 212, 285-290. | 2.0 | 64 |
| 45 | Health-Related Quality of Life in Stable, Long-Term Survivors of Low-Grade Glioma. Journal of Clinical Oncology, 2015, 33, 1023-1029. | 0.8 | 64 |
| 46 | Proliferative Retinopathy in Type 1 Diabetes Is Associated With Cerebral Microbleeds, Which Is Part of Generalized Microangiopathy. Diabetes Care, 2014, 37, 1165-1168. | 4.3 | 61 |
| 47 | CODEL: phase III study of RT, RT + TMZ, or TMZ for newly diagnosed 1p/19q codeleted oligodendroglioma. Analysis from the initial study design. Neuro-Oncology, 2021, 23, 457-467. | 0.6 | 58 |
| 48 | Levetiracetam improves verbal memory in high-grade glioma patients. Neuro-Oncology, 2013, 15, 216-223. | 0.6 | 57 |
| 49 | Health-related quality of life of significant others of patients with malignant CNS versus non-CNS tumors: a comparative study. Journal of Neuro-Oncology, 2013, 115, 87-94. | 1.4 | 47 |
| 50 | EORTC QLQ-C15-PAL: the new standard in the assessment of health-related quality of life in advanced cancer?. Palliative Medicine, 2006, 20, 1-2. | 1.3 | 46 |
| 51 | Docosahexaenoic Acid Slows Visual Field Progression in X-Linked Retinitis Pigmentosa: Ancillary Outcomes of the DHAX Trial. , 2015, 56, 6646. | | 46 |
| 52 | Internet-based guided self-help for glioma patients with depressive symptoms: a randomized controlled trial. Journal of Neuro-Oncology, 2018, 137, 191-203. | 1.4 | 46 |
| 53 | Cortical Thickness, Surface Area and Subcortical Volume Differentially Contribute to Cognitive Heterogeneity in Parkinson's Disease. PLoS ONE, 2016, 11, e0148852. | 1.1 | 44 |
| 54 | Detecting Mild Cognitive Deficits in <scp>P</scp> arkinson's <scp>D</scp> isease: <scp>C</scp> omparison of <scp>N</scp> europsychological <scp>T</scp> ests. Movement Disorders, 2018, 33, 1750-1759. | 2.2 | 42 |

| # | Article | IF | CITATIONS |
|----|---|------------------|------------------|
| 55 | Understanding cognitive functioning in glioma patients: The relevance of IDHâ€mutation status and functional connectivity. Brain and Behavior, 2019, 9, e01204. | 1.0 | 42 |
| 56 | Cerebrospinal fluid levels of Alzheimer's disease biomarkers in middle-aged patients with type 1 diabetes. Diabetologia, 2014, 57, 2208-2214. | 2.9 | 40 |
| 57 | Dynamic hub load predicts cognitive decline after resective neurosurgery. Scientific Reports, 2017, 7, 42117. | 1.6 | 39 |
| 58 | Linking late cognitive outcome with glioma surgery location using resection cavity maps. Human Brain Mapping, 2018, 39, 2064-2074. | 1.9 | 38 |
| 59 | Trial design on prophylaxis and treatment of brain metastases: Lessons learned from the EORTC Brain Metastases Strategic Meeting 2012. European Journal of Cancer, 2012, 48, 3439-3447. | 1.3 | 37 |
| 60 | The association between cognitive functioning and health-related quality of life in low-grade glioma patients. Neuro-Oncology Practice, 2014, 1, 40-46. | 1.0 | 37 |
| 61 | Cognition and health-related quality of life in a well-defined subgroup of patients with partial epilepsy. Journal of Neurology, 2002, 249, 294-299. | 1.8 | 36 |
| 62 | How Useful Is the IQCODE for Discriminating between Alzheimer's Disease, Mild Cognitive Impairment and Subjective Memory Complaints?. Dementia and Geriatric Cognitive Disorders, 2010, 30, 411-416. | 0.7 | 35 |
| 63 | Association between tumor location and neurocognitive functioning using tumor localization maps. Journal of Neuro-Oncology, 2019, 144, 573-582. | 1.4 | 35 |
| 64 | Functional Outcomes and Health-Related Quality of Life Following Glioma Surgery. Neurosurgery, 2021, 88, 720-732. | 0.6 | 35 |
| 65 | Altered eigenvector centrality is related to local restingâ€state network functional connectivity in patients with longstanding type 1 diabetes mellitus. Human Brain Mapping, 2017, 38, 3623-3636. | 1.9 | 33 |
| 66 | The Measurement of Cognitive Functioning in Low-Grade Glioma Patients After Radiotherapy. Journal of Clinical Oncology, 2004, 22, 966-967. | 0.8 | 32 |
| 67 | Ventral Striatum, but Not Cortical Volume Loss, Is Related to Cognitive Dysfunction in Type 1 Diabetic Patients With and Without Microangiopathy. Diabetes Care, 2014, 37, 2483-2490. | 4.3 | 31 |
| 68 | Oscillatory brain activity associates with neuroligin-3 expression and predicts progression free survival in patients with diffuse glioma. Journal of Neuro-Oncology, 2018, 140, 403-412. | 1.4 | 31 |
| 69 | The Lesioned Brain: Still a Small-World?. Frontiers in Human Neuroscience, 2010, 4, 174. | 1.0 | 29 |
| 70 | Impact of neurocognitive deficits on patient–proxy agreement regarding health-related quality of life in low-grade glioma patients. Quality of Life Research, 2017, 26, 869-880. | 1.5 | 29 |
| 71 | Functional connectivity in the brain before and during intra-arterial amobarbital injection (Wada) Tj ETQq1 10.78 | 4314 rgBT 2.1 | Overlock 28 |
| 72 | Cognition and health-related quality of life in chronic well-controlled patients with partial epilepsy on carbamazepine monotherapy. Epilepsy and Behavior, 2002, 3, 316-321. | 0.9 | 26 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | Determining priority signs and symptoms for use as clinical outcomes assessments in trials including patients with malignant gliomas: Panel 1 Report. Neuro-Oncology, 2016, 18, ii1-ii12. | 0.6 | 26 |
| 74 | Alleviation of Brain Edema and Restoration of Functional Independence by Bevacizumab in Brain-Metastatic Breast Cancer: A Case Report. Breast Care, 2014, 9, 134-134. | 0.8 | 25 |
| 75 | Disrupted subjectâ€specific gray matter network properties and cognitive dysfunction in type 1 diabetes patients with and without proliferative retinopathy. Human Brain Mapping, 2016, 37, 1194-1208. | 1.9 | 25 |
| 76 | Memory in low-grade glioma patients treated with radiotherapy or temozolomide: a correlative analysis of EORTC study 22033-26033. Neuro-Oncology, 2021, 23, 803-811. | 0.6 | 25 |
| 77 | Evaluation of the Hippocampal Normal Tissue Complication Model in a Prospective Cohort of Low Grade Glioma Patients—An Analysis Within the EORTC 22033 Clinical Trial. Frontiers in Oncology, 2019, 9, 991. | 1.3 | 24 |
| 78 | Attitudes and preferences toward monitoring symptoms, distress, and quality of life in glioma patients and their informal caregivers. Supportive Care in Cancer, 2016, 24, 3011-22. | 1.0 | 23 |
| 79 | Individual changes in neurocognitive functioning and health-related quality of life in patients with brain oligometastases treated with stereotactic radiotherapy. Journal of Neuro-Oncology, 2018, 139, 359-368. | 1.4 | 23 |
| 80 | Processing speed is related to striatal dopamine transporter availability in Parkinson's disease. NeuroImage: Clinical, 2020, 26, 102257. | 1.4 | 22 |
| 81 | Detection of memory impairment in the general population: screening by questionnaire and telephone compared to subsequent face-to-face assessment. International Journal of Geriatric Psychiatry, 2007, 22, 203-210. | 1.3 | 21 |
| 82 | Internet-based guided self-help for glioma patients with depressive symptoms: design of a randomized controlled trial. BMC Neurology, 2014, 14, 81. | 0.8 | 21 |
| 83 | The association between preoperative edema and postoperative cognitive functioning and health-related quality of life in WHO grade I meningioma patients. Acta Neurochirurgica, 2019, 161, 579-588. | 0.9 | 20 |
| 84 | Understanding Global Brain Network Alterations in Glioma Patients. Brain Connectivity, 2021, 11, 865-874. | 0.8 | 20 |
| 85 | Changes in MEG resting-state networks are related to cognitive decline in type 1 diabetes mellitus patients. NeuroImage: Clinical, 2014, 5, 69-76. | 1.4 | 19 |
| 86 | Postoperative oscillatory brain activity as an add-on prognostic marker in diffuse glioma. Journal of Neuro-Oncology, 2020, 147, 49-58. | 1.4 | 19 |
| 87 | Cognitive functioning and functional brain networks in postoperative WHO grade I meningioma patients. Journal of Neuro-Oncology, 2018, 140, 605-613. | 1.4 | 17 |
| 88 | Prevalence of Prodromal Symptoms of Parkinson's Disease in the Late Middle-Aged Population. Journal of Parkinson's Disease, 2022, 12, 967-974. | 1.5 | 17 |
| 89 | Subgenual Cingulate Cortex Functional Connectivity in Relation to Depressive Symptoms and Cognitive Functioning in Type 1 Diabetes Mellitus Patients. Psychosomatic Medicine, 2016, 78, 740-749. | 1.3 | 16 |
| 90 | Lesion momentum as explanation for preoperative neurocognitive function in patients with malignant glioma. Neuro-Oncology, 2016, 18, 1595-1596. | 0.6 | 16 |

| # | Article | IF | CITATIONS |
|-----|--|-----------------|-----------------|
| 91 | Healthcare utilization and productivity loss in glioma patients and family caregivers: the impact of treatable psychological symptoms. Journal of Neuro-Oncology, 2020, 147, 485-494. | 1.4 | 16 |
| 92 | Improving the Accuracy and Precision of Cognitive Testing in Mild Dementia. Journal of the International Neuropsychological Society, 2012, 18, 314-322. | 1.2 | 15 |
| 93 | Profiling cognitive and neuropsychiatric heterogeneity in Parkinson's disease. Parkinsonism and Related Disorders, 2016, 28, 130-136. | 1.1 | 15 |
| 94 | Long-Term Disease Burden and Survivorship Issues After Surgery and Radiotherapy of Intracranial Meningioma Patients. Neurosurgery, 2021, 88, 155-164. | 0.6 | 15 |
| 95 | Implementation of Guidelines for the Treatment of Acute ST-Elevation Myocardial Infarction. Circulation: Cardiovascular Interventions, 2008, 1, 95-102. | 1.4 | 14 |
| 96 | Determinants and predictors for the long-term disease burden of intracranial meningioma patients. Journal of Neuro-Oncology, 2021, 151, 201-210. | 1.4 | 14 |
| 97 | The effectiveness of psychological interventions for patients with relatively well-controlled epilepsy. Epilepsy and Behavior, 2002, 3, 420-426. | 0.9 | 13 |
| 98 | Objective neurocognitive functioning and neurocognitive complaints in patients with high-grade glioma: Evidence of cognitive awareness from the European Organisation for Research and Treatment of Cancer brain tumour clinical trials. European Journal of Cancer, 2021, 144, 162-168. | 1.3 | 12 |
| 99 | Agranulocytosis Secondary to Chlorthalidone Therapy. JAMA - Journal of the American Medical Association, 1963, 184, 310. | 3.8 | 9 |
| 100 | Accelerated executive functions decline and gray matter structural changes in middleâ€aged type 1 diabetes mellitus patients with proliferative retinopathy. Journal of Diabetes, 2018, 10, 835-846. | 0.8 | 9 |
| 101 | The presence of cerebral white matter lesions and lower skin microvascular perfusion predicts lower cognitive performance in type 1 diabetes patients with retinopathy but not in healthy controls—A longitudinal study. Microcirculation, 2019, 26, e12530. | 1.0 | 8 |
| 102 | Development of Randomized Trials in Adults with Medulloblastoma—The Example of EORTC 1634-BTG/NOA-23. Cancers, 2021, 13, 3451. | 1.7 | 8 |
| 103 | Subjective Ratings vs. Objective Measurement of Cognitive Function: In Regard to Van Beek etÂal. (Int J) Tj ETQq1 2008, 70, 961-962. | 1 0.7843 0.4 | 14 rgBT /O 7 |
| 104 | Longitudinal proxy measurements in multiple sclerosis: patient-proxy agreement on the impact of MS on daily life over a period of two years. BMC Neurology, 2008, 8, 2. | 0.8 | 7 |
| 105 | The size of the treatment effect: do patients and proxies agree?. BMC Neurology, 2009, 9, 12. | 0.8 | 7 |
| 106 | Improved, personalized treatment of glioma necessitates long-term follow-up of cognitive functioning. Pharmacogenomics, 2012, 13, 1667-1669. | 0.6 | 7 |
| 107 | CSF Biomarkers Reflecting Protein Pathology and Axonal Degeneration Are Associated with Memory, Attentional, and Executive Functioning in Early-Stage Parkinson′s Disease. International Journal of Molecular Sciences, 2020, 21, 8519. | 1.8 | 7 |
| 108 | Corticosteroids use and neurocognitive functioning in patients with recurrent glioblastoma: Evidence from European Organization for Research and Treatment of Cancer (EORTC) trial 26101. Neuro-Oncology Practice, 2022, 9, 310-316. | 1.0 | 7 |

| # | Article | IF | CITATIONS |
|-----|---|-------------------|-------------|
| 109 | Differential impact of subclinical carotid artery disease on cerebral structure and functioning in type 1 diabetes patients with versus those without proliferative retinopathy. Cardiovascular Diabetology, 2014, 13, 58. | 2.7 | 6 |
| 110 | The level of reporting of neurocognitive outcomes in randomised controlled trials of brain tumour patients: AÂsystematic review. European Journal of Cancer, 2018, 100, 104-125. | 1.3 | 6 |
| 111 | Cellular Substrates of Functional Network Integration and Memory in Temporal Lobe Epilepsy. Cerebral Cortex, 2022, 32, 2424-2436. | 1.6 | 6 |
| 112 | The long-term caregiver burden in World Health Organization grade I and II meningioma: It is not just the patient. Neuro-Oncology Advances, 2021, 3, vdaa169. | 0.4 | 6 |
| 113 | Reducing severe fatigue in patients with diffuse glioma: a study protocol for an RCT on the effect of blended cognitive behavioural therapy. Trials, 2022, 23, . | 0.7 | 6 |
| 114 | Evaluation of cognitive functions and quality of life. Handbook of Clinical Neurology / Edited By P J Vinken and G W Bruyn, 2012, 104, 173-183. | 1.0 | 5 |
| 115 | NCOG-07. MEMORY FUNCTIONING IN LOW-GRADE GLIOMA PATIENTS TREATED WITH EITHER RADIOTHERAPY (RT) OR TEMOZOLOMIDE (TMZ) CHEMOTHERAPY. AÂCORRELATIVE ANALYSIS OF EUROPEAN ORGANIZATION FOR RESEARCH AND TREATMENT (EORTC) STUDY 22033-26033. Neuro-Oncology, 2017, 19, vi139-vi139. | 0.6 | 5 |
| 116 | Development of an EORTC questionnaire measuring instrumental activities of daily living (IADL) in patients with brain tumours: phase l–III. Quality of Life Research, 2021, 30, 1491-1502. | 1.5 | 5 |
| 117 | Impaired Set-Shifting from Dorsal Stream Disconnection: Insights from a European Series of Right Parietal Lower-Grade Glioma Resection. Cancers, 2021, 13, 3337. | 1.7 | 5 |
| 118 | Long-term impact of adult WHO grade II or III gliomas on health-related quality of life: A systematic review. Neuro-Oncology Practice, 2022, 9, 3-17. | 1.0 | 5 |
| 119 | ls poor sleep quality associated with poor neurocognitive outcome in cancer survivors? A systematic review. Journal of Cancer Survivorship, 2022, , 1. | 1.5 | 5 |
| 120 | A Web-Based Lifestyle Intervention Aimed at Improving Cognition in Patients With Cancer Returning to Work in an Outpatient Setting: Protocol for a Randomized Controlled Trial. JMIR Research Protocols, 2021, 10, e22670. | 0.5 | 3 |
| 121 | Cognitive Functioning and Hippocampal Connectivity in Patients With Longstanding Type 1 Diabetes and Apolipoprotein E ε4. Diabetes Care, 2021, 44, 2388-2396. | 4.3 | 3 |
| 122 | CTNI-29. CODEL: PHASE III TRIAL OF RT ALONE, RT PLUS TMZ, OR TMZ ALONE FOR NEWLY-DIAGNOSED, 1p/19q CODELETED ANAPLASTIC OLIGODENDROGLIOMA. ANALYSIS FROM THE INITIAL STUDY DESIGN. (NCCTG N0577,) | T∳ <u>I</u> ETQq0 | 030 rgBT /0 |
| 123 | Prognostic Significance of DNA Methylation Profiles at MRI Enhancing Tumor Recurrence: a Report from the EORTC 26091 TAVAREC Trial. Clinical Cancer Research, 2022, 28, 2440-2448. | 3.2 | 3 |
| 124 | In Reference to Maschio et al. (Neuro-Oncology. 2008;10:106–107). Neuro-Oncology, 2008, 10, 1172-1172. | 0.6 | 2 |
| 125 | An Informant Questionnaire for Detecting Alzheimer's Disease: Are Some Items Better Than Others?. Journal of the International Neuropsychological Society, 2011, 17, 674-681. | 1.2 | 2 |
| | | | |

126 Quality of Life in Patients with Diffuse Low-Grade Glioma. , 2017, , 235-252.

2

| # | Article | IF | CITATIONS |
|-----|--|-------------------------------|-----------|
| 127 | Title is missing!. Epilepsy and Behavior, 2003, 4, 91. | 0.9 | 1 |
| 128 | MNGI-26. THE DISEASE BURDEN OF MENINGIOMA PATIENTS: LONG-TERM RESULTS ON WORK PRODUCTIVITY AND HEALTHCARE CONSUMPTION. Neuro-Oncology, 2018, 20, vi154-vi154. | 0.6 | 1 |
| 129 | Immunobiological and Neural Substrates of Cancer-Related Neurocognitive Deficits. , 2009, , 327-340. | | 1 |
| 130 | Do neurocognitive impairments explain the differences between brain tumor patients and their proxies when assessing the patient's IADL?. Neuro-Oncology Practice, 2022, 9, 271-283. | 1.0 | 1 |
| 131 | PS1 - 6. Cerebral resting-state network changes in patients with type 1 diabetes with and without microangiopathy relate to cognitive functions. Nederlands Tijdschrift Voor Diabetologie, 2011, 9, 94-94. | 0.0 | 0 |
| 132 | PS11 - 50. Depressive symptoms exacerbate the negative effects of type 1 diabetes and microangiopathy on brain functioning. Nederlands Tijdschrift Voor Diabetologie, 2012, 10, 133-133. | 0.0 | 0 |
| 133 | ACTR-18. MOLECULAR GENETIC, HOST-DERIVED AND CLINICAL DETERMINANTS OF LONG-TERM SURVIVAL IN GLIOBLASTOMA: AN UPDATE FROM THE BRAIN TUMOR FUNDERS' COLLABORATIVE CONSORTIUM. Neuro-Oncology, 2016, 18, vi5-vi5. | 0.6 | 0 |
| 134 | MNGI-25. THE CAREGIVER BURDEN IN MENINGIOMA: LONG-TERM RESULTS AND ITS EFFECTS ON CAREGIVER' HEALTH-RELATED QUALITY OF LIFE, ANXIETY AND DEPRESSION. Neuro-Oncology, 2018, 20, vi154-vi154. | ¹ S _{0.6} | 0 |
| 135 | MNGI-27. THE LONG-TERM DISEASE BURDEN OF MENINGIOMA PATIENTS: RESULTS ON HEALTH-RELATED QUALITY OF LIFE, COGNITIVE FUNCTION, ANXIETY AND DEPRESSION. Neuro-Oncology, 2018, 20, vi154-vi155. | 0.6 | 0 |
| 136 | NCOG-09. THE LEVEL OF REPORTING OF NEUROCOGNITIVE OUTCOMES IN RANDOMIZED CONTROLLED TRIALS OF BRAIN TUMOR PATIENTS: A SYSTEMATIC REVIEW. Neuro-Oncology, 2018, 20, vi174-vi174. | 0.6 | 0 |
| 137 | Assessment of Neurocognitive Functioning in Clinical Practice and for Trial Purposes. , 2019, , 121-131. | | 0 |
| 138 | In Reply: Functional Outcomes and Health-Related Quality of Life Following Glioma Surgery. Neurosurgery, 2021, 89, E189-E189. | 0.6 | 0 |
| 139 | QOLP-14. LONG-TERM IMPACT OF ADULT GLIOMA ON HEALTH-RELATED QUALITY OF LIFE: A SYSTEMATIC REVIEW. Neuro-Oncology, 2021, 23, vi185-vi186. | 0.6 | 0 |
| 140 | QOLP-05. HEALTH-RELATED QUALITY OF LIFE IN LOW-GRADE GLIOMA SURVIVORS 26 YEARS AFTER DIAGNOSIS. Neuro-Oncology, 2021, 23, vi183-vi183. | 0.6 | 0 |
| 141 | NCOG-49. UNMET NEEDS AND WISH FOR SUPPORT OF FAMILY CAREGIVERS OF PRIMARY BRAIN TUMOUR PATIENTS. Neuro-Oncology, 2021, 23, vi162-vi163. | 0.6 | 0 |
| 142 | Slowing as a multidomain and vascular geriatric syndrome: Apathy symptoms, gait speed and information processing speed in a geriatric memory clinic population. Alzheimer's and Dementia, 2021, 17, . | 0.4 | 0 |