Vladislav Yu Kuznetsov

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2655280/publications.pdf

Version: 2024-02-01

42 papers

554 citations

623734 14 h-index 22 g-index

43 all docs 43 docs citations

43 times ranked

607 citing authors

| # | Article | IF | CITATIONS |
|----|---|-----------|------------|
| 1 | Seafloor Massive Sulfides from the Northern Equatorial Mid-Atlantic Ridge: New Discoveries and Perspectives. Marine Georesources and Geotechnology, 2010, 28, 222-239. | 2.1 | 62 |
| 2 | Composition and characteristics of the ferromanganese crusts from the western Arctic Ocean. Ore Geology Reviews, 2017, 87, 88-99. | 2.7 | 43 |
| 3 | Massive sulfide ores of the northern equatorial Mid-Atlantic Ridge. Oceanology, 2013, 53, 607-619. | 1.2 | 42 |
| 4 | Ice Complex permafrost of MIS5 age in the Dmitry Laptev Strait coastal region (East Siberian Arctic). Quaternary Science Reviews, 2016, 147, 298-311. | 3.0 | 37 |
| 5 | Sulfide geochronology along the Northern Equatorial Mid-Atlantic Ridge. Ore Geology Reviews, 2017, 87, 147-154. | 2.7 | 37 |
| 6 | Two New Hydrothermal Fields at the Mid-Atlantic Ridge. Marine Georesources and Geotechnology, 2008, 26, 308-316. | 2.1 | 34 |
| 7 | 230Th/U chronology of ore formation within the semyenov hydrothermal district (13°31′ N) at the Mid-Atlantic ridge. Geochronometria, 2011, 38, 72-76. | 0.8 | 26 |
| 8 | Ice Complex formation on Bol'shoy Lyakhovsky Island (New Siberian Archipelago, East Siberian Arctic) since about 200 ka. Quaternary Research, 2019, 92, 530-548. | 1.7 | 26 |
| 9 | Environmental and climate reconstructions of the Fore-Baikal area during MIS 5-1: Multiproxy record from terrestrial sediments of the Ust-Oda section (Siberia, Russia). Journal of Asian Earth Sciences, 2016, 129, 220-230. | 2.3 | 20 |
| 10 | Landscape evolution in the periglacial zone of Eastern Europe since MIS5: Proxies from paleosols and sediments of the Cheremoshnik key site (Upper Volga, Russia). Quaternary International, 2015, 365, 26-41. | 1.5 | 18 |
| 11 | New hydrothermal ore fields in the Mid-Atlantic Ridge: Zenith-Victoria (20°08′ N) and Petersburg (19°52′ |) ŢįĘTQq1 | 1,0.784314 |
| 12 | Palaeoecological investigations and 230Th/U dating of Eemian interglacial peat sequence of Banzin (Mecklenburg-Western Pomerania, NE-Germany). Quaternary International, 2015, 386, 122-136. | 1.5 | 17 |
| 13 | Late Pleistocene paleosols in the extra-glacial regions of Northwestern Eurasia: Pedogenesis, post-pedogenic transformation, paleoenvironmental inferences. Quaternary International, 2019, 501, 174-192. | 1.5 | 17 |
| 14 | Late Quaternary marine terraces in the Mediterranean coastal area of Syria: Geochronology and neotectonics. Quaternary International, 2008, 190, 158-170. | 1.5 | 15 |
| 15 | Palaeoecological investigations and 230Th/U dating of the Eemian Interglacial peat sequence from Neubrandenburg-Hinterste Mýhle (Mecklenburg-Western Pomerania, NE Germany). Quaternary International, 2018, 467, 62-78. | 1.5 | 15 |
| 16 | The 230Th/U dating of sulfide ores in the ocean: Methodical possibilities, measurement results, and perspectives of application. Doklady Earth Sciences, 2007, 417, 1202-1205. | 0.7 | 14 |
| 17 | The oldest seafloor massive sulfide deposits at the Mid-Atlantic Ridge: ²³⁰ Th/U chronology and composition. Geochronometria, 2015, 42, . | 0.8 | 12 |
| 18 | First 230Th/U date of Middle Pleistocene peat bog in Siberia (key section Krivosheino, Western Siberia). Geochronometria, 2012, 39, 241-251. | 0.8 | 11 |

| # | Article | IF | Citations |
|----|--|-----|-----------|
| 19 | The first case study of 230Th/U and 14C dating of mid-valdai organic deposits. Doklady Earth Sciences, 2011, 438, 598-602. | 0.7 | 10 |
| 20 | Stratigraphy of bottom sediments in the Mendeleev Ridge area (Arctic Ocean). Doklady Earth Sciences, 2013, 450, 602-606. | 0.7 | 10 |
| 21 | Assessment of the long-term safety of radioactive waste disposal: 1. Paleoreconstruction of groundwater formation conditions. Water Resources, 2009, 36, 206-213. | 0.9 | 9 |
| 22 | The terrestrial Eemian to late Weichselian sediment record at Beckentin (NE-Germany): First results from lithostratigraphic, palynological and geochronological analyses. Quaternary International, 2019, 501, 90-108. | 1.5 | 9 |
| 23 | Geochronology and landscape-climatic environments of the Early Zyryanian Interstadial in West Siberia. Doklady Earth Sciences, 2008, 421, 796-799. | 0.7 | 7 |
| 24 | A new approach to isotope dating of buried organic-rich deposits with an example from the Kuryador section, upper Vychegda valley. Doklady Earth Sciences, 2015, 462, 570-574. | 0.7 | 7 |
| 25 | Mass-wasting processes input in proximal metalliferous sediments: A case study of the Pobeda hydrothermal fields. Marine Geology, 2021, 438, 106517. | 2.1 | 7 |
| 26 | Last interglacial climate changes and environments of the Lesser Kuril arc, north-western Pacific. Quaternary International, 2011, 241, 35-50. | 1.5 | 6 |
| 27 | Paleoclimates and chronology of the middle $W\tilde{A}^{1}\!\!/\!\!4$ rm megainterstadial on the West Siberian Plain. Doklady Earth Sciences, 2006, 411, 1457-1461. | 0.7 | 5 |
| 28 | New outcrop of buried Kazantsevo peat at lower reaches of the Irtysh River. Doklady Earth Sciences, 2008, 419, 200-204. | 0.7 | 5 |
| 29 | Geochronology and Palaeomagnetic Records of the SnaigupÄ—lÄ— Section in South Lithuania. Geochronometria, 2015, 42, . | 0.8 | 4 |
| 30 | The first uranium-thorium dating of the Middle Neopleistocene peat in West Siberia. Doklady Earth Sciences, 2010, 433, 915-919. | 0.7 | 3 |
| 31 | The first find of buried low-temperature hydrothermal deposits in the Mid-Atlantic Ridge rift valley. Doklady Earth Sciences, 2009, 424, 1-6. | 0.7 | 2 |
| 32 | U–Th age of the Kazantsevo (MIS 5) Horizon of the Upper Neopleistocene Ust Oda reference section, Baikal Region. Doklady Earth Sciences, 2017, 473, 266-270. | 0.7 | 2 |
| 33 | Chronostratigraphy of the Cheremoshnik key section (Yaroslavl Volga region) based on new geochronological, palynological, and paleosol data. Doklady Earth Sciences, 2017, 472, 244-247. | 0.7 | 2 |
| 34 | Origin of high 234U/238U ratio in post-permafrost aquifers. , 2006, , 847-856. | | 2 |
| 35 | Last interglacial environment of the Baikal Region (Southern Siberia, Russia) based on analysis of fossil invertebrates and plants. Palaeoentomology, 2021, 4, . | 1.0 | 1 |
| 36 | Middle Pleistocene warming phase based on the deposits of a buried oyster reef, Southern Lesser Kuril Islands. Doklady Earth Sciences, 2014, 455, 376-382. | 0.7 | 0 |

| # | Article | IF | CITATIONS |
|----|---|-------------------|-----------------------------|
| 37 | Comparative ²³⁰ Th/U and ¹⁴ C Dating of a Buried Stump Layer (Western) Tj ETQq1 1 C |).784314 r 0.8 | gBT /Over <mark>l</mark> oc |
| 38 | Environmental changes at final warming of Middle Pleistocene (MIS 7) in South Kurils. Quaternary International, 2015, 355, 90-100. | 1.5 | 0 |
| 39 | Climatic Stratigraphy of the Kazantsevo Horizon (as an Analogue of MIS-5) in the Boreal Zone of Western Siberia. Springer Geology, 2014, , 965-968. | 0.3 | O |
| 40 | The First Case Study of 230Th–U Dating of Buried Wood Remnants from Siberia. Springer Geology, 2014, , 293-296. | 0.3 | 0 |
| 41 | Uranium-thorium dating of high sea terraces of the Spitsbergen Archipelago. Vestnik of Saint Petersburg University Geology Geography, 2016, , 54-64. | 0.0 | 0 |
| 42 | POLLEN COMPLEXES OF THE MIKULINO (EEMIAN) INTERGLACIAL'S INITIAL PHASE IN THE UPPER VOLGA BASIN (ACCORDING TO THE STUDY OF THE MALAYA KOSHA RIVER SECTION). , 2022, , . | | 0 |