

Jian-Yi Li

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2661243/publications.pdf>

Version: 2024-02-01

18
papers

304
citations

1040056

9
h-index

940533

16
g-index

19
all docs

19
docs citations

19
times ranked

500
citing authors

#	ARTICLE	IF	CITATIONS
1	The heterogeneous subclones might be induced by cycling hypoxia which was aggravated along with the luminal A tumor growth. <i>Tissue and Cell</i> , 2022, , 101844.	2.2	0
2	Long-term tracing and staining of carbon nanoparticles for axillary lymph node in patients with local advanced breast cancer treated with neoadjuvant chemotherapy. <i>Asian Journal of Surgery</i> , 2021, , .	0.4	3
3	A Novel Model Incorporating Tumor Stiffness, Blood Flow Characteristics, and Ki-67 Expression to Predict Responses After Neoadjuvant Chemotherapy in Breast Cancer. <i>Frontiers in Oncology</i> , 2020, 10, 603574.	2.8	5
4	Efficacy of shear-wave elastography versus dynamic optical breast imaging for predicting the pathological response to neoadjuvant chemotherapy in breast cancer. <i>European Journal of Radiology</i> , 2020, 129, 109098.	2.6	10
5	Comparison of strain and shear-wave ultrasonographic elastography in predicting the pathological response to neoadjuvant chemotherapy in breast cancers. <i>European Radiology</i> , 2017, 27, 2282-2291.	4.5	43
6	Combination of shear wave elastography and Ki-67 index as a novel predictive modality for the pathological response to neoadjuvant chemotherapy in patients with invasive breast cancer. <i>European Journal of Cancer</i> , 2016, 69, 86-101.	2.8	42
7	Partial axillary lymph node dissection inferior to the intercostobrachial nerves complements sentinel node biopsy in patients with clinically node-negative breast cancer. <i>BMC Surgery</i> , 2015, 15, 79.	1.3	17
8	Contrast enhanced computed tomography is indicative for angiogenesis pattern and display prognostic significance in breast cancer. <i>BMC Cancer</i> , 2014, 14, 672.	2.6	9
9	A New Technique That Complements Sentinel Lymph Node Biopsy: Lymph Node Dissection Under the Intercostobrachial Nerves in Early-Stage Breast Cancer. <i>Clinical Breast Cancer</i> , 2013, 13, 212-218.	2.4	3
10	Intercostobrachial Nerves as a Novel Anatomic Landmark for Dividing the Axillary Space in Lymph Node Dissection. <i>ISRN Oncology</i> , 2013, 2013, 1-7.	2.1	7
11	Survival Analysis Based on Clinicopathological Data from a Single Institution: Chemotherapy Intensity Would Be Enhanced in Patients with Positive Hormone Receptors and Positive HER2 in China Who Cannot Afford the Target Therapy. <i>ISRN Oncology</i> , 2013, 2013, 1-8.	2.1	3
12	Genetic Heterogeneity of Breast Cancer Metastasis May Be Related to miR-21 Regulation of TIMP-3 in Translation. <i>International Journal of Surgical Oncology</i> , 2013, 2013, 1-7.	0.6	28
13	Effects of Differential Distribution of Microvessel Density, Possibly Regulated by miR-374a, on Breast Cancer Prognosis. <i>Asian Pacific Journal of Cancer Prevention</i> , 2013, 14, 1715-1720.	1.2	21
14	Differential Distribution of microRNAs in Breast Cancer Grouped by Clinicopathological Subtypes. <i>Asian Pacific Journal of Cancer Prevention</i> , 2013, 14, 3197-3203.	1.2	27
15	Differential Distribution of miR-20a and miR-20b may Underly Metastatic Heterogeneity of Breast Cancers. <i>Asian Pacific Journal of Cancer Prevention</i> , 2012, 13, 1901-1906.	1.2	40
16	Melatonin protects liver from intestine ischemia reperfusion injury in rats. <i>World Journal of Gastroenterology</i> , 2008, 14, 7392.	3.3	18
17	Melatonin abates liver ischemia/reperfusion injury by improving the balance between nitric oxide and endothelin. <i>Hepatobiliary and Pancreatic Diseases International</i> , 2006, 5, 574-9.	1.3	28
18	Estrogen Receptor and Claudin-6 Might Play Vital Roles for Long-Term Prognosis in Patients With Luminal A Breast Cancer Who Underwent Neoadjuvant Chemotherapy. <i>Frontiers in Oncology</i> , 0, 12, .	2.8	0