

# Houping Liu

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2665915/publications.pdf>

Version: 2024-02-01

14

papers

404

citations

1040056

9

h-index

1058476

14

g-index

14

all docs

14

docs citations

14

times ranked

246

citing authors

#	ARTICLE	IF	CITATIONS
1	Infrared thermography for insect detection: lighting up the spotted lanternfly in the field. <i>Journal of Pest Science</i> , 2021, 94, 231-240.	3.7	3
2	Nondestructive Sampling for Spotted Lanternfly (Hemiptera: Fulgoridae) Egg Masses in Woodlands Based on Fixed-Radius Plots. <i>Journal of Economic Entomology</i> , 2021, 114, 1353-1361.	1.8	3
3	Biology and ecology of the Northern walkingstick, <i>&lt; i&gt;Diapheromera femorata&lt;/i&gt;</i> (Say) (Phasmatodea). <i>Tj ETQq1 1.0.784314 rgBT /Overlock 10</i>	1.8	6
4	Seasonal Development, Cumulative Growing Degree-Days, and Population Density of Spotted Lanternfly (Hemiptera: Fulgoridae) on Selected Hosts and Substrates. <i>Environmental Entomology</i> , 2020, 49, 1171-1184.	1.4	21
5	Spatial Distribution of <i>Lycorma delicatula</i> (Hemiptera: Fulgoridae) Egg Masses on Tree-of-Heaven, Black Walnut, and Siberian Elm in North America. <i>Journal of Economic Entomology</i> , 2020, 113, 1028-1032.	1.8	16
6	Oviposition Substrate Selection, Egg Mass Characteristics, Host Preference, and Life History of the Spotted Lanternfly (Hemiptera: Fulgoridae) in North America. <i>Environmental Entomology</i> , 2019, 48, 1452-1468.	1.4	56
7	Occurrence, Seasonal Abundance, and Superparasitism of <i>Ooencyrtus kuvanae</i> (Hymenoptera: Encyrtidae). <i>Tj ETQq1 1.0.784314 rgBT /Overlock 10</i>	2.1	18
8	Forests, 2019, 10, 79.		
9	Under Siege: Ash Management in the Wake of the Emerald Ash Borer. <i>Journal of Integrated Pest Management</i> , 2018, 9, .	2.0	8
10	An Old Remedy for a New Problem? Identification of <i>Ooencyrtus kuvanae</i> (Hymenoptera: Encyrtidae). <i>Tj ETQq1 1.0.784314 rgBT /Overlock 10</i>	1.5	30
11	Journal of Insect Science, 2017, 17, 18.		
12	Seasonal abundance and development of the Asian longhorned beetle and natural enemy prevalence in different forest types in China. <i>Biological Control</i> , 2016, 103, 154-164.	3.0	8
13	Assessing deposition and persistence of <i>Beauveria bassiana</i> GHA (Ascomycota: Hypocreales) applied for control of the emerald ash borer, <i>Agrilus planipennis</i> (Coleoptera: Buprestidae), in a commercial tree nursery. <i>Biological Control</i> , 2010, 54, 61-67.	3.0	32
14	Characterization of <i>Beauveria bassiana</i> (Ascomycota: Hypocreales) isolates associated with <i>Agrilus planipennis</i> (Coleoptera: Buprestidae) populations in Michigan. <i>Biological Control</i> , 2010, 54, 135-140.	3.0	28
15	Microbial control of emerald ash borer, <i>Agrilus planipennis</i> (Coleoptera: Buprestidae) with <i>Beauveria bassiana</i> strain GHA: Greenhouse and field trials. <i>Biological Control</i> , 2008, 45, 124-132.	3.0	42
16	Seasonal abundance of <i>Agrilus planipennis</i> (Coleoptera: Buprestidae) and its natural enemies <i>Oobius agrili</i> (Hymenoptera: Encyrtidae) and <i>Tetrastichus planipennisi</i> (Hymenoptera: Eulophidae) in China. <i>Biological Control</i> , 2007, 42, 61-71.	3.0	133