Agnieszka Adamczyk

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2666372/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Comparison of eight commercially available kits for DNA extraction from formalin-fixed paraffin-embedded tissues. Analytical Biochemistry, 2015, 476, 8-10.	2.4	43
2	Original paper Prognostic value of PIK3CA mutation status, PTEN and androgen receptor expression for metastasis-free survival in HER2-positive breast cancer patients treated with trastuzumab in adjuvant setting. Polish Journal of Pathology, 2015, 2, 133-141.	0.3	25
3	CD44/CD24 as potential prognostic markers in node-positive invasive ductal breast cancer patients treated with adjuvant chemotherapy. Journal of Molecular Histology, 2014, 45, 35-45.	2.2	24
4	Relationship between HER2 gene status and selected potential biological features related to trastuzumab resistance and its influence on survival of breast cancer patients undergoing trastuzumab adjuvant treatment. OncoTargets and Therapy, 2018, Volume 11, 4525-4535.	2.0	21
5	Prognostic Significance of Intratumour Microvessel Density and Haemoglobin Level in Carcinoma of the Uterine Cervix. Acta Oncológica, 2002, 41, 437-443.	1.8	19
6	Tumor Grade and Matrix Metalloproteinase 2 Expression in Stromal Fibroblasts Help to Stratify the High-Risk Group of Patients With Early Breast Cancer Identified on the Basis of St Gallen Recommendations. Clinical Breast Cancer, 2013, 13, 119-128.	2.4	17
7	Proteins Involved in HER2 Signalling Pathway, Their Relations and Influence on Metastasis-Free Survival in HER2-Positive Breast Cancer Patients Treated with Trastuzumab in Adjuvant Setting. Journal of Cancer, 2017, 8, 131-139.	2.5	16
8	Relationships Between Immunophenotype, Ki-67 Index, Microvascular Density, Ep-CAM/P-cadherin, and MMP-2 Expression in Early-stage Invasive Ductal Breast Cancer. Applied Immunohistochemistry and Molecular Morphology, 2012, 20, 550-560.	1.2	15
9	Cardiotoxicity Induced by Protein Kinase Inhibitors in Patients with Cancer. International Journal of Molecular Sciences, 2022, 23, 2815.	4.1	15
10	Triple-negative, Basal Marker-expressing, and High-grade Breast Carcinomas are Characterized by High Lymphatic Vessel Density and the Expression of Podoplanin in Stromal Fibroblasts. Applied Immunohistochemistry and Molecular Morphology, 2014, 22, 10-16.	1.2	14
11	Bromodeoxyuridine Labeling Index as an Indicator of Early Tumor Response to Preoperative Radiotherapy in Patients with Rectal Cancer. Journal of Gastrointestinal Surgery, 2007, 11, 520-528.	1.7	13
12	Expression of ER/PR/HER2, basal markers and adhesion molecules in primary breast cancer and in lymph nodes metastases: a comparative immunohistochemical analysis. Polish Journal of Pathology, 2012, 4, 228-234.	0.3	13
13	HPV16 detection by qPCR method in relation to quantity and quality of DNA extracted from archival formalin fixed and paraffin embedded head and neck cancer tissues by three commercially available kits. Journal of Virological Methods, 2016, 236, 157-163.	2.1	12
14	Survival of breast cancer patients according to changes in expression of selected markers between primary tumor and lymph node metastases. Biomarkers in Medicine, 2016, 10, 219-228.	1.4	10
15	Gender-Related Differences in Pathological and Clinical Tumor Response Based on Immunohistochemical Proteins Expression in Rectal Cancer Patients Treated with Short Course of Preoperative Radiotherapy. Journal of Gastrointestinal Surgery, 2014, 18, 1306-1318.	1.7	9
16	Androgen receptor in male breast cancer. Polish Journal of Pathology, 2015, 4, 347-352.	0.3	8
17	Intensity and Pattern of Enhancement on CESM: Prognostic Significance and its Relation to Expression of Podoplanin in Tumor Stroma – A Preliminary Report. Anticancer Research, 2018, 38, 1085-1095.	1.1	8
18	Lymphangiogenesis assessed using three methods is related to tumour grade, breast cancer subtype and expression of basal marker. Polish Journal of Pathology, 2012, 3, 165-171.	0.3	6

#	Article	IF	CITATIONS
19	Gender-related prognostic significance of clinical and biological tumor features in rectal cancer patients receiving short-course preoperative radiotherapy. Reports of Practical Oncology and Radiotherapy, 2017, 22, 368-377.	0.6	6
20	Prognostic role of lymphatic vessel density and lymphovascular invasion in chemotherapy-naive and chemotherapy-treated patients with invasive breast cancer. American Journal of Translational Research (discontinued), 2017, 9, 1435-1447.	0.0	6
21	Synovial sarcoma of the stomach: case report and systematic review of the literature. Polish Journal of Pathology, 2020, 71, 181-193.	0.3	5
22	Comparison of mutation profile between primary phyllodes tumors of the breast and their paired local recurrences. Polish Journal of Pathology, 2020, 71, 7-12.	0.3	5
23	Podoplanin-positive Cancer-associated Stromal Fibroblasts in Primary Tumor and Synchronous Lymph Node Metastases of HER2-overexpressing Breast Carcinomas. Anticancer Research, 2018, 38, 1957-1965.	1.1	5
24	BCX–Ki-67 Index as a Supplementary Marker to MIB-1 Index, Enabling More Precise Distinction Between Luminal A and B Subtypes of Breast Carcinoma and Eliminating the Problem of Membranous/Cytoplasmic MIB-1 Staining. American Journal of Clinical Pathology, 2015, 143, 419-429.	0.7	3
25	The tumour border on contrast-enhanced spectral mammography and its relation to histological characteristics of invasive breast cancer. Polish Journal of Pathology, 2016, 3, 295-299.	0.3	3
26	Comet assay is not useful to predict normal tissue response after radiochemotherapy in cervical and larynx cancer patients. Polish Journal of Pathology, 2018, 69, 410-421.	0.3	3
27	Low frequency of HPV positivity in breast tumors among patients from south-central Poland. Infectious Agents and Cancer, 2021, 16, 67.	2.6	3
28	Distribution of Podoplanin-Positive Tumor Vessels Predicts Disease-Specific Survival of Node-Positive Breast Cancer Patients Treated with Anthracyclines and/or Taxanes. Cancer Investigation, 2014, 32, 168-177.	1.3	2
29	Gender-related significance of time interval between radiotherapy and surgery in hypofractionated preoperative radiotherapy for rectal cancer patients' survival. Reports of Practical Oncology and Radiotherapy, 2016, 21, 174-180.	0.6	1
30	High Ki-67 expression is a marker of poor survival in apocrine breast carcinoma. Polish Journal of Pathology, 2020, 71, 107-119.	0.3	1