

Byeong Mo Kim

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/2673482/byeong-mo-kim-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

32
papers

1,283
citations

19
h-index

35
g-index

35
ext. papers

1,524
ext. citations

6.3
avg, IF

4.33
L-index

#	Paper	IF	Citations
32	Thymoquinone Selectively Kills Hypoxic Renal Cancer Cells by Suppressing HIF-1 β -Mediated Glycolysis. <i>International Journal of Molecular Sciences</i> , 2019 , 20,	6.3	19
31	A negative feedback loop between XBP1 and Fbw7 regulates cancer development. <i>Oncogenesis</i> , 2019 , 8, 12	6.6	9
30	Therapeutic advantage of genetically engineered <i>Salmonella typhimurium</i> carrying short hairpin RNA against inhibin alpha subunit in cancer treatment. <i>Annals of Oncology</i> , 2018 , 29, 2010-2017	10.3	16
29	TNF- β -Induced YAP/TAZ Activity Mediates Leukocyte-Endothelial Adhesion by Regulating VCAM1 Expression in Endothelial Cells. <i>International Journal of Molecular Sciences</i> , 2018 , 19,	6.3	27
28	Emodin Sensitizes Hepatocellular Carcinoma Cells to the Anti-Cancer Effect of Sorafenib through Suppression of Cholesterol Metabolism. <i>International Journal of Molecular Sciences</i> , 2018 , 19,	6.3	43
27	The Role of Saikosaponins in Therapeutic Strategies for Age-Related Diseases. <i>Oxidative Medicine and Cellular Longevity</i> , 2018 , 2018, 8275256	6.7	15
26	Death-associated protein kinase 1 phosphorylates NDRG2 and induces neuronal cell death. <i>Cell Death and Differentiation</i> , 2017 , 24, 238-250	12.7	30
25	Fascaplysin Sensitizes Anti-Cancer Effects of Drugs Targeting AKT and AMPK. <i>Molecules</i> , 2017 , 23,	4.8	14
24	Application of genetically engineered <i>Salmonella typhimurium</i> for interferon-gamma-induced therapy against melanoma. <i>European Journal of Cancer</i> , 2017 , 70, 48-61	7.5	44
23	Vanillin Suppresses Cell Motility by Inhibiting STAT3-Mediated HIF-1 β mRNA Expression in Malignant Melanoma Cells. <i>International Journal of Molecular Sciences</i> , 2017 , 18,	6.3	19
22	Caspase-4 is essential for saikosaponin a-induced apoptosis acting upstream of caspase-2 and β H2AX in colon cancer cells. <i>Oncotarget</i> , 2017 , 8, 100433-100448	3.3	20
21	Role of Protein Kinases and Their Inhibitors in Radiation Response of Tumor Cells. <i>Current Pharmaceutical Design</i> , 2017 , 23, 4259-4280	3.3	3
20	Inhibition of death-associated protein kinase 1 attenuates the phosphorylation and amyloidogenic processing of amyloid precursor protein. <i>Human Molecular Genetics</i> , 2016 , 25, 2498-2513	5.6	23
19	Inhibition of glutamine utilization sensitizes lung cancer cells to apigenin-induced apoptosis resulting from metabolic and oxidative stress. <i>International Journal of Oncology</i> , 2016 , 48, 399-408	4.4	50
18	Small-molecule inhibitors of USP7 induce apoptosis through oxidative and endoplasmic reticulum stress in cancer cells. <i>Biochemical and Biophysical Research Communications</i> , 2016 , 470, 181-186	3.4	23
17	Antiobesity and Antidiabetes Effects of a <i>Cudrania tricuspidata</i> Hydrophilic Extract Presenting PTP1B Inhibitory Potential. <i>BioMed Research International</i> , 2016 , 2016, 8432759	3	20
16	A potential therapeutic effect of saikosaponin C as a novel dual-target anti-Alzheimer agent. <i>Journal of Neurochemistry</i> , 2016 , 136, 1232-1245	6	22

15	Therapeutic Implications for Overcoming Radiation Resistance in Cancer Therapy. <i>International Journal of Molecular Sciences</i> , 2015 , 16, 26880-913	6.3	110
14	Pin1 cysteine-113 oxidation inhibits its catalytic activity and cellular function in Alzheimer's disease. <i>Neurobiology of Disease</i> , 2015 , 76, 13-23	7.5	62
13	Cell-cycle-regulated activation of Akt kinase by phosphorylation at its carboxyl terminus. <i>Nature</i> , 2014 , 508, 541-5	50.4	232
12	β-Hydroxyundec-9-enoic acid induces apoptosis through ROS-mediated endoplasmic reticulum stress in non-small cell lung cancer cells. <i>Biochemical and Biophysical Research Communications</i> , 2014 , 448, 267-73	3.4	22
11	Saikosaponin C inhibits lipopolysaccharide-induced apoptosis by suppressing caspase-3 activation and subsequent degradation of focal adhesion kinase in human umbilical vein endothelial cells. <i>Biochemical and Biophysical Research Communications</i> , 2014 , 445, 615-21	3.4	15
10	Catechin-7-O-xyloside induces apoptosis via endoplasmic reticulum stress and mitochondrial dysfunction in human non-small cell lung carcinoma H1299 cells. <i>Oncology Reports</i> , 2014 , 31, 314-20	3.5	6
9	Co-stimulation of TLR4 and Dectin-1 Induces the Production of Inflammatory Cytokines but not TGF-β for Th17 Cell Differentiation. <i>Immune Network</i> , 2014 , 14, 30-7	6.1	10
8	The recombinant kringle domain of urokinase plasminogen activator inhibits VEGF165-induced angiogenesis of HUVECs by suppressing VEGFR2 dimerization and subsequent signal transduction. <i>IUBMB Life</i> , 2012 , 64, 259-65	4.7	6
7	5-Phenylselenenyl- and 5-methylselenenyl-methyl-2-Deoxyuridine induce oxidative stress, DNA damage, and caspase-2-dependent apoptosis in cancer cells. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2012 , 17, 200-16	5.4	26
6	p38 mitogen-activated protein kinase is a key regulator of 5-phenylselenenyl- and 5-methylselenenyl-methyl-2-Deoxyuridine-induced apoptosis in human HL-60 cells. <i>Biochemical and Biophysical Research Communications</i> , 2012 , 417, 237-44	3.4	9
5	Combined treatment with the Cox-2 inhibitor niflumic acid and PPAR α ligand ciglitazone induces ER stress/caspase-8-mediated apoptosis in human lung cancer cells. <i>Cancer Letters</i> , 2011 , 300, 134-44	9.9	36
4	Sequential caspase-2 and caspase-8 activation is essential for saikosaponin a-induced apoptosis of human colon carcinoma cell lines. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2011 , 16, 184-97	5.4	35
3	Potent radiosensitizing agents: 5-methylselenenyl- and 5-phenylselenenyl-methyl-2-Deoxyuridine. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2011 , 21, 1151-4	2.9	4
2	Cytotoxicity and genotoxicity of titanium dioxide nanoparticles in UVA-irradiated normal peripheral blood lymphocytes. <i>Drug and Chemical Toxicology</i> , 2011 , 34, 277-84	2.3	20
1	Titanium dioxide nanoparticles trigger p53-mediated damage response in peripheral blood lymphocytes. <i>Environmental and Molecular Mutagenesis</i> , 2008 , 49, 399-405	3.2	290