

Alfred Uchman

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2677973/publications.pdf>

Version: 2024-02-01

200
papers

4,383
citations

94433

37
h-index

161849

54
g-index

200
all docs

200
docs citations

200
times ranked

1973
citing authors

#	ARTICLE	IF	CITATIONS
1	Arthropod trackways and their preservational variants from the Bagh Formation (Upper Cretaceous), India. <i>Cretaceous Research</i> , 2022, 130, 105038.	1.4	2
2	Ichnology of Lower Cretaceous prodelta and delta front deposits of the Sidi Khalif Formation, Central Tunisia. <i>Facies</i> , 2022, 68, 1.	1.4	1
3	Were Pleistocene proglacial lakes biological deserts? Insights from varved clays in Lithuania. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2022, 594, 110928.	2.3	1
4	<i>Segmentichnus mohri</i> gen. et sp. nov., a giant new trace fossil from the Culm facies (lower) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 547	0.5	0
5	Crowded <i>Trichophycus</i> ichnofabrics in the early Ordovician successions of central Iran: insight into the Ordovician radiation. <i>Lethaia</i> , 2021, 54, 314-329.	1.4	6
6	Pliocene and late Pleistocene (MIS 5e) decapod crustaceans from Santa Maria Island (Azores) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 547 <i>Quaternary Science</i> , 2021, 36, 91-109.	2.1	6
7	Burrows of the common field-cricket <i>Gryllus campestris</i> Linnaeus, 1758 (Orthoptera: Gryllidae) from Dajti Mountain, Albania. <i>Ichnos</i> , 2021, 28, 46-55.	0.5	2
8	Bioerosion structures from the Pliocene of the Agua Amarga Subbasin (Almería, SE Spain): Palaeoecological and palaeoenvironmental implications. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2021, 562, 110071.	2.3	5
9	Extra-large grains in Late Glacial " Early Holocene aeolian inland dune deposits of cold climate, European Sand Belt, Poland: An evidence of hurricane-speed frontal winds. <i>Sedimentary Geology</i> , 2021, 415, 105847.	2.1	11
10	Diverse bioerosion structures in lower Pliocene deposits from a volcanic oceanic island: Baía de Nossa Senhora section, Santa Maria Island, Azores (central North Atlantic). <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2021, 569, 110284.	2.3	3
11	Dinoflagellate cyst biostratigraphy of Upper Cretaceous turbiditic deposits from a part of the Białowiec section in the Skole Nappe (Outer Carpathians, southern Poland). <i>Cretaceous Research</i> , 2021, 123, 104780.	1.4	3
12	Paleocene-Eocene volcanic segmentation of the Norwegian-Greenland seaway reorganized high-latitude ocean circulation. <i>Communications Earth & Environment</i> , 2021, 2, .	6.8	10
13	Ichnology, sedimentology, and orbital cycles in the hemipelagic Early Jurassic Laurusian Seaway (Pliensbachian, Cardigan Bay Basin, UK). <i>Global and Planetary Change</i> , 2021, 207, 103648.	3.5	9
14	Palaeoenvironmental changes after the Messinian Salinity Crisis in the Mediterranean Almería-Nájar Basin (SE Spain) recorded by benthic foraminifera. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2021, 577, 110536.	2.3	0
15	Pleistocene coralline algal buildups on a mid-ocean rocky shore " Insights into the MIS 5e record of the Azores. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2021, 579, 110598.	2.3	2
16	Environmental conditions during the late Oligocene transgression in the North Alpine Foreland Basin (Eferding Formation, Egerian) " A multidisciplinary approach. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2021, 580, 110527.	2.3	5
17	New Ediacaran fossils from the Ukraine, some with a putative tunicate relationship. <i>Palaontologische Zeitschrift</i> , 2021, 95, 623-639.	1.6	2
18	Comment on "Decadal to millennial variations in water column parameters in pelagic marine environments of the Western Tethys (Carpathian realm) during Middle "Late Jurassic " Evidence from the radiolarian record" by M. Bălk, K. Bălk and M. Michalik. <i>Global and Planetary Change</i> , 2020, 193, 102855.	3.5	0

#	ARTICLE	IF	CITATIONS
19	Ichnological analysis of the Cenomanian–Turonian boundary interval in a collapsing slope setting: A case from the Rio Fardes section, southern Spain. <i>Cretaceous Research</i> , 2020, 106, 104262.	1.4	7
20	Bivalve bioerosion in Cretaceous-Neogene amber around the globe, with implications for the ichnogenera <i>Teredolites</i> and <i>Apectoichnus</i> . <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2020, 538, 109410.	2.3	7
21	<i>Estrellichnus jacaensis</i> from the Eocene Jaca Basin of NE Spain: new locality and new ethological interpretation. <i>Lethaia</i> , 2020, 53, 129-143.	1.4	2
22	Taxis behaviour of burrowing organisms recorded in an Ediacaran trace fossil from Ukraine. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2020, 538, 109441.	2.3	5
23	Oxygen as a factor controlling palynological record: An example from the Cenomanian–Turonian transition in the Rybie section, Polish Carpathians. <i>Marine and Petroleum Geology</i> , 2020, 112, 104067.	3.3	3
24	The 9th International Bioerosion Workshop. <i>Facies</i> , 2020, 66, 1.	1.4	0
25	Provenance of Upper Oligocene to Lower Miocene Krosno Formation sandstones in the Skole Nappe (southeast Poland): New insights from heavy minerals. <i>Geological Journal</i> , 2020, 55, 4625-4641.	1.3	1
26	Neogene marine sediments and biota encapsulated between lava flows on Santa Maria Island (Azores). <i>Sedimentology</i> , 2020, 67, 3595-3618.	3.1	9
27	Abundant trace fossil <i>Polykampton</i> in Palaeogene deep-sea flysch deposits of the Lesser Caucasus in Georgia: Palaeoecological and palaeoenvironmental implications. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2020, 558, 109958.	2.3	3
28	Upper Cretaceous bottom current deposits, north-east Greenland. <i>Sedimentology</i> , 2020, 67, 3619-3654.	3.1	11
29	Avian diversity and behavior in an Eocene coastal plain, Svalbard: the ichnological evidence. <i>Ichnos</i> , 2020, 27, 334-343.	0.5	2
30	The upper Miocene Deurne Member of the Diest Formation revisited: unexpected results from the study of a large temporary outcrop near Antwerp International Airport, Belgium. <i>Geologica Belgica</i> , 2020, 23, 219-252.	1.1	7
31	The trace fossil <i>Circulichnis</i> as a record of feeding exploration: New data from deep-sea Oligocene–Miocene deposits of northern Italy. <i>Comptes Rendus - Palevol</i> , 2019, 18, 1-12.	0.2	8
32	Ecological snapshot of a population of <i>Panoepa</i> within their traces (Pliocene, Agua Amarga subbasin). <i>Sedimentology</i> , 2019, 66, 1095-1106.	2.3	6
33	A NEW TEICHICHNOID TRACE FOSSIL SYRINGOMORPHA CYPRENSIS FROM THE MIOCENE OF CYPRUS. <i>Palaios</i> , 2019, 34, 506-514.	1.3	6
34	BURROWS OF THE POLYCHAETE PERINEREIS AIBUHIUTENSIS ON A TIDAL FLAT OF THE YELLOW RIVER DELTA IN CHINA: IMPLICATIONS FOR THE ICHNOFOSSILS POLYKLADICHNUS AND ARCHAEONASSA. <i>Palaios</i> , 2019, 34, 271-279.	1.3	12
35	Ichnological and Sedimentological Characteristics of Submarine Fan-Delta Deposits in a Half-Graben, Lower Cretaceous Palnatokes Bjerg Formation, NE Greenland. <i>Ichnos</i> , 2019, 26, 28-57.	0.5	7
36	Cold-seep fossil macrofaunal assemblages from Vestnesa Ridge, eastern Fram Strait, during the past 45 000 years. <i>Polar Research</i> , 2019, 38, .	1.6	10

#	ARTICLE	IF	CITATIONS
37	Alternating stripmining and sequestration in deep-sea sediments: The trace fossil <i>Polykampton</i> – an ecologic and ichnotaxonomic evaluation. <i>Palaeontologia Electronica</i> , 2019, 22, .	0.9	1
38	New interpretation of the provenance of crystalline material from Oligocene flysch deposits of the Skole Nappe, Poland: evidence from heavy minerals and clasts in the Nowy Borek section. <i>Geologos</i> , 2019, 25, 163-174.	0.6	1
39	Topological analysis of graphoglyptid trace fossils, a study of macrobenthic solitary and collective animal behaviors in the deep-sea environment. <i>Paleobiology</i> , 2018, 44, 306-325.	2.0	14
40	Lower Jurassic Bahamian-type facies in the ChoÅ•Nappe (Tatra Mts, West Carpathians, Poland) influenced by paleocirculation in the Western Tethys. <i>Facies</i> , 2018, 64, 1.	1.4	4
41	Peter Pervesler (1.09.1951 – 25.10.2015) as an Ichnologist and Colleague. <i>Ichnos</i> , 2018, 25, 295-298.	0.5	1
42	Dasycladacean alga <i>Palaeodasycladus</i> in the northern Tethys (West Carpathians, Poland) and its new palaeogeographic range during the Early Jurassic. <i>Swiss Journal of Geosciences</i> , 2018, 111, 305-315.	1.2	3
43	Subfossil markers of climate change during the Roman Warm Period of the late Holocene. <i>Die Naturwissenschaften</i> , 2018, 105, 6.	1.6	5
44	<i>Spirolites radwanskii</i> n. igen. n. isp.: vermetid gastropod attachment etching trace from the middle Miocene rocky coast of the Paratethys, Poland. <i>Journal of Paleontology</i> , 2018, 92, 883-895.	0.8	6
45	Spider burrows in ichnological context: a review of literature data and burrows of the wolf spider <i>Trochosa hispanica</i> Simon, 1870 from Albania. <i>Rendiconti Lincei</i> , 2018, 29, 67-79.	2.2	8
46	The Trace Fossil <i>Polykampton cabellae</i> isp. nov. from the Pagliaro Formation (Paleocene), Northern Apennines, Italy: A Record of Nutritional Sediment Sequestration by a Deep Sea Invertebrate. <i>Ichnos</i> , 2018, 25, 1-10.	0.5	4
47	The former presence of organic matter caused its later absence: Burn-down of organic matter in oceanic red beds enhanced by bioturbation (Eocene Variegated Shale, Carpathians). <i>Sedimentology</i> , 2018, 65, 1504-1519.	3.1	8
48	Trace fossils of an amalgamated storm-bed succession from the Jurassic of the Kachchh Basin, India: The significance of time-averaging in ichnology. <i>Journal of Palaeogeography</i> , 2018, 7, 14-31.	1.9	34
49	A RECORD OF SEQUESTRATION OF PLANT MATERIAL BY MARINE BURROWING ANIMALS AS A NEW FEEDING STRATEGY UNDER OLIGOTROPHIC CONDITIONS EVIDENCED BY PYRITE MICROTEXTURES. <i>Palaios</i> , 2018, 33, 312-322.	1.3	1
50	The bivalve boring <i>Cuenulites amygdaloides</i> nov. isp. in siliceous sponges from the Upper Cretaceous of Germany. <i>Geobios</i> , 2018, 51, 481-486.	1.4	3
51	FEEDING TRACES OF RECENT RAY FISH AND OCCURRENCES OF THE TRACE FOSSIL <i>PISCICHNUS WAITEMATA</i> FROM THE PLIOCENE OF SANTA MARIA ISLAND, AZORES (NORTHEAST ATLANTIC). <i>Palaios</i> , 2018, 33, 361-375.	1.3	27
52	Large cruzianid trace fossils in the Ordovician of the peri-Baltic area: the case of the BukÅ•wka Formation (Holy Cross Mountains, Poland). <i>Geological Quarterly</i> , 2018, 62, .	0.2	3
53	Ichnological record of the Frasnian – Famennian boundary interval: two examples from the Holy Cross Mts (Central Poland). <i>International Journal of Earth Sciences</i> , 2017, 106, 157-170.	1.8	11
54	<i>Rhizocorallium hamatum</i> (Fischer-Ooster 1858), a <i>Zoophycos</i> -like trace fossil from deep-sea Cretaceous-Neogene sediments. <i>Historical Biology</i> , 2017, 29, 395-410.	1.4	5

#	ARTICLE	IF	CITATIONS
55	Evolutionary trend of Zoophycos morphotypes from the Upper Cretaceous-Lower Miocene in the type pelagic sections of Gubbio, Italy. <i>Lethaia</i> , 2017, 50, 41-57.	1.4	22
56	Life beneath ammonite shells – A unique Late Cretaceous habitat for the trace maker of Chondrites and its impact on taphonomy of the shells. <i>Cretaceous Research</i> , 2017, 72, 151-160.	1.4	2
57	Intense hurricane transports sand onshore: Example from the Pliocene Malbusca section on Santa Maria Island (Azores, Portugal). <i>Marine Geology</i> , 2017, 385, 244-249.	2.1	18
58	Macroboring, their tracemakers and nestlers in clasts of a fan delta: the Savignone Conglomerate (Lower Oligocene), Northern Apennines, Italy. <i>Neues Jahrbuch Fur Geologie Und Palaontologie - Abhandlungen</i> , 2017, 283, 35-51.	0.4	13
59	The invertebrate trace fossil <i>Labyrinthichnus</i> in the Late Triassic red beds of the Argana Basin (Western Tj ETQq1 1 0,784314 ggBT / Over	2.0	15
60	Mayfly Burrows in Firmground of Recent Rivers from the Czech Republic and Poland, with Some Comments on Ephemeropteran Burrows in General. <i>Ichnos</i> , 2017, 24, 191-203.	0.5	10
61	The trace fossil <i>Diopatrighnus</i> <i>antamariensis</i> nov. isp. – A shell armored tube from Pliocene sediments of Santa Maria Island, Azores (NE Atlantic Ocean). <i>Geobios</i> , 2017, 50, 459-469.	1.4	10
62	Slumping in the Upper Jurassic Baisakhi Formation of the Jaisalmer Basin, western India: Sign of syndimentary tectonics?. <i>Journal of Palaeogeography</i> , 2017, 6, 321-332.	1.9	13
63	The Faraoni event (latest Hauterivian) in ichnological record: The RÃo Argos section of southern Spain. <i>Cretaceous Research</i> , 2017, 79, 109-121.	1.4	15
64	From morphology to behaviour: Quantitative morphological study of the trace fossil <i>Helminthorhaphe</i> . <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2017, 485, 946-955.	2.3	7
65	Evaluating macrobenthic response to the Cretaceous – Palaeogene event: A high-resolution ichnological approach at the Agost section (SE Spain). <i>Cretaceous Research</i> , 2017, 70, 96-110.	1.4	24
66	A glimpse of a fish face – An exceptional fish feeding trace fossil from the Lower Devonian of the Holy Cross Mountains, Poland. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2016, 454, 113-124.	2.3	10
67	Vertically-oriented trace fossil <i>Macaronichnus</i> <i>segregatis</i> from Neogene of Santa Maria Island (Azores; NE Atlantic) records vertical fluctuations of the coastal groundwater mixing zone on a small oceanic island. <i>Geobios</i> , 2016, 49, 229-241.	1.4	30
68	Late Ordovician Trace Fossils from Offshore to Shallow Water Mixed Siliciclastic and Carbonate Facies in the Ringerike Area, Oslo Region, Norway. <i>Ichnos</i> , 2016, 23, 189-221.	0.5	13
69	Rocking around a volcanic island shelf: Pliocene Rhodolith beds from Malbusca, Santa Maria Island (Azores, NE Atlantic). <i>Facies</i> , 2016, 62, 1.	1.4	19
70	Depositional environment, ichnological features and oxygenation of Permian to earliest Triassic marine sediments in central Spitsbergen, Svalbard. <i>Polar Research</i> , 2016, 35, 24782.	1.6	20
71	Ichnology of prodelta deposits of the Mezardere Formation (late Eocene – early Oligocene) in the Çankaya Island, western Turkey. <i>Geodinamica Acta</i> , 2016, 28, 86-100.	2.2	7
72	Underground Miners Come Out to the Surface – Trails of Earthworms. <i>Ichnos</i> , 2016, 23, 99-107.	0.5	7

#	ARTICLE	IF	CITATIONS
73	The Probable Isopod Burrow <i>Sinusichnus seilacheri</i> isp. n. from the Middle Triassic of Germany: An Example of Behavioral Convergence. <i>Ichnos</i> , 2016, 23, 138-146.	0.5	15
74	The End-Cretaceous Extinction and Ecosystem Change. <i>Topics in Geobiology</i> , 2016, , 265-300.	0.5	11
75	The Stebnyk Formation (Miocene) in the Boryslav-Pokuttya and Sambir nappes of the Ukrainian Carpathians: a record of environmental change in the Carpathian Foredeep. <i>Geological Quarterly</i> , 2016, , .	0.2	4
76	GELÄ°BOLU YARIMADASI GEÄ± EOSEN YAÄZLI CEYLAN FORMASYONU DERÄ°N DENÄ°Z Ä±Ä–KELLERÄ° Ä°Z FOSÄ°LLERÄ° (GB) Tj ETQqC	0.1	0
77	Eocene flora and trace fossils from the Hruby Regiel section in the Tatra Mountains (Poland): Taxonomic revision of the Wiktor KuÄ±niar fossil plant collection. <i>Acta Geologica Polonica</i> , 2015, 65, 215-238.	0.9	3
78	Fossilized bioelectric wire â€” the trace fossil <i>Trichichnus</i> . <i>Biogeosciences</i> , 2015, 12, 2301-2309.	3.3	37
79	Borings in gneiss boulders in the Miocene (Upper Tortonian) of the Sorbas Basin, SE Spain. <i>Geological Magazine</i> , 2015, 152, 287-297.	1.5	12
80	A delayed response of the trace fossil community at the Cretaceous-Paleogene boundary in the Bottaccione section, Gubbio, Central Italy. <i>Geobios</i> , 2015, 48, 137-145.	1.4	23
81	The trace fossil <i>Gyrophyllites</i> in deep-sea siliciclastic deposits of the Istebna Formation (Upper) Tj ETQq1 1 0.784314 rgBT /Overlock 1 Palaeogeography, Palaeoclimatology, Palaeoecology, 2015, 426, 260-274.	2.3	7
82	The trace fossil <i>Lepidenteron lewesiensis</i> : a taphonomic window on diversity of Late Cretaceous fishes. <i>Palaontologische Zeitschrift</i> , 2015, 89, 795-806.	1.6	12
83	Palaeoecology, taphonomy, and preservation of a lower Pliocene shell bed (coquina) from a volcanic oceanic island (Santa Maria Island, Azores). <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2015, 430, 57-73.	2.3	44
84	Deep Endichnial <i>Cruziana</i> from the Lower-Middle Ordovician of Spain â€” A Unique Trace Fossil Record of Trilobitomorph Deep Burrowing Behavior. <i>Ichnos</i> , 2015, 22, 12-18.	0.5	5
85	Last occurrence of <i>Abathomphalus mayaroensis</i> (Bolli) foraminiferid index of the Cretaceousâ€”Paleogene boundary: the calcareous nannofossil proof. <i>Geologica Carpathica</i> , 2015, 66, 181-195.	0.7	3
86	How bioturbation obscured the Cretaceousâ€”Palaeogene boundary record. <i>Terra Nova</i> , 2015, 27, 225-230.	2.1	34
87	Comment on â€œOccurrence of faecal pellet-filled simple and composite burrows in cold seep carbonates: A glimpse of a complex benthic ecosystemâ€”by A. Mazumdar, R.K. Joshi and M. Kocherla [Marine Geology 289 (2011) 117â€”121]. <i>Marine Geology</i> , 2015, 364, 65-67.	2.1	2
88	Lower Ordovician (Arenig) shallow-marine trace fossils of the Pochico Formation, southern Spain: palaeoenvironmental and palaeogeographic implications at the Gondwanan and peri-Gondwanan realm. <i>Journal of Iberian Geology</i> , 2014, 40, .	1.3	12
89	Cambrian trace fossils of the <i>Cruziana</i> ichnofacies from the Bikaner-Nagaur Basin, north western Indian Craton. <i>Journal of Asian Earth Sciences</i> , 2014, 81, 129-141.	2.3	26
90	Deep-sea trace fossils of the Oligoceneâ€”Miocene Numidian Formation, northern Tunisia. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2014, 414, 155-177.	2.3	26

#	ARTICLE	IF	CITATIONS
91	Unusually well preserved casts of halite crystals: A case from the Upper Frasnian of northern Lithuania. <i>Sedimentary Geology</i> , 2014, 308, 44-52.	2.1	4
92	Differential Effects of Bioturbation on Benthic Foraminiferal Distribution Across the Cretaceous-Palaeogene (K-Pg) Boundary at Bidart (Southwestern France). <i>Springer Geology</i> , 2014, , 61-63.	0.3	1
93	Ichnological characteristics of Late Cretaceous hemipelagic and pelagic sediments in a submarine high around the OAE-2 event: A case from the Rybie section, Polish Carpathians. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2013, 370, 222-231.	2.3	38
94	Exceptionally favourable life conditions for macrobenthos during the Late Cenomanian OAE-2 event: Ichnological record from the Bonarelli Level in the Grajcarek Unit, Polish Carpathians. <i>Cretaceous Research</i> , 2013, 46, 1-10.	1.4	24
95	Scratch circles associated with the large foraminifer <i>Bathysiphon</i> from deep-sea turbiditic sediments of the Pagliaro Formation (Palaeocene), Northern Apennines, Italy. <i>Sedimentary Geology</i> , 2013, 289, 115-123.	2.1	5
96	The new trace fossil <i>Gyrolithes lorcaensis</i> isp. n. from the Miocene of SE Spain and a critical review of the <i>Gyrolithes</i> ichnospecies. <i>Stratigraphy and Geological Correlation</i> , 2013, 21, 312-322.	0.8	17
97	The trace fossil <i>Lepidenteron lewesiensis</i> (Mantell, 1822) from the Upper Cretaceous of southern Poland. <i>Acta Geologica Polonica</i> , 2013, 63, 611-623.	0.9	12
98	First record of catacrinid crinoid from the Lower Permian of Spitsbergen. <i>Polish Polar Research</i> , 2013, 34, 139-150.	0.9	2
99	<i>Cladichnus parallelum</i> isp. nov. - a Mid- to Deep-Tier Feeding Burrow System. <i>Ichnos</i> , 2013, 20, 120-128.	0.5	8
100	Late Maastrichtian foraminiferids and diatoms from the Polish Carpathians (Ropianka Formation,). <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50</i> 2013, 63, 515-525.	0.9	6
101	Conventional and high-resolution heavy mineral analyses applied to flysch deposits: comparative provenance studies of the Ropianka (Upper Cretaceous-Paleocene) and Menilite (Oligocene) formations (Skole Nappe, Polish Carpathians). <i>Geological Quarterly</i> , 2013, 57, .	0.2	7
102	ICHOLOGICAL ANALYSIS OF LATERAL ENVIRONMENTAL HETEROGENEITY WITHIN THE BONARELLI LEVEL (UPPERMOST CENOMANIAN) IN THE CLASSICAL LOCALITIES NEAR GUBBIO, CENTRAL APENNINES, ITALY. <i>Palaios</i> , 2012, 27, 48-54.	1.3	40
103	Ichnology of Upper Cretaceous deep-sea thick-bedded flysch sandstones: Lower Istebna Beds, Silesian Unit (Outer Carpathians, southern Poland). <i>Geologica Carpathica</i> , 2012, 63, 107-120.	0.7	10
104	Integrated ichnology and ichthyology of the Oligocene Menilite Formation, Skole and Subsilesian nappes, Polish Carpathians: A proxy to oxygenation history. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2012, 331-332, 104-118.	2.3	57
105	The trace fossil <i>Nummipera eocenica</i> from the Tatra Mountains, Poland: morphology and palaeoenvironmental implications. <i>Lethaia</i> , 2012, 45, 342-355.	1.4	14
106	Bio-events, foraminiferal and nannofossil biostratigraphy of the Cenomanian/Turonian boundary interval in the Subsilesian Nappe, Rybie section, Polish Carpathians. <i>Cretaceous Research</i> , 2012, 35, 181-198.	1.4	18
107	The late Barremian <i>Halimedes</i> horizon of the Dolomites (Southern Alps, Italy). <i>Cretaceous Research</i> , 2012, 35, 199-207.	1.4	16
108	A History of Ideas in Ichnology. <i>Developments in Sedimentology</i> , 2012, 64, 3-43.	0.5	17

#	ARTICLE	IF	CITATIONS
109	Deep-Sea Fans. <i>Developments in Sedimentology</i> , 2012, 64, 643-671.	0.5	39
110	Hemipelagic and Pelagic Basin Plains. <i>Developments in Sedimentology</i> , 2012, , 673-701.	0.5	36
111	Glacial Environments. <i>Developments in Sedimentology</i> , 2012, , 299-327.	0.5	12
112	Late Pleistoceneâ€“early Holocene polychaete borings in NE Spitsbergen and their palaeoecological and climatic implications: an example from the Basissletta area. <i>Boreas</i> , 2012, 41, 42-55.	2.4	16
113	The miniature echinoid trace fossil <i>Bichordites kuzunensis</i> isp. nov. from early Oligocene prodelta sediments of the Mezardere Formation, GÅŕkÅšeada Island, NW Turkey. <i>Acta Geologica Polonica</i> , 2012, 62, 205-215.	0.9	9
114	Probable root structures and associated trace fossils from the Lower Pleistocene calcarenites of Favignana Island, southern Italy: dilemmas of interpretation. <i>Geological Quarterly</i> , 2012, 56, 745-756.	0.2	6
115	Biostratigraphy and palaeoenvironment of the Kimmeridgian-Lower Tithonian pelagic deposits of the KrÅŕ3/4na Nappe, Lejowa Valley, Tatra Mts. (southern Poland). <i>Geological Quarterly</i> , 2012, 56, 773-788.	0.2	14
116	Foreland provenance of thick conglomerates in the early stage of the Carpathian Foredeep development: the case of the Sloboda Conglomerate (Lower Miocene), western Ukraine. <i>Geological Quarterly</i> , 2012, 56, 789-802.	0.2	10
117	Heavy minerals from Oligocene sandstones of the Menilite Formation of the Skole Nappe, SE Poland: a tool for provenance specification. <i>Geological Quarterly</i> , 2012, 56, 803-820.	0.2	7
118	ICHTHOLOGICAL RECORD OF ENVIRONMENTAL CHANGES IN EARLY QUATERNARY (GELASIAN-CALABRIAN) MARINE DEPOSITS OF THE STIRONE SECTION, NORTHERN ITALY. <i>Palaios</i> , 2011, 26, 578-593.	1.3	19
119	The Trace Fossil <i>Diplopodichnus</i> from the Lower Jurassic Lacustrine Sediments of Central China and the Isopod <i>Armadillidium vulgare</i> (Pillbug) Lebensspuren as its Recent Analogue. <i>Ichnos</i> , 2011, 18, 147-155.	0.5	13
120	Deep-Sea Ichnology: The Relationships Between Depositional Environment and Endobenthic Organisms. <i>Developments in Sedimentology</i> , 2011, 63, 517-556.	0.5	46
121	The new complex helical trace fossil <i>Avetoichnus luisae</i> igen. n. et isp. n. from the Cainozoic deep-sea sediments of the Alpine realm: a non-graphoglyptid mid-tier agrichnion. <i>Neues Jahrbuch Fur Geologie Und Palaontologie - Abhandlungen</i> , 2011, 260, 319-330.	0.4	6
122	The Cretaceous-Paleogene boundary in turbiditic deposits identified to the bed: a case study from the Skole Nappe (Outer Carpathians, southern Poland). <i>Geologica Carpathica</i> , 2011, 62, 333-343.	0.7	6
123	Vertical displacement and taphonomic filtering of nannofossils by bioturbation in the Cretaceousâ€“Palaeogene boundary section at Caravaca, SE Spain. <i>Lethaia</i> , 2011, 44, 321-328.	1.4	30
124	Mollusc trace fossils <i>Ptychoplasma</i> Fenton and Fenton, 1937 and <i>Oravaichnium</i> PliÅŕka and UhrovÅŕj, 1990: Their type material and ichnospecies. <i>Geobios</i> , 2011, 44, 387-397.	1.4	37
125	Ichnological data as a useful tool for deep-sea environmental characterization: a brief overview and an application to recognition of small-scale oxygenation changes during the Cenomanianâ€“Turonian anoxic event. <i>Geo-Marine Letters</i> , 2011, 31, 525-536.	1.1	21
126	Impact of the Paleoceneâ€“Eocene Thermal Maximum on the macrobenthic community: Ichnological record from the Zumaia section, northern Spain. <i>Marine Geology</i> , 2011, 282, 178-187.	2.1	26

#	ARTICLE	IF	CITATIONS
127	Ichnological analysis of the Bidart and Sopelana Cretaceous/Paleogene (K/Pg) boundary sections (Basque Basin, W Pyrenees): Refining eco-sedimentary environment. <i>Sedimentary Geology</i> , 2011, 234, 42-55.	2.1	26
128	Trace fossils in Quaternary glacial varved clays near Uppsala, Sweden. <i>Gff</i> , 2011, 133, 135-140.	1.2	6
129	The Global Stratotype Section and Point (GSSP) for the base of the Lutetian Stage at the Gorrondatxe section, Spain. <i>Episodes</i> , 2011, 34, 86-108.	1.2	69
130	<i>Phymatoderma melvillensis</i> isp. nov. and other trace fossils from the Cape Melville Formation (Lower Miocene) of King George Island, Antarctica. <i>Polish Polar Research</i> , 2010, 31, 83-99.	0.9	18
131	Bioturbational redistribution of Danian calcareous nannofossils in the uppermost Maastrichtian across the K-Pg boundary at Bidart, SW France. <i>Geobios</i> , 2010, 43, 569-579.	1.4	42
132	A new ichnogenus <i>Skolichnus</i> for <i>Chondrites hoernesii</i> Ettingshausen, 1863, a deep-sea radial trace fossil from the Upper Cretaceous of the Polish Flysch Carpathians: Its taxonomy and palaeoecological interpretation as a deep-tier chemichnion. <i>Cretaceous Research</i> , 2010, 31, 515-523.	1.4	8
133	Sea-level dynamics and palaeoecological factors affecting trace fossil distribution in Eocene turbiditic deposits (Gorrondatxe section, N Spain). <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2010, 285, 50-65.	2.3	45
134	A hypersaline ichnoassemblage from the Middle Triassic carbonate ramp of the Tatricum domain in the Tatra Mountains, Southern Poland. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2010, 292, 71-81.	2.3	39
135	ICHTNOFABRIC EVIDENCE FOR THE LACK OF BOTTOM ANOXIA DURING THE LOWER TOARCIAN OCEANIC ANOXIC EVENT IN THE FUENTE DE LA VIDRIERA SECTION, BETIC CORDILLERA, SPAIN. <i>Palaios</i> , 2010, 25, 576-587.	1.3	56
136	<i>Arthrodendron maguricum</i> n. sp., a new larger agglutinated foraminifer from the Eocene Magura flysch of the Polish Carpathians and its relationship to komokiaceans and trace fossils. <i>Journal of Paleontology</i> , 2010, 84, 1015-1021.	0.8	1
137	<i>Omanichnus</i> and <i>Vitichnus</i> – Two New Graphoglyptid Ichnogenera from Upper Triassic Deep-Sea Fan Deposits in Oman. <i>Ichnos</i> , 2009, 16, 179-185.	0.5	1
138	Cyclostratigraphic dating in the Lower Badenian (Middle Miocene) of the Vienna Basin (Austria): the Baden-Sooss core. <i>International Journal of Earth Sciences</i> , 2009, 98, 915-930.	1.8	35
139	Upper bathyal trace fossils document palaeoclimate changes. <i>Terra Nova</i> , 2009, 21, 229-236.	2.1	3
140	Oceanic Anoxic Event at the Cenomanian – Turonian boundary interval (OAE-2): ichnological approach from the Betic Cordillera, southern Spain. <i>Lethaia</i> , 2009, 42, 407-417.	1.4	53
141	Nutrient spatial variation during intrabasinal upwelling at the Cenomanian – Turonian oceanic anoxic event in the westernmost Tethys: An ichnological and facies approach. <i>Sedimentary Geology</i> , 2009, 215, 83-93.	2.1	38
142	Large chambered sponge borings on a Late Cretaceous abrasion platform at Cracow, Poland. <i>Cretaceous Research</i> , 2009, 30, 149-160.	1.4	18
143	Trace fossils from Late Pleistocene varved lacustrine sediments in eastern Lithuania. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2009, 272, 199-211.	2.3	57
144	The <i>Ophiomorpha rudis</i> ichnosubfacies of the Nereites ichnofacies: Characteristics and constraints. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2009, 276, 107-119.	2.3	92

#	ARTICLE	IF	CITATIONS
145	Field expressions of the transformation of debris flows into turbidity currents, with examples from the Polish Carpathians and the French Maritime Alps. <i>Marine and Petroleum Geology</i> , 2009, 26, 2011-2020.	3.3	36
146	Rheotactic <i>Macaronichnus</i> , and Human and Cattle Trackways in Holocene Beachrock, Greece: Reconstruction of Paleoshoreline Orientation. <i>Ichnos</i> , 2009, 16, 103-117.	0.5	42
147	Latest Maastrichtian foraminiferal assemblages from the Huszár region (Skole Nappe, Outer Tethyan) BT / Overlock 10	0.7	9
148	A New Y-Shaped Trace Fossil Attributed to Upogebiid Crustaceans from Early Pleistocene of Italy. <i>Acta Palaeontologica Polonica</i> , 2009, 54, 135-142.	0.4	35
149	A larger agglutinated foraminifer originally described as a marine plant: the case of <i>Arthrodendron</i> Ulrich, 1904 (Foraminifera), its synonyms and homonyms. <i>Journal of Micropalaeontology</i> , 2008, 27, 103-110.	3.6	3
150	Bioturbational disturbance of the Cretaceous-Palaeogene (K-Pg) boundary layer: Implications for the interpretation of the K-Pg boundary impact event. <i>Geobios</i> , 2008, 41, 661-667.	1.4	46
151	Palaeoenvironmental turnover across the Ypresian-Lutetian transition at the Agost section, Southeastern Spain: In search of a marker event to define the Stratotype for the base of the Lutetian Stage. <i>Marine Micropaleontology</i> , 2008, 69, 297-313.	1.2	31
152	Ichnological record of deep-sea palaeoenvironmental changes around the Oceanic Anoxic Event 2 (Cenomanian-Turonian boundary): An example from the Barnasiwka section, Polish Outer Carpathians. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2008, 262, 61-71.	2.3	52
153	Trace fossils from the Upper Pleistocene glaciolacustrine laminated sediments of Lithuania. <i>Geologija</i> , 2008, 50, 212-226.	0.1	11
154	A HIGHLY DIVERSE ICHNOFAUNA IN LATE TRIASSIC DEEP-SEA FAN DEPOSITS OF OMAN. <i>Palaios</i> , 2007, 22, 567-576.	1.3	46
155	Deep-sea trace fossils from the mixed carbonate-siliciclastic flysch of the Monte Antola Formation (Late Campanian-Maastrichtian), North Apennines, Italy. <i>Cretaceous Research</i> , 2007, 28, 980-1004.	1.4	38
156	Deep-Sea Ichnology: Development of Major Concepts. , 2007, , 248-267.		22
157	Comments on the paper "Reconnaissance of Upper Jurassic Morrison Formation ichnofossils, Rocky Mountain Region, USA: Palaeoenvironmental, stratigraphic, and paleoclimatic significance of terrestrial and freshwater ichnocoenoses" by Stephen T. Hasiotis. <i>Sedimentary Geology</i> , 2007, 200, 141-150.	2.1	38
158	<i>Bichordites</i> and <i>Bichordites-Rosselia</i> ; Ichnoassemblages from the Lower Pleistocene Tursi Sandstone (Southern Italy). , 2007, , .		6
159	SURFACE LEBENSSPUREN PRODUCED BY AMPHIPODS AND ISOPODS (CRUSTACEANS) FROM THE ISONZO DELTA TIDAL FLAT, ITALY. <i>Palaios</i> , 2006, 21, 384-390.	1.3	49
160	Ichnological analysis of the Cretaceous-Palaeogene boundary interval at the Caravaca section, SE Spain. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2006, 242, 313-325.	2.3	62
161	Names for trace fossils: a uniform approach. <i>Lethaia</i> , 2006, 39, 265-286.	1.4	400
162	<i>Persichnus</i> gen. nov. and Associated Ichnofossils From the Upper Cretaceous to Eocene Deep-Sea Deposits of the Sanandaj Area, West Iran. <i>Ichnos</i> , 2005, 12, 141-149.	0.5	4

#	ARTICLE	IF	CITATIONS
163	Ordovician Bathyal Trace Fossils From Metasiliciclastics in Central Norway and Their Sedimentological and Paleogeographical Implications. <i>Ichnos</i> , 2005, 12, 105-133.	0.5	26
164	Revision of the ichnogenussabellarifexrichter, 1921 and its relationship toskolithoshaldeman, 1840 andpolykladichnusfÄ¼rsich, 1981. <i>Journal of Systematic Palaeontology</i> , 2005, 3, 115-131.	1.5	71
165	Trace fossils after the KT boundary event from the Agost section, SE Spain. <i>Geological Magazine</i> , 2004, 141, 429-440.	1.5	65
166	Lithostratigraphy of the Lower Cambrian metaclastics and their age based on trace fossils in the SandÄ±rÄ± region, southwestern Turkey. <i>Geobios</i> , 2004, 37, 346-360.	1.4	37
167	Oligocene trace fossils from temporary fluvial plain ponds: An example from the Freshwater Molasse of Switzerland. <i>Eclogae Geologicae Helveticae</i> , 2004, 97, 133-148.	0.6	42
168	Phanerozoic history of deep-sea trace fossils. <i>Geological Society Special Publication</i> , 2004, 228, 125-139.	1.3	41
169	Ichnotaxonomic analysis of the Cretaceous/Palaeogene boundary interval in the Agost section, south-east Spain. <i>Cretaceous Research</i> , 2004, 25, 635-647.	1.4	43
170	Large microbial-foraminiferal oncoids from condensed Lowerâ€œMiddle Jurassic deposits: a case study from the Tatra Mountains, Poland. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2004, 213, 133-151.	2.3	36
171	Large microbial-foraminiferal oncoids from condensed Lowerâ€œMiddle Jurassic deposits: a case study from the Tatra Mountains, Poland. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2004, 213, 133-151.	2.3	20
172	Early Devonian trace fossils in marine to non-marine redbeds in Podolia, Ukraine: palaeoenvironmental implications. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2004, 214, 67-83.	2.3	34
173	Trends in diversity, frequency and complexity of graphoglyptid trace fossils: evolutionary and palaeoenvironmental aspects. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2003, 192, 123-142.	2.3	85
174	<i>Hillichnus lobosensis</i> igen. et isp. nov., a complex trace fossil produced by tellinacean bivalves, Paleocene, Monterey, California, USA. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2003, 192, 157-186.	2.3	50
175	Sequential colonization of muddy turbidites in the Eocene BeloveÅ¾a Formation, Carpathians, Poland. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2001, 168, 171-186.	2.3	127
176	The oldest deep-sea Ophiomorpha and Scolicia and associated trace fossils from the Upper Jurassicâ€œLower Cretaceous deep-water turbidite deposits of SW Bulgaria. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2001, 169, 85-99.	2.3	73
177	Upper Triassic (Keuper) non-marine trace fossils from the HaÃŸberge area (Franconia, south-eastern) <i>Tj ETQq1 1 0.784314 rgBT /Overlock_10 Tf 50 1.</i>	1.6	15
178	<i>Estrellichnus jacaensis</i> nov. igen., nov. isp. - a large radial trace fossil from Eocene flysch (Hecho) <i>Tj ETQq0 0 0 rgBT /Overlock_10 Tf 50 1.</i>	1.4	15
179	The<i>glossifungites</i> ichnofacies in the area of its nomenclatural archetype, lviv, Ukraine. <i>Ichnos</i> , 2000, 7, 183-193.	0.5	27
180	An aberrant, helicoidal trace fossil <i>Chondrites</i> Sternberg. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 1999, 146, 165-169.	2.3	15

#	ARTICLE	IF	CITATIONS
181	A <i>zoophycos</i> group trace fossil from miocene flysch in Southern Turkey: Evidence for a U-shaped causative burrow. <i>Ichnos</i> , 1999, 6, 251-259.	0.5	31
182	Lower and Middle Jurassic flysch trace fossils from the eastern Stara Planina Mountains, Bulgaria: A contribution to the evolution of Mesozoic ichnodiversity. <i>Neues Jahrbuch Fur Geologie Und Palaontologie - Abhandlungen</i> , 1999, 213, 169-199.	0.4	10
183	<i>Ichnogenus treptichnus</i> in eocene flysch, carpathians, Poland: Taxonomy and preservation. <i>Ichnos</i> , 1998, 5, 269-275.	0.5	17
184	Deep-Sea Benthic Food Content Recorded by Ichnofabrics: A Conceptual Model Based on Observations from Paleogene Flysch, Carpathians, Poland. <i>Palaaios</i> , 1998, 13, 533.	1.3	61
185	Ichnology of deep-sea fan overbank deposits of the Ganei slates (Eocene, Switzerland) – a classical flysch trace fossil locality studied first by Oswald Heer. <i>Ichnos</i> , 1997, 5, 139-162.	0.5	36
186	New biostratigraphic data on the Early Cretaceous platform carbonates of the Tatra Mountains, Western Carpathians, Poland. <i>Cretaceous Research</i> , 1997, 18, 713-729.	1.4	14
187	Trace fossils and facies changes in Cretaceous–Eocene flysch deposits of the Julian Prealps (Italy and) Tj ETQq1 1 0.784314 rgBT /Overlock 10 T	0.5	31
188	Ichnology of eocene flysch deposits of the Istria peninsula, croatia and slovenia. <i>Ichnos</i> , 1996, 5, 1-22.	0.5	22
189	Use of trace fossils in delineating sequence stratigraphic surfaces (Tertiary Venetian Basin,) Tj ETQq1 1 0.784314 rgBT /Overlock 10 T	2.3	56
190	Trace fossils indicating bottom aeration changes: Folsz Limestone, Oligocene, Outer Carpathians, Poland. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 1996, 121, 79-87.	2.3	11
191	Note on rediscovered type and figured material relating to <i>Muensteria</i> sternberg 1833. <i>Ichnos</i> , 1996, 4, 305-309.	0.5	6
192	Tiering patterns of trace fossils in the Palaeogene flysch deposits of the Carpathians, Poland. <i>Geobios</i> , 1995, 28, 389-394.	1.4	27
193	Trace fossils from Lower Miocene (Ottngian) molasse deposits of Upper Austria. <i>Palaontologische Zeitschrift</i> , 1995, 69, 503-524.	1.6	46
194	Trace fossils from interdune deposits – an example from the lower triassic aeolian Tumlin Sandstone, central Poland. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 1994, 108, 121-138.	2.3	37
195	Biogenic structures of organics – poor siliciclastic sediments: Examples from Paleogene variegated shales, Polish Carpathians. <i>Ichnos</i> , 1993, 2, 267-275.	0.5	21
196	Diverse tiering patterns in Paleogene flysch trace fossils, Magura nappe, Carpathian Mountains, Poland. <i>Ichnos</i> , 1991, 1, 287-292.	0.5	19
197	The trace fossil <i>Polykampton recurvum</i> n. isp. (<i>sequestrichnia</i>) from the Maastrichtian – Paleocene deep-sea deposits of NW Italy. <i>Journal of Paleontology</i> , 0, , 1-15.	0.8	0
198	3D sine wave-like geometry of soft sediment deformation structures: Possible record of the Late Pleistocene seismic activity over the Teisseyre – Tornquist Zone. <i>Terra Nova</i> , 0, , .	2.1	2

#	ARTICLE	IF	CITATIONS
199	Arthropod trace fossils from Eocene cold climate continental strata of King George Island, West Antarctica. <i>Acta Palaeontologica Polonica</i> , 0, 63, .	0.4	7
200	The Dobrotiv Formation (Miocene) in the Boryslav-Pokuttya and Sambir nappes of the Ukrainian Carpathians: a record of sedimentary environmental change in the development of the Carpathian Foredeep basin. <i>Geological Quarterly</i> , 0, , .	0.2	4