

Sergio Arrabal

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2682352/publications.pdf>

Version: 2024-02-01

13
papers

407
citations

759233

12
h-index

1125743

13
g-index

14
all docs

14
docs citations

14
times ranked

916
citing authors

#	ARTICLE	IF	CITATIONS
1	Adipocyte cannabinoid CB1 receptor deficiency alleviates high fat diet-induced memory deficit, depressive-like behavior, neuroinflammation and impairment in adult neurogenesis. <i>Psychoneuroendocrinology</i> , 2019, 110, 104418.	2.7	17
2	Oleylethanolamide restores alcohol-induced inhibition of neuronal proliferation and microglial activity in striatum. <i>Neuropharmacology</i> , 2019, 146, 184-197.	4.1	12
3	Acetaminophen-Induced Liver Injury Alters the Acyl Ethanolamine-Based Anti-Inflammatory Signaling System in Liver. <i>Frontiers in Pharmacology</i> , 2017, 8, 705.	3.5	18
4	Cannabinoid CB1 Receptors Are Localized in Striated Muscle Mitochondria and Regulate Mitochondrial Respiration. <i>Frontiers in Physiology</i> , 2016, 7, 476.	2.8	89
5	Antiobesity efficacy of GLP-1 receptor agonist liraglutide is associated with peripheral tissue-specific modulation of lipid metabolic regulators. <i>BioFactors</i> , 2016, 42, 600-611.	5.4	33
6	Pharmacological Blockade of Cannabinoid CB1 Receptors in Diet-Induced Obesity Regulates Mitochondrial Dihydropyridinyl Dehydrogenase in Muscle. <i>PLoS ONE</i> , 2015, 10, e0145244.	2.5	31
7	Chronic administration of recombinant IL-6 upregulates lipogenic enzyme expression and aggravates high fat diet-induced steatosis in IL-6 deficient mice. <i>DMM Disease Models and Mechanisms</i> , 2015, 8, 721-31.	2.4	34
8	Oleylethanolamide enhances β -adrenergic-mediated thermogenesis and white-to-brown adipocyte phenotype in epididymal white adipose tissue in rat. <i>DMM Disease Models and Mechanisms</i> , 2014, 7, 129-41.	2.4	51
9	Localization of peroxisome proliferator-activated receptor alpha (PPAR α) and N-acyl phosphatidylethanolamine phospholipase D (NAPE-PLD) in cells expressing the Ca ²⁺ -binding proteins calbindin, calretinin, and parvalbumin in the adult rat hippocampus. <i>Frontiers in Neuroanatomy</i> , 2014, 8, 12.	1.7	16
10	Localization of the cannabinoid CB1 receptor and the 2-AG synthesizing (DAGL β) and degrading (MAGL) in the adult rat hippocampus. <i>Frontiers in Neuroanatomy</i> , 2014, 8, 56.	1.7	27
11	Pharmacological blockade of either cannabinoid CB1 or CB2 receptors prevents both cocaine-induced conditioned locomotion and cocaine-induced reduction of cell proliferation in the hippocampus of adult male rat. <i>Frontiers in Integrative Neuroscience</i> , 2014, 7, 106.	2.1	45
12	Effects of acute versus repeated cocaine exposure on the expression of endocannabinoid signaling-related proteins in the mouse cerebellum. <i>Frontiers in Integrative Neuroscience</i> , 2014, 8, 22.	2.1	19
13	CB1 Blockade Potentiates Down-Regulation of Lipogenic Gene Expression in Perirenal Adipose Tissue in High Carbohydrate Diet-Induced Obesity. <i>PLoS ONE</i> , 2014, 9, e90016.	2.5	15