

Min Liang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2688300/publications.pdf>

Version: 2024-02-01

52
papers

903
citations

567144

15
h-index

526166

27
g-index

56
all docs

56
docs citations

56
times ranked

1244
citing authors

#	ARTICLE	IF	CITATIONS
1	Urinary albumin-to-creatinine ratio and the risk of first stroke in Chinese hypertensive patients treated with angiotensin-converting enzyme inhibitors. <i>Hypertension Research</i> , 2022, 45, 116-124.	1.5	4
2	U-shaped association between dietary copper intake and new-onset hypertension. <i>Clinical Nutrition</i> , 2022, 41, 536-542.	2.3	24
3	Association of urinary albumin:creatinine ratio with incident frailty in older populations. <i>CKJ: Clinical Kidney Journal</i> , 2022, 15, 1093-1099.	1.4	1
4	Dietary protein intake and the risk of all-cause and cardiovascular mortality in maintenance hemodialysis patients: A multicenter, prospective cohort study. <i>Nutrition</i> , 2022, 95, 111564.	1.1	6
5	Change in the Estimated Glomerular Filtration Rate Over Time and Risk of First Stroke in Hypertensive Patients. <i>Journal of Epidemiology</i> , 2022, , .	1.1	0
6	Folate intake and incident chronic kidney disease: a 30-year follow-up study from young adulthood to midlife. <i>American Journal of Clinical Nutrition</i> , 2022, 116, 599-607.	2.2	4
7	Relationship of visceral adiposity index with new-onset proteinuria in hypertensive patients. <i>Clinical Nutrition</i> , 2021, 40, 438-444.	2.3	10
8	Silencing of Central (Pro)renin Receptor Ameliorates Salt-Induced Renal Injury in Chronic Kidney Disease. <i>Antioxidants and Redox Signaling</i> , 2021, 35, 93-112.	2.5	4
9	Interaction of serum calcium and folic acid treatment on first stroke in hypertensive males. <i>Clinical Nutrition</i> , 2021, 40, 2381-2388.	2.3	2
10	Higher dietary fibre intake is associated with lower CVD mortality risk among maintenance haemodialysis patients: a multicentre prospective cohort study. <i>British Journal of Nutrition</i> , 2021, 126, 1510-1518.	1.2	9
11	Neutrophil counts and the risk of first stroke in general hypertensive adults. <i>Hypertension Research</i> , 2021, 44, 830-839.	1.5	3
12	Association of estimated glomerular filtration rate from serum creatinine and cystatin C with new-onset diabetes: a nationwide cohort study in China. <i>Acta Diabetologica</i> , 2021, 58, 1269-1276.	1.2	1
13	Association of visceral adiposity index with new-onset type 2 diabetes and impaired fasting glucose in hypertensive Chinese adults. <i>Eating and Weight Disorders</i> , 2021, , 1.	1.2	2
14	Dietary Plant Protein and Mortality Among Patients Receiving Maintenance Hemodialysis: A Cohort Study. <i>American Journal of Kidney Diseases</i> , 2021, 78, 649-657.e1.	2.1	9
15	Association of Depressive Symptoms with Rapid Kidney Function Decline in Adults with Normal Kidney Function. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2021, 16, 889-897.	2.2	25
16	Compound Heterozygosity for KLF1 Mutations Causing Hemolytic Anemia in Children: A Case Report and Literature Review. <i>Frontiers in Genetics</i> , 2021, 12, 691461.	1.1	7
17	The Effects of Oral Energy-Dense Supplements on Nutritional Status in Nondiabetic Maintenance Hemodialysis Patients. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2021, 16, 1228-1236.	2.2	10
18	Relationship of several serum folate forms with the risk of mortality: A prospective cohort study. <i>Clinical Nutrition</i> , 2021, 40, 4255-4262.	2.3	14

#	ARTICLE	IF	CITATIONS
19	Prospective association between baseline plasma zinc concentration and development of proteinuria in Chinese hypertensive patients. <i>Journal of Trace Elements in Medicine and Biology</i> , 2021, 66, 126755.	1.5	0
20	Interaction of Serum Alkaline Phosphatase and Folic Acid Treatment on Chronic Kidney Disease Progression in Treated Hypertensive Adults. <i>Frontiers in Pharmacology</i> , 2021, 12, 753803.	1.6	2
21	U-shaped association between serum albumin and development of chronic kidney disease in general hypertensive patients. <i>Clinical Nutrition</i> , 2020, 39, 258-264.	2.3	12
22	Visit-to-visit variability in blood pressure and the development of chronic kidney disease in treated general hypertensive patients. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, 1739-1746.	0.4	12
23	The association between dietary energy intake and the risk of mortality in maintenance haemodialysis patients: a multi-centre prospective cohort study. <i>British Journal of Nutrition</i> , 2020, 123, 437-445.	1.2	11
24	A tetranucleotide deletion in the ANK1 gene causes hereditary spherocytosis; a case of misdiagnosis. <i>Gene</i> , 2020, 726, 144226.	1.0	6
25	Association of visit-to-visit variability in blood pressure and first stroke risk in hypertensive patients with chronic kidney disease. <i>Journal of Hypertension</i> , 2020, 38, 610-617.	0.3	13
26	Relationship between Serum Uric Acid and Mortality Risk in Hemodialysis Patients: A Multicenter Prospective Cohort Study. <i>American Journal of Nephrology</i> , 2020, 51, 823-832.	1.4	7
27	Degree of blood pressure control and the risk of new-onset hyperuricemia in treated hypertensive patients. <i>Annals of Translational Medicine</i> , 2020, 8, 1434-1434.	0.7	5
28	Interaction of neutrophil counts and folic acid treatment on new-onset proteinuria in hypertensive patients. <i>British Journal of Nutrition</i> , 2020, 126, 1-8.	1.2	2
29	Positive association of serum uric acid with new-onset diabetes in Chinese women with hypertension in a retrospective analysis of the China Stroke Primary Prevention Trial. <i>Diabetes, Obesity and Metabolism</i> , 2020, 22, 1598-1606.	2.2	8
30	Relationship of Body Mass Index and Waist Circumference With Risk of New-Onset Proteinuria in Hypertensive Patients. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020, 105, e511-e519.	1.8	6
31	Interaction of serum vitamin B ₁₂ and folate with MTHFR genotypes on risk of ischemic stroke. <i>Neurology</i> , 2020, 94, e1126-e1136.	1.5	40
32	Plasma selenium levels and risk of new-onset diabetes in hypertensive adults. <i>Journal of Trace Elements in Medicine and Biology</i> , 2019, 56, 6-12.	1.5	12
33	Comparison of three nutritional screening tools for predicting mortality in maintenance hemodialysis patients. <i>Nutrition</i> , 2019, 67-68, 110532.	1.1	16
34	Quantitative reduction in short-chain fatty acids, especially butyrate, contributes to the progression of chronic kidney disease. <i>Clinical Science</i> , 2019, 133, 1857-1870.	1.8	88
35	Positive association between baseline brachial-ankle pulse wave velocity and the risk of new-onset diabetes in hypertensive patients. <i>Cardiovascular Diabetology</i> , 2019, 18, 111.	2.7	27
36	Aldosterone Antagonists Reduce the Risk of Cardiovascular Mortality in Dialysis Patients: A Meta-Analysis. <i>Evidence-based Complementary and Alternative Medicine</i> , 2019, 2019, 1-9.	0.5	4

#	ARTICLE	IF	CITATIONS
37	A novel 223Åkb deletion in the betaâ€globin gene cluster was identified in a Chinese thalassemia major patient. International Journal of Laboratory Hematology, 2019, 41, 456-460.	0.7	9
38	Serum Lipids and Risk of Rapid Renal Function Decline in Treated Hypertensive Adults With Normal Renal Function. American Journal of Hypertension, 2019, 32, 393-401.	1.0	8
39	Impact of achieved blood pressure on renal function decline and first stroke in hypertensive patients with chronic kidney disease. Nephrology Dialysis Transplantation, 2018, 33, 409-417.	0.4	18
40	BMI is associated with the development of chronic kidney diseases in hypertensive patients with normal renal function. Journal of Hypertension, 2018, 36, 2085-2091.	0.3	18
41	Independent and combined effect of bilirubin and smoking on the progression of chronic kidney disease. Clinical Epidemiology, 2018, Volume 10, 121-132.	1.5	14
42	Folic acid therapy reduces serum uric acid in hypertensive patients: a substudy of the China Stroke Primary Prevention Trial (CSPPT). American Journal of Clinical Nutrition, 2017, 105, 882-889.	2.2	43
43	Advanced oxidation protein products promote NADPH oxidase-dependent Î²-cell destruction and dysfunction through the Bcl-2/Bax apoptotic pathway. Laboratory Investigation, 2017, 97, 792-805.	1.7	16
44	Effects of Folic Acid Therapy on the New-Onset Proteinuria in Chinese Hypertensive Patients. Hypertension, 2017, 70, 300-306.	1.3	15
45	Impact of Achieved Blood Pressure on First Stroke in Uncomplicated Grade 1 Hypertension. Journal of the American Heart Association, 2017, 6, .	1.6	6
46	Elevated Homocysteine Concentrations Decrease the Antihypertensive Effect of Angiotensin-Converting Enzyme Inhibitors in Hypertensive Patients. Arteriosclerosis, Thrombosis, and Vascular Biology, 2017, 37, 166-172.	1.1	38
47	Increased Apoptosis in the Paraventricular Nucleus Mediated by AT1R/Ras/ERK1/2 Signaling Results in Sympathetic Hyperactivity and Renovascular Hypertension in Rats after Kidney Injury. Frontiers in Physiology, 2017, 8, 41.	1.3	22
48	Efficacy of Folic Acid Therapy on the Progression of Chronic Kidney Disease. JAMA Internal Medicine, 2016, 176, 1443.	2.6	133
49	Efficacy and safety of mycophenolate mofetil treatment in IgA nephropathy: a systematic review. BMC Nephrology, 2014, 15, 193.	0.8	21
50	Renal expression of advanced oxidative protein products predicts progression of renal fibrosis in patients with IgA nephropathy. Laboratory Investigation, 2014, 94, 966-977.	1.7	13
51	Increased plasma advanced oxidation protein products is an early marker of endothelial dysfunction in type 2 diabetes patients without albuminuria	0.8	28
52	Advanced oxidation protein products decrease expression of nephrin and podocin in podocytes via ROS-dependent activation of p38 MAPK. Science China Life Sciences, 2010, 53, 68-77.	2.3	13