

# Eder Gambeta

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2691021/publications.pdf>

Version: 2024-02-01

17  
papers

402  
citations

933447

10  
h-index

888059

17  
g-index

18  
all docs

18  
docs citations

18  
times ranked

413  
citing authors

#	ARTICLE	IF	CITATIONS
1	Regulation of CaV3.2 channels by the receptor for activated C kinase 1 (Rack-1). <i>Pflugers Archiv European Journal of Physiology</i> , 2022, 474, 447-454.	2.8	7
2	A Synthetically Accessible Small-Molecule Inhibitor of USP5-Cav3.2 Calcium Channel Interactions with Analgesic Properties. <i>ACS Chemical Neuroscience</i> , 2022, 13, 524-536.	3.5	12
3	CaV3.2 calcium channels contribute to trigeminal neuralgia. <i>Pain</i> , 2022, 163, 2315-2325.	4.2	22
4	An orbitofrontal cortex to midbrain projection modulates hypersensitivity after peripheral nerve injury. <i>Cell Reports</i> , 2021, 35, 109033.	6.4	11
5	A CACNA1A variant associated with trigeminal neuralgia alters the gating of Cav2.1 channels. <i>Molecular Brain</i> , 2021, 14, 4.	2.6	11
6	Trigeminal neuralgia: An overview from pathophysiology to pharmacological treatments. <i>Molecular Pain</i> , 2020, 16, 174480692090189.	2.1	153
7	Dopamine Inputs from the Ventral Tegmental Area into the Medial Prefrontal Cortex Modulate Neuropathic Pain-Associated Behaviors in Mice. <i>Cell Reports</i> , 2020, 31, 107812.	6.4	47
8	Discovery of Michael acceptor containing 1,4-dihydropyridines as first covalent inhibitors of L-/T-type calcium channels. <i>Bioorganic Chemistry</i> , 2019, 91, 103187.	4.1	16
9	Posttraumatic stress disorder-type behaviors in streptozotocin-induced diabetic rats can be prevented by prolonged treatment with vitamin E. <i>Behavioural Brain Research</i> , 2019, 359, 749-754.	2.2	15
10	Comparison of antinociceptive effects of plain lidocaine versus lidocaine complexed with hydroxypropyl- $\beta$ -cyclodextrin in animal models of acute and persistent orofacial pain. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2019, 392, 573-583.	3.0	4
11	Anxiety- but not depressive-like behaviors are related to facial hyperalgesia in a model of trigeminal neuropathic pain in rats. <i>Physiology and Behavior</i> , 2018, 191, 131-137.	2.1	19
12	Blockade of endothelin receptors reduces tumor-induced ongoing pain and evoked hypersensitivity in a rat model of facial carcinoma induced pain. <i>European Journal of Pharmacology</i> , 2018, 818, 132-140.	3.5	13
13	Chlorpyrifos induces anxiety-like behavior in offspring rats exposed during pregnancy. <i>Neuroscience Letters</i> , 2017, 641, 94-100.	2.1	30
14	Facial pain and anxiety-like behavior are reduced by pregabalin in a model of facial carcinoma in rats. <i>Neuropharmacology</i> , 2017, 125, 263-271.	4.1	9
15	A serotonergic deficit in the dorsal periaqueductal gray matter may underpin enhanced panic-like behavior in diabetic rats. <i>Behavioural Pharmacology</i> , 2017, 28, 558-564.	1.7	3
16	Evaluation of heat hyperalgesia and anxiety like-behaviors in a rat model of orofacial cancer. <i>Neuroscience Letters</i> , 2016, 619, 100-105.	2.1	11
17	Reestablishment of the hyperglycemia to the normal levels seems not to be essential to the anxiolytic-like effect induced by insulin. <i>Metabolic Brain Disease</i> , 2016, 31, 563-571.	2.9	19