

Iordanis N Papadopoulos

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/2691696/iordanis-n-papadopoulos-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

39
papers

659
citations

18
h-index

25
g-index

41
ext. papers

755
ext. citations

3.4
avg, IF

3.9
L-index

#	Paper	IF	Citations
39	The long-term need for calcium supplementation after incidental parathyroidectomy.. <i>Journal of Taibah University Medical Sciences</i> , 2022 , 17, 214-219	1.7	
38	High Expression of a tRNAPro Derivative Associates with Poor Survival and Independently Predicts Colorectal Cancer Recurrence. <i>Biomedicines</i> , 2022 , 10, 1120	4.8	0
37	Femoral fractures are an indicator of increased severity of injury for road traffic collision victims: an autopsy-based case-control study on 4895 fatalities. <i>Archives of Orthopaedic and Trauma Surgery</i> , 2021 , 1	3.6	
36	High clusterin (CLU) mRNA expression levels in tumors of colorectal cancer patients predict a poor prognostic outcome. <i>Clinical Biochemistry</i> , 2020 , 75, 62-69	3.5	15
35	Heat shock protein beta 3 (HSPB3) is an unfavorable molecular biomarker in colorectal adenocarcinoma. <i>Molecular Carcinogenesis</i> , 2020 , 59, 116-125	5	8
34	TYMS/KRAS/BRAF molecular profiling predicts survival following adjuvant chemotherapy in colorectal cancer. <i>World Journal of Gastrointestinal Oncology</i> , 2019 , 11, 551-566	3.4	10
33	Osteoporosis-pseudoglioma syndrome: clinical, genetic, and treatment-response study of 10 new cases in Greece. <i>European Journal of Pediatrics</i> , 2019 , 178, 323-329	4.1	9
32	High microRNA-28-5p expression in colorectal adenocarcinoma predicts short-term relapse of node-negative patients and poor overall survival of patients with non-metastatic disease. <i>Clinical Chemistry and Laboratory Medicine</i> , 2018 , 56, 990-1000	5.9	20
31	miR-15a-5p, A Novel Prognostic Biomarker, Predicting Recurrent Colorectal Adenocarcinoma. <i>Molecular Diagnosis and Therapy</i> , 2017 , 21, 453-464	4.5	33
30	miR-34a overexpression predicts poor prognostic outcome in colorectal adenocarcinoma, independently of clinicopathological factors with established prognostic value. <i>Clinical Biochemistry</i> , 2017 , 50, 918-924	3.5	18
29	Elevated expression of miR-24-3p is a potentially adverse prognostic factor in colorectal adenocarcinoma. <i>Clinical Biochemistry</i> , 2017 , 50, 285-292	3.5	27
28	Upregulated miR-16 expression is an independent indicator of relapse and poor overall survival of colorectal adenocarcinoma patients. <i>Clinical Chemistry and Laboratory Medicine</i> , 2017 , 55, 737-747	5.9	25
27	Early Diagnosis and Surgical Treatment for Necrotizing Fasciitis: A Multicenter Study. <i>Frontiers in Surgery</i> , 2017 , 4, 5	2.3	38
26	High miR-96 levels in colorectal adenocarcinoma predict poor prognosis, particularly in patients without distant metastasis at the time of initial diagnosis. <i>Tumor Biology</i> , 2016 , 37, 11815-11824	2.9	36
25	miR-224 overexpression is a strong and independent prognosticator of short-term relapse and poor overall survival in colorectal adenocarcinoma. <i>International Journal of Oncology</i> , 2015 , 46, 849-59	4.4	32
24	The effect of metronomic versus standard chemotherapy on the regulatory to effector T-cell equilibrium in cancer patients. <i>Experimental Hematology and Oncology</i> , 2014 , 3, 3	7.8	18
23	Kallikrein-related peptidase-6 (KLK6) mRNA expression is an independent prognostic tissue biomarker of poor disease-free and overall survival in colorectal adenocarcinoma. <i>Tumor Biology</i> , 2014 , 35, 4673-85	2.9	28

22	KLK11 mRNA expression predicts poor disease-free and overall survival in colorectal adenocarcinoma patients. <i>Biomarkers in Medicine</i> , 2014 , 8, 671-85	2.3	23
21	Acute abdominal pain due to intussusception of the small intestine in a 17-year-old patient. <i>Surgery</i> , 2014 , 156, 1282-3	3.6	
20	Enhanced miR-182 transcription is a predictor of poor overall survival in colorectal adenocarcinoma patients. <i>Clinical Chemistry and Laboratory Medicine</i> , 2014 , 52, 1217-27	5.9	31
19	Clinical significance of kallikrein-related peptidase (KLK10) mRNA expression in colorectal cancer. <i>Clinical Biochemistry</i> , 2013 , 46, 1453-61	3.5	37
18	Kallikrein-related peptidase 4 (KLK4) mRNA predicts short-term relapse in colorectal adenocarcinoma patients. <i>Cancer Letters</i> , 2013 , 330, 106-12	9.9	28
17	L-Dopa decarboxylase (DDC) constitutes an emerging biomarker in predicting patients' survival with stomach adenocarcinomas. <i>Journal of Cancer Research and Clinical Oncology</i> , 2013 , 139, 297-306	4.9	6
16	A structured autopsy-based audit of 370 firearm fatalities: Contribution to inform policy decisions and the probability of the injured arriving alive at a hospital and receiving definitive care. <i>Accident Analysis and Prevention</i> , 2013 , 50, 667-77	6.1	11
15	Alcohol and psychoactive drugs increased the pre-hospital mortality in 655 fall-related fatalities in Greece: a call for management protocols. <i>Injury</i> , 2012 , 43, 1522-6	2.5	8
14	Asymptomatic omental granuloma following spillage of gallstones during laparoscopic cholecystectomy protects patients and influences surgeons' decisions: a review. <i>BMJ Case Reports</i> , 2012 , 2012,	0.9	5
13	Enterocolitis with multiple ulcers of ileum and right colon in a patient with leukaemia attributed to cytomegalovirus. <i>BMJ Case Reports</i> , 2012 , 2012,	0.9	1
12	Patients with pelvic fractures due to falls: A paradigm that contributed to autopsy-based audit of trauma in Greece. <i>Journal of Trauma Management and Outcomes</i> , 2011 , 5, 2		5
11	Perforated mucinous cystadenoma of the vermiform appendix: an overview in reasoning clinical decisions. <i>BMJ Case Reports</i> , 2011 , 2011,	0.9	2
10	Motor vehicle collision fatalities involving alcohol and illicit drugs in Greece: the need for management protocols and a reassessment of surveillance. <i>Addiction</i> , 2010 , 105, 1952-61	4.6	6
9	Primary epithelial splenic cyst with micro-rupture and raised carbohydrate antigen CA 19-9: a paradigm of management. <i>BMJ Case Reports</i> , 2010 , 2010,	0.9	2
8	Molecular analysis and prognostic impact of the novel apoptotic gene BCL2L12 in gastric cancer. <i>Biochemical and Biophysical Research Communications</i> , 2010 , 391, 214-8	3.4	29
7	Kallikrein-related peptidase 13 (KLK13) gene expressional status contributes significantly in the prognosis of primary gastric carcinomas. <i>Clinical Biochemistry</i> , 2010 , 43, 1205-11	3.5	21
6	Primary epithelial splenic cyst with micro-rupture and raised carbohydrate antigen CA 19-9: a paradigm of management. <i>BMJ Case Reports</i> , 2010 , 2010,	0.9	4
5	Fibroepithelial polyp in an anal fistulous track: a sign of chronic pathology. <i>BMJ Case Reports</i> , 2010 , 2010,	0.9	1

4	Status and perspectives of hospital mortality in a public urban Hellenic hospital, based on a five-year review. <i>BMC Public Health</i> , 2008 , 8, 28	4.1	10
3	Quantitative expression analysis and prognostic significance of the novel apoptosis-related gene BCL2L12 in colon cancer. <i>Biological Chemistry</i> , 2008 , 389, 1467-75	4.5	36
2	Auditing 655 fatalities with pelvic fractures by autopsy as a basis to evaluate trauma care. <i>Journal of the American College of Surgeons</i> , 2006 , 203, 30-43	4.4	74
1	Delayed reaction to the Dacron buttress used in Stamey bladder neck suspension. <i>International Urology and Nephrology</i> , 2006 , 38, 269-72	2.3	2