

# Huan-Xin Lin

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2692201/publications.pdf>

Version: 2024-02-01

80  
papers

1,771  
citations

304602

22  
h-index

345118

36  
g-index

84  
all docs

84  
docs citations

84  
times ranked

2950  
citing authors

#	ARTICLE	IF	CITATIONS
1	Overexpression of GOLPH3 Promotes Proliferation and Tumorigenicity in Breast Cancer via Suppression of the FOXO1 Transcription Factor. <i>Clinical Cancer Research</i> , 2012, 18, 4059-4069.	3.2	129
2	Effect of Capecitabine Maintenance Therapy Using Lower Dosage and Higher Frequency vs Observation on Disease-Free Survival Among Patients With Early-Stage Triple-Negative Breast Cancer Who Had Received Standard Treatment. <i>JAMA - Journal of the American Medical Association</i> , 2021, 325, 50.	3.8	113
3	Serum levels of CEA and CA15-3 in different molecular subtypes and prognostic value in Chinese breast cancer. <i>Breast</i> , 2014, 23, 88-93.	0.9	90
4	Thymosin beta 10 is a key regulator of tumorigenesis and metastasis and a novel serum marker in breast cancer. <i>Breast Cancer Research</i> , 2017, 19, 15.	2.2	89
5	Tumor cells PD-L1 expression as a favorable prognosis factor in nasopharyngeal carcinoma patients with pre-existing intratumor-infiltrating lymphocytes. <i>Oncolmmunology</i> , 2017, 6, e1312240.	2.1	68
6	Comparison of clinical outcomes of squamous cell carcinoma, adenocarcinoma, and adenosquamous carcinoma of the uterine cervix after definitive radiotherapy: a population-based analysis. <i>Journal of Cancer Research and Clinical Oncology</i> , 2017, 143, 115-122.	1.2	59
7	Upregulation of E2F8 promotes cell proliferation and tumorigenicity in breast cancer by modulating G1/S phase transition. <i>Oncotarget</i> , 2016, 7, 23757-23771.	0.8	46
8	The Value of Prognostic Nutritional Index (PNI) in Predicting Survival and Guiding Radiotherapy of Patients With T1-2N1 Breast Cancer. <i>Frontiers in Oncology</i> , 2019, 9, 1562.	1.3	45
9	Decreased Expression of Beclin 1 Correlates Closely with Bcl-xL Expression and Poor Prognosis of Ovarian Carcinoma. <i>PLoS ONE</i> , 2013, 8, e60516.	1.1	44
10	Optimal cumulative cisplatin dose in nasopharyngeal carcinoma patients based on induction chemotherapy response. <i>Radiotherapy and Oncology</i> , 2019, 137, 83-94.	0.3	44
11	TIMELESS confers cisplatin resistance in nasopharyngeal carcinoma by activating the Wnt/ $\beta$ -catenin signaling pathway and promoting the epithelial mesenchymal transition. <i>Cancer Letters</i> , 2017, 402, 117-130.	3.2	42
12	Prognostic Value of Preoperative Systemic Immune-Inflammation Index in Breast Cancer: A Propensity Score-Matching Study. <i>Frontiers in Oncology</i> , 2020, 10, 580.	1.3	38
13	The preoperative systemic inflammation response index (SIRI) independently predicts survival in postmenopausal women with breast cancer. <i>Current Problems in Cancer</i> , 2020, 44, 100560.	1.0	34
14	ALG3 contributes to stemness and radioresistance through regulating glycosylation of TGF- $\beta$ 2 receptor II in breast cancer. <i>Journal of Experimental and Clinical Cancer Research</i> , 2021, 40, 149.	3.5	34
15	Prognostic Value of Ki-67 in Breast Cancer Patients with Positive Axillary Lymph Nodes: A Retrospective Cohort Study. <i>PLoS ONE</i> , 2014, 9, e87264.	1.1	33
16	Overexpression of Kinesin Family Member 20A Correlates with Disease Progression and Poor Prognosis in Human Nasopharyngeal Cancer: A Retrospective Analysis of 105 Patients. <i>PLoS ONE</i> , 2017, 12, e0169280.	1.1	32
17	Surgery Combined with Radiotherapy Improved Survival in Metastatic Esophageal Cancer in a Surveillance Epidemiology and End Results Population-based Study. <i>Scientific Reports</i> , 2016, 6, 28280.	1.6	31
18	Tumor location is a prognostic factor for survival of Chinese women with T1-2N0M0 breast cancer. <i>International Journal of Surgery</i> , 2014, 12, 394-398.	1.1	28

#	ARTICLE	IF	CITATIONS
19	Identification of two microRNA signatures in whole blood as novel biomarkers for diagnosis of nasopharyngeal carcinoma. <i>Journal of Translational Medicine</i> , 2019, 17, 186.	1.8	27
20	Prognosis of patients with esophageal squamous cell carcinoma after esophagectomy using the log odds of positive lymph nodes. <i>Oncotarget</i> , 2015, 6, 36911-36922.	0.8	26
21	Pan-Immune-Inflammation Value: A New Prognostic Index in Operative Breast Cancer. <i>Frontiers in Oncology</i> , 2022, 12, 830138.	1.3	26
22	Randomized study of sinusoidal chronomodulated versus flat intermittent induction chemotherapy with cisplatin and 5-fluorouracil followed by traditional radiotherapy for locoregionally advanced nasopharyngeal carcinoma. <i>Chinese Journal of Cancer</i> , 2013, 32, 502-511.	4.9	24
23	The diagnostic and prognostic values of plasma Epstein-Barr virus DNA for residual cervical lymphadenopathy in nasopharyngeal carcinoma patients: a retrospective study. <i>Cancer Communications</i> , 2019, 39, 1-13.	3.7	24
24	Use of CEA and CA15-3 to Predict Axillary Lymph Node Metastasis in Patients with Breast Cancer. <i>Journal of Cancer</i> , 2016, 7, 37-41.	1.2	23
25	Patterns of distant metastasis in Chinese women according to breast cancer subtypes. <i>Oncotarget</i> , 2016, 7, 47975-47984.	0.8	23
26	Prognostic value of lymph node ratio in stage IIIC epithelial ovarian cancer with node-positive in a SEER population-based study. <i>Oncotarget</i> , 2016, 7, 7952-7959.	0.8	22
27	Patterns of Regional Lymph Node Recurrence After Radical Surgery for Thoracic Esophageal Squamous Cell Carcinoma. <i>Annals of Thoracic Surgery</i> , 2016, 101, 551-557.	0.7	22
28	Postmastectomy Radiotherapy Improves Disease-Free Survival of High Risk of Locoregional Recurrence Breast Cancer Patients with T1-2 and 1 to 3 Positive Nodes. <i>PLoS ONE</i> , 2015, 10, e0119105.	1.1	22
29	Adjuvant radiation therapy and survival for adenoid cystic carcinoma of the breast. <i>Breast</i> , 2017, 31, 214-218.	0.9	21
30	Sarcopenia is associated with higher toxicity and poor prognosis of nasopharyngeal carcinoma. <i>Therapeutic Advances in Medical Oncology</i> , 2020, 12, 175883592094761.	1.4	21
31	Using the Lymph Node Ratio to Evaluate the Prognosis of Stage II/III Breast Cancer Patients Who Received Neoadjuvant Chemotherapy and Mastectomy. <i>Cancer Research and Treatment</i> , 2015, 47, 757-764.	1.3	20
32	Number of negative lymph nodes should be considered for incorporation into staging for breast cancer. <i>American Journal of Cancer Research</i> , 2015, 5, 844-53.	1.4	20
33	Prognostic Value of Different Lymph Node Staging Methods in Esophageal Squamous Cell Carcinoma After Esophagectomy. <i>Annals of Thoracic Surgery</i> , 2015, 99, 284-290.	0.7	19
34	Efficacy of controlled-release oxycodone for reducing pain due to oral mucositis in nasopharyngeal carcinoma patients treated with concurrent chemoradiotherapy: a prospective clinical trial. <i>Supportive Care in Cancer</i> , 2019, 27, 3759-3767.	1.0	18
35	The local treatment modalities in FIGO stage I-II small cell carcinoma of the cervix are determined by disease stage and lymph node status. <i>Cancer Medicine</i> , 2016, 5, 1108-1115.	1.3	17
36	Patients with Old Age or Proximal Tumors Benefit from Metabolic Syndrome in Early Stage Gastric Cancer. <i>PLoS ONE</i> , 2014, 9, e89965.	1.1	17

#	ARTICLE	IF	CITATIONS
37	Clinicopathological features of small cell carcinoma of the uterine cervix in the surveillance, epidemiology, and end results database. <i>Oncotarget</i> , 2017, 8, 40425-40433.	0.8	17
38	Dosimetric analysis of the brachial plexus among patients with breast cancer treated with post-mastectomy radiotherapy to the ipsilateral supraclavicular area: report of 3 cases of radiation-induced brachial plexus neuropathy. <i>Radiation Oncology</i> , 2014, 9, 292.	1.2	16
39	Lymph node ratio may predict the benefit of postoperative radiotherapy in node-positive cervical cancer. <i>Oncotarget</i> , 2016, 7, 29420-29428.	0.8	16
40	A prognostic survival model based on metabolism-related gene expression in plasma cell myeloma. <i>Leukemia</i> , 2021, 35, 3212-3222.	3.3	15
41	The putative tumor activator ARHGEF3 promotes nasopharyngeal carcinoma cell pathogenesis by inhibiting cellular apoptosis. <i>Oncotarget</i> , 2016, 7, 25836-25848.	0.8	15
42	Impact of the number of resected lymph nodes on survival after preoperative radiotherapy for esophageal cancer. <i>Oncotarget</i> , 2016, 7, 22497-22507.	0.8	14
43	Overexpression of acylglycerol kinase is associated with poorer prognosis and lymph node metastasis in nasopharyngeal carcinoma. <i>Tumor Biology</i> , 2016, 37, 3349-3357.	0.8	14
44	The effect of local treatment modalities in patients with early-stage adenocarcinoma of the uterine cervix: A population-based analysis. <i>International Journal of Surgery</i> , 2017, 41, 16-22.	1.1	14
45	Prostate Tumor Overexpressed 1 (PTOV1) Is a Novel Prognostic Marker for Nasopharyngeal Carcinoma Progression and Poor Survival Outcomes. <i>PLoS ONE</i> , 2015, 10, e0136448.	1.1	14
46	Comparable Survival between Additional Radiotherapy and Local Surgery in Occult Breast Cancer after Axillary Lymph Node Dissection: A Population-based Analysis. <i>Journal of Cancer</i> , 2017, 8, 3849-3855.	1.2	13
47	Liposomal paclitaxel versus docetaxel in induction chemotherapy using Taxanes, cisplatin and 5-fluorouracil for locally advanced nasopharyngeal carcinoma. <i>BMC Cancer</i> , 2018, 18, 1279.	1.1	13
48	A Novel Inflammatory-Nutritional Prognostic Scoring System for Patients with Early-Stage Breast Cancer. <i>Journal of Inflammation Research</i> , 2022, Volume 15, 381-394.	1.6	13
49	Number of negative lymph nodes can predict survival of breast cancer patients with four or more positive lymph nodes after postmastectomy radiotherapy. <i>Radiation Oncology</i> , 2014, 9, 284.	1.2	12
50	Differences in esophageal cancer characteristics and survival between Chinese and Caucasian patients in the SEER database. <i>OncoTargets and Therapy</i> , 2016, Volume 9, 6435-6444.	1.0	12
51	Prognostic value of skeletal muscle index and monocyte-to-lymphocyte ratio for lymph node-positive breast cancer patients after mastectomy. <i>Annals of Translational Medicine</i> , 2019, 7, 775-775.	0.7	12
52	Prognostic significance of the skeletal muscle index and an inflammation biomarker in patients with breast cancer who underwent postoperative adjuvant radiotherapy. <i>Current Problems in Cancer</i> , 2020, 44, 100513.	1.0	12
53	Incorporation of the number of positive lymph nodes leads to better prognostic discrimination of node-positive early stage cervical cancer. <i>Oncotarget</i> , 2017, 8, 26057-26065.	0.8	12
54	Number of negative lymph nodes is associated with disease-free survival in patients with breast cancer. <i>BMC Cancer</i> , 2015, 15, 43.	1.1	10

#	ARTICLE	IF	CITATIONS
55	The survival benefits of local surgery in stage IV breast cancer are not affected by breast cancer subtypes: a population-based analysis. <i>Oncotarget</i> , 2017, 8, 67851-67860.	0.8	10
56	Progesterone receptor loss identifies hormone receptor-positive and HER2-negative breast cancer subgroups at higher risk of relapse: a retrospective cohort study. <i>OncoTargets and Therapy</i> , 2016, 9, 1707.	1.0	9
57	21-Gene Recurrence Score Assay Could Not Predict Benefit of Post-mastectomy Radiotherapy in T1-2 N1mic ER-Positive HER2-Negative Breast Cancer. <i>Frontiers in Oncology</i> , 2019, 9, 270.	1.3	8
58	Comparing three induction chemotherapy regimens for patients with locoregionally advanced nasopharyngeal carcinoma based on TNM stage and plasma Epstein-Barr virus DNA level. <i>BMC Cancer</i> , 2020, 20, 89.	1.1	8
59	Optimal cumulative cisplatin dose in nasopharyngeal carcinoma patients based on plasma Epstein-Barr virus DNA level after induction chemotherapy. <i>Aging</i> , 2020, 12, 4931-4944.	1.4	8
60	Ovarian Ablation Using Goserelin Improves Survival of Premenopausal Patients with Stage II/III Hormone Receptor-Positive Breast Cancer without Chemotherapy-Induced Amenorrhea. <i>Cancer Research and Treatment</i> , 1970, 47, 55-63.	1.3	8
61	Effect of blood type on survival of Chinese patients with esophageal squamous cell carcinoma. <i>OncoTargets and Therapy</i> , 2015, 8, 947.	1.0	7
62	Clinicopathological characteristics, treatment, and survival outcomes of cystadenocarcinoma of the salivary gland: a population-based study. <i>OncoTargets and Therapy</i> , 2016, Volume 9, 6569-6572.	1.0	7
63	Therapeutic role of axillary lymph node dissection in patients with stage IV breast cancer: a population-based analysis. <i>Journal of Cancer Research and Clinical Oncology</i> , 2017, 143, 467-474.	1.2	7
64	The effect of postmastectomy radiotherapy in node-positive triple-negative breast cancer. <i>BMC Cancer</i> , 2020, 20, 1146.	1.1	7
65	The value of radiotherapy in breast cancer patients with isolated ipsilateral supraclavicular lymph node metastasis without distant metastases at diagnosis: a retrospective analysis of Chinese patients. <i>OncoTargets and Therapy</i> , 2014, 7, 281.	1.0	6
66	Lymph node dissection improved survival in patients with metastatic thoracic esophageal cancer: An analysis of 220 patients from the SEER database. <i>International Journal of Surgery</i> , 2016, 35, 13-18.	1.1	6
67	Clinicopathologic characteristics and clinical outcomes of pure type and mixed type of tubular carcinoma of the breast: a single-institution cohort study. <i>Cancer Management and Research</i> , 2018, Volume 10, 4509-4515.	0.9	6
68	Nomogram Predicting the Benefits of Adding Concurrent Chemotherapy to Intensity-Modulated Radiotherapy After Induction Chemotherapy in Stages IIa-IVb Nasopharyngeal Carcinoma. <i>Frontiers in Oncology</i> , 2020, 10, 539321.	1.3	6
69	Low Skeletal Muscle Mass Impairs Quality of Life in Nasopharyngeal Carcinoma Patients Treated With Concurrent Chemoradiotherapy. <i>Frontiers in Nutrition</i> , 2019, 6, 195.	1.6	5
70	Establishment and validation of a prognostic nomogram to predict early metastasis in nasopharyngeal carcinoma patients within six months after radiotherapy and to guide intensive treatment. <i>Radiotherapy and Oncology</i> , 2021, 162, 202-211.	0.3	5
71	Clinicopathologic features and treatment of breast metastasis from nasopharyngeal carcinoma: A report of two cases and literature review. <i>Oncology Letters</i> , 2015, 10, 3675-3681.	0.8	4
72	Effect of postoperative radiotherapy for squamous cell cancer of the breast in a surveillance epidemiology and end results population-based study. <i>Oncotarget</i> , 2016, 7, 10684-10693.	0.8	4

#	ARTICLE	IF	CITATIONS
73	Number of Negative Lymph Nodes Can Predict Survival after Postmastectomy Radiotherapy According to Different Breast Cancer Subtypes. <i>Journal of Cancer</i> , 2015, 6, 261-269.	1.2	3
74	The prognosis of neck residue nasopharyngeal carcinoma (NPC) patients: results from a case-cohort study. <i>Journal of Cancer</i> , 2018, 9, 1765-1772.	1.2	3
75	Prognostic value of lymph node ratio in patients with small-cell carcinoma of the cervix based on data from a large national registry. <i>OncoTargets and Therapy</i> , 2015, 9, 67.	1.0	2
76	Establishment of prognostic nomograms based on skeletal muscle index and serum biomarker in breast cancer patients receiving radiotherapy. <i>Clinical and Translational Medicine</i> , 2020, 10, e115.	1.7	2
77	Accelerated Partial Breast Irradiation with Intensity-Modulated Radiotherapy Is Feasible for Chinese Breast Cancer Patients. <i>Journal of Breast Cancer</i> , 2014, 17, 256.	0.8	1
78	A Model Combining Skeletal Muscle Mass and a Hematological Biomarker to Predict Survival in Patients With Nasopharyngeal Carcinoma Undergoing Concurrent Chemoradiotherapy. <i>Frontiers in Oncology</i> , 2021, 11, 644676.	1.3	1
79	Systematic Construction of an Autophagic Risk Model in Bone Marrow for Prognostic Prediction in Multiple Myeloma. <i>Blood</i> , 2021, 138, 4713-4713.	0.6	0
80	An Immune Score Model of the Bone Marrow Micro-Environment Predicts Survival in Chronic Lymphocytic Leukaemia. <i>Blood</i> , 2020, 136, 38-38.	0.6	0