

Victor Ortiz-García-a de la Foz

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2694945/publications.pdf>

Version: 2024-02-01

83
papers

1,747
citations

304743

22
h-index

345221

36
g-index

90
all docs

90
docs citations

90
times ranked

3000
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 1 | Cortical thickness across the lifespan: Data from 17,075 healthy individuals aged 3â€“90â€“years. Human Brain Mapping, 2022, 43, 431-451. | 3.6 | 143 |
| 2 | Virtual Histology of Cortical Thickness and Shared Neurobiology in 6 Psychiatric Disorders. JAMA Psychiatry, 2021, 78, 47. | 11.0 | 136 |
| 3 | Predicting relapse after a first episode of non-affective psychosis: A three-year follow-up study. Journal of Psychiatric Research, 2012, 46, 1099-1105. | 3.1 | 118 |
| 4 | Using Machine Learning and Structural Neuroimaging to Detect First Episode Psychosis: Reconsidering the Evidence. Schizophrenia Bulletin, 2020, 46, 17-26. | 4.3 | 76 |
| 5 | Genetic variants associated with longitudinal changes in brain structure across the lifespan. Nature Neuroscience, 2022, 25, 421-432. | 14.8 | 75 |
| 6 | Subcortical volumes across the lifespan: Data from 18,605 healthy individuals aged 3â€“90â€“years. Human Brain Mapping, 2022, 43, 452-469. | 3.6 | 72 |
| 7 | Towards Precision Medicine in Psychosis: Benefits and Challenges of Multimodal Multicenter Studiesâ€“PSYSCAN: Translating Neuroimaging Findings From Research into Clinical Practice. Schizophrenia Bulletin, 2020, 46, 432-441. | 4.3 | 56 |
| 8 | Neuroharmony: A new tool for harmonizing volumetric MRI data from unseen scanners. NeuroImage, 2020, 220, 117127. | 4.2 | 48 |
| 9 | Clinical Outcome After Antipsychotic Treatment Discontinuation in Functionally Recovered First-Episode Nonaffective Psychosis Individuals. Journal of Clinical Psychiatry, 2016, 77, 492-500. | 2.2 | 47 |
| 10 | Incidence of non-alcoholic fatty liver disease and metabolic dysfunction in first episode schizophrenia and related psychotic disorders: a 3-year prospective randomized interventional study. Psychopharmacology, 2016, 233, 3947-3952. | 3.1 | 40 |
| 11 | Comparison of the anti-inflammatory effect of aripiprazole and risperidone in 75 drug-naïve first episode psychosis individuals: A 3â€“months randomized study. Schizophrenia Research, 2018, 202, 226-233. | 2.0 | 37 |
| 12 | Understanding sex differences in long-term outcomes after a first episode of psychosis. NPJ Schizophrenia, 2020, 6, 33. | 3.6 | 36 |
| 13 | Prediction of acute clinical response following a first episode of non affective psychosis: Results of a cohort of 375 patients from the Spanish PAFIP study. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2013, 44, 162-167. | 4.8 | 35 |
| 14 | Comparison of metabolic effects of aripiprazole, quetiapine and ziprasidone after 12weeks of treatment in first treated episode of psychosis. Schizophrenia Research, 2014, 159, 90-94. | 2.0 | 34 |
| 15 | Brain structural and clinical changes after first episode psychosis: Focus on cannabinoid receptor 1 polymorphisms. Psychiatry Research - Neuroimaging, 2015, 233, 112-119. | 1.8 | 34 |
| 16 | Tobacco smoking and its association with cognition in first episode psychosis patients. Schizophrenia Research, 2018, 192, 269-273. | 2.0 | 34 |
| 17 | Rates and predictors of relapse in first-episode non-affective psychosis: a 3-year longitudinal study in a specialized intervention program (PAFIP). European Archives of Psychiatry and Clinical Neuroscience, 2017, 267, 315-323. | 3.2 | 32 |
| 18 | Long-term metabolic effects of aripiprazole, ziprasidone and quetiapine: a pragmatic clinical trial in drug-naïve patients with a first-episode of non-affective psychosis. Psychopharmacology, 2018, 235, 245-255. | 3.1 | 30 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Effect of antipsychotic drugs on cortical thickness. A randomized controlled one-year follow-up study of haloperidol, risperidone and olanzapine. <i>Schizophrenia Research</i> , 2012, 141, 22-28. | 2.0 | 28 |
| 20 | Incidence and risk factors of acute akathisia in 493 individuals with first episode non-affective psychosis: a 6-week randomised study of antipsychotic treatment. <i>Psychopharmacology</i> , 2017, 234, 2563-2570. | 3.1 | 27 |
| 21 | Long-term (3-year) effectiveness of haloperidol, risperidone and olanzapine: results of a randomized, flexible-dose, open-label comparison in first-episode nonaffective psychosis. <i>Psychopharmacology</i> , 2012, 219, 225-233. | 3.1 | 24 |
| 22 | Three-year longitudinal population-based volumetric MRI study in first-episode schizophrenia spectrum patients. <i>Psychological Medicine</i> , 2014, 44, 1591-1604. | 4.5 | 23 |
| 23 | Stopping cannabis use benefits outcome in psychosis: findings from 10-year follow-up study in the PAFIP cohort. <i>Acta Psychiatrica Scandinavica</i> , 2019, 140, 349-359. | 4.5 | 23 |
| 24 | Neuroanatomical abnormalities in first-episode psychosis across independent samples: a multi-centre mega-analysis. <i>Psychological Medicine</i> , 2021, 51, 340-350. | 4.5 | 23 |
| 25 | Plasma prolactin levels are associated with the severity of illness in drug-naive first-episode psychosis female patients. <i>Archives of Women's Mental Health</i> , 2019, 22, 367-373. | 2.6 | 22 |
| 26 | Aripiprazole, Ziprasidone and Quetiapine in the treatment of first-episode nonaffective psychosis: A 12-week randomized, flexible-dose, open-label trial. <i>Schizophrenia Research</i> , 2013, 147, 375-382. | 2.0 | 21 |
| 27 | Treatment of first-episode non-affective psychosis: a randomized comparison of aripiprazole, quetiapine and ziprasidone over 1 year. <i>Psychopharmacology</i> , 2014, 231, 357-366. | 3.1 | 21 |
| 28 | Aripiprazole, Ziprasidone, and Quetiapine in the Treatment of First-Episode Nonaffective Psychosis. <i>Journal of Clinical Psychopharmacology</i> , 2013, 33, 215-220. | 1.4 | 20 |
| 29 | Effects of aripiprazole, quetiapine and ziprasidone on plasma prolactin levels in individuals with first episode nonaffective psychosis: Analysis of a randomized open-label 1 year study. <i>Schizophrenia Research</i> , 2017, 189, 134-141. | 2.0 | 20 |
| 30 | Cannabis consumption and non-alcoholic fatty liver disease. A three years longitudinal study in first episode non-affective psychosis patients. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2019, 95, 109677. | 4.8 | 20 |
| 31 | Sex-specific variation of MRI-based cortical morphometry in adult healthy volunteers: The effect on cognitive functioning. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2011, 35, 616-623. | 4.8 | 19 |
| 32 | The effect of excess weight on circulating inflammatory cytokines in drug-naïve first-episode psychosis individuals. <i>Journal of Neuroinflammation</i> , 2018, 15, 63. | 7.2 | 19 |
| 33 | A 3-year prospective study on the metabolic effect of aripiprazole, quetiapine and ziprasidone: A pragmatic clinical trial in first episode psychosis patients. <i>European Neuropsychopharmacology</i> , 2020, 39, 46-55. | 0.7 | 19 |
| 34 | Effect of cannabis on weight and metabolism in first-episode non-affective psychosis: Results from a three-year longitudinal study. <i>Journal of Psychopharmacology</i> , 2019, 33, 284-294. | 4.0 | 18 |
| 35 | Duration of active psychosis and functional outcomes in first-episode non-affective psychosis. <i>European Psychiatry</i> , 2018, 52, 29-37. | 0.2 | 17 |
| 36 | No progression of the alterations in the cortical thickness of individuals with schizophrenia-spectrum disorder: a three-year longitudinal magnetic resonance imaging study of first-episode patients. <i>Psychological Medicine</i> , 2015, 45, 2861-2871. | 4.5 | 16 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | A cross-sectional and longitudinal structural magnetic resonance imaging study of the post-central gyrus in first-episode schizophrenia patients. <i>Psychiatry Research - Neuroimaging</i> , 2015, 231, 42-49. | 1.8 | 16 |
| 38 | Long-Term Antipsychotic Effectiveness in First Episode of Psychosis: A 3-Year Follow-Up Randomized Clinical Trial Comparing Aripiprazole, Quetiapine, and Ziprasidone. <i>International Journal of Neuropsychopharmacology</i> , 2018, 21, 1090-1101. | 2.1 | 16 |
| 39 | 10Kin1day: A Bottom-Up Neuroimaging Initiative. <i>Frontiers in Neurology</i> , 2019, 10, 425. | 2.4 | 15 |
| 40 | Dissecting the functional outcomes of first episode schizophrenia spectrum disorders: a 10-year follow-up study in the PAFIP cohort. <i>Psychological Medicine</i> , 2021, 51, 264-277. | 4.5 | 15 |
| 41 | Brain grey matter abnormalities in first episode non-affective psychosis patients with suicidal behaviours: The role of neurocognitive functioning. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2020, 102, 109948. | 4.8 | 14 |
| 42 | Treatment Discontinuation Impact on Long-Term (10-Year) Weight Gain and Lipid Metabolism in First-Episode Psychosis: Results From the PAFIP-10 Cohort. <i>International Journal of Neuropsychopharmacology</i> , 2021, 24, 1-7. | 2.1 | 12 |
| 43 | Understanding the direct and indirect costs of a first episode of psychosis program: Insights from PAFIP of Cantabria, Spain, during the first year of intervention. <i>Microbial Biotechnology</i> , 2019, 13, 1182-1190. | 1.7 | 11 |
| 44 | Duration of active psychosis during early phases of the illness and functional outcome: The PAFIP 10-year follow-up study. <i>Schizophrenia Research</i> , 2020, 220, 240-247. | 2.0 | 11 |
| 45 | One year longitudinal study of the straight gyrus morphometry in first-episode schizophrenia-spectrum patients. <i>Psychiatry Research - Neuroimaging</i> , 2012, 202, 80-83. | 1.8 | 9 |
| 46 | Progressive Structural Brain Changes and NRG1 Gene Variants in First-Episode Nonaffective Psychosis. <i>Neuropsychobiology</i> , 2015, 71, 103-111. | 1.9 | 9 |
| 47 | Comparison of aripiprazole and risperidone effectiveness in first episode non-affective psychosis: Rationale and design of a prospective, randomized, 3-phase, investigator-initiated study (PAFIP-3). <i>Revista De Psiquiatría Y Salud Mental</i> , 2021, 14, 157-163. | 1.8 | 9 |
| 48 | The effects of tobacco smoking on age of onset of psychosis and psychotic symptoms in a first-episode psychosis population. <i>Addiction</i> , 2017, 112, 526-532. | 3.3 | 8 |
| 49 | Long term cortical thickness changes after a first episode of non-affective psychosis: The 10-year follow-up of the PAFIP cohort. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2021, 108, 110180. | 4.8 | 8 |
| 50 | Education and long-term outcomes in first episode psychosis: 10-year follow-up study of the PAFIP cohort. <i>Psychological Medicine</i> , 2023, 53, 66-77. | 4.5 | 8 |
| 51 | Active psychosis and pro-inflammatory cytokines in first-episode of psychosis. <i>Journal of Psychiatric Research</i> , 2021, 134, 150-157. | 3.1 | 7 |
| 52 | Aripiprazole vs Risperidone for the acute-phase treatment of first-episode psychosis: A 6-week randomized, flexible-dose, open-label clinical trial. <i>European Neuropsychopharmacology</i> , 2021, 47, 74-85. | 0.7 | 7 |
| 53 | BDNF Val66Met variants and brain volume changes in non-affective psychosis patients and healthy controls: A 3-year follow-up study. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2013, 45, 201-206. | 4.8 | 6 |
| 54 | Age of onset of Cannabis use and cognitive function in first-episode non-affective psychosis patients: Outcome at three-year follow-up. <i>Schizophrenia Research</i> , 2018, 201, 159-166. | 2.0 | 6 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Expression and Functionality Study of 9 Toll-Like Receptors in 33 Drug-Naïve Non-Affective First Episode Psychosis Individuals: A 3-Month Study. <i>International Journal of Molecular Sciences</i> , 2020, 21, 6106. | 4.1 | 6 |
| 56 | Classification of first-episode psychosis using cortical thickness: A large multicenter MRI study. <i>European Neuropsychopharmacology</i> , 2021, 47, 34-47. | 0.7 | 6 |
| 57 | Prolactin, metabolic and immune parameters in naïve subjects with a first episode of psychosis. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2021, 110, 110332. | 4.8 | 6 |
| 58 | A family study on first episode of psychosis patients: Exploring neuropsychological performance as an endophenotype. <i>Acta Psychiatrica Scandinavica</i> , 2022, 145, 384-396. | 4.5 | 6 |
| 59 | Aripiprazole and Risperidone Present Comparable Long-Term Metabolic Profiles: Data From a Pragmatic Randomized Controlled Trial in Drug-Naïve First-Episode Psychosis. <i>International Journal of Neuropsychopharmacology</i> , 2022, 25, 795-806. | 2.1 | 5 |
| 60 | Straight gyrus morphology in first-episode schizophrenia-spectrum patients. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2011, 35, 84-90. | 4.8 | 4 |
| 61 | Comparison of aripiprazole and risperidone effectiveness in first episode non-affective psychosis: Rationale and design of a prospective, randomized, 3-phase, investigator-initiated study (PAFIP-3). <i>Revista De Psiquiatría Y Salud Mental (English Edition)</i> , 2021, 14, 157-163. | 0.3 | 4 |
| 62 | Predictors of diagnostic stability in brief psychotic disorders: Findings from a 3-year longitudinal study. <i>Acta Psychiatrica Scandinavica</i> , 2021, 144, 578-588. | 4.5 | 4 |
| 63 | Data regarding the effect of cannabis consumption on liver function in the prospective PAFIP cohort of first episode psychosis. <i>Data in Brief</i> , 2019, 27, 104415. | 1.0 | 3 |
| 64 | Cognitive reserve as a moderator of outcomes in five clusters of first episode psychosis patients: a 10-year follow-up study of the PAFIP cohort. <i>Psychological Medicine</i> , 2023, 53, 1891-1905. | 4.5 | 3 |
| 65 | Disentangling early and late onset of psychosis in women: identifying new targets for treatment. <i>Archives of Women's Mental Health</i> , 2022, 25, 335-344. | 2.6 | 3 |
| 66 | Elevated risk of liver steatosis in first-episode psychosis patients: Results from a 3-year prospective study. <i>Schizophrenia Research</i> , 2022, 246, 30-38. | 2.0 | 3 |
| 67 | S101. DIFFERENTIATING NON-AFFECTIVE ACUTE REMITTING PSYCHOSIS (NARP) FROM SCHIZOPHRENIA IN FIRST-EPISODE PSYCHOSIS: FINDINGS FROM A 3-YEAR LONGITUDINAL STUDY IN THE PAFIP COHORT. <i>Schizophrenia Bulletin</i> , 2020, 46, S72-S73. | 4.3 | 2 |
| 68 | Long-term clinical and functional outcome after antipsychotic discontinuation in early phases of non-affective psychosis: Results from the PAFIP-10 cohort. <i>Schizophrenia Research</i> , 2021, 232, 28-30. | 2.0 | 2 |
| 69 | Long-term metabolic effect of second-generation antipsychotics in first episode of psychosis. <i>European Psychiatry</i> , 2017, 41, S388-S388. | 0.2 | 1 |
| 70 | Predictive value of prolactin in first episode psychosis at ten years follow-up. <i>Revista De Psiquiatría Y Salud Mental</i> , 2021, 14, 179-180. | 1.8 | 1 |
| 71 | Stability of schizophrenia diagnosis in a 10-year longitudinal study on first episode of non-affective psychosis: Conclusions from the PAFIP cohort. <i>Acta Psychiatrica Scandinavica</i> , 2021, 144, 342-357. | 4.5 | 1 |
| 72 | Different neurocognitive profiles of risperidone and aripiprazole in the FIRST episode of psychosis: A 3-year follow-up comparison. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2021, 110, 110309. | 4.8 | 1 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | Naturalistic study on the use of clozapine in the early phases of non-affective psychosis: A 10-year follow-up study in the PAFIP-10 cohort. <i>Journal of Psychiatric Research</i> , 2022, 153, 292-299. | 3.1 | 1 |
| 74 | A Cross-Sectional and Longitudinal Magnetic Resonance study of Insular Cortex in First-Episode Non-Affective Psychosis. Effect of Gender and Diagnostic specificity.. <i>NeuroImage</i> , 2009, 47, S174. | 4.2 | 0 |
| 75 | Effect of Disrupted-in-Schizophrenia 1 gene on treatment response in patients with a first episode of psychosis. <i>European Psychiatry</i> , 2016, 33, S183-S183. | 0.2 | 0 |
| 76 | Lack of differential long-term metabolic profile of aripiprazole, quetiapine and ziprasidone in first episode of psychosis. <i>European Psychiatry</i> , 2017, 41, S388-S388. | 0.2 | 0 |
| 77 | O6.7. COMMON NEUROANATOMICAL ABNORMALITIES IN FIRST EPISODE PSYCHOSIS ACROSS SEVERAL INDEPENDENT SAMPLES. <i>Schizophrenia Bulletin</i> , 2018, 44, S92-S92. | 4.3 | 0 |
| 78 | T175. A 10-YEAR LONGITUDINAL STUDY OF GREY MATTER VOLUME IN FIRST EPISODE OF NON-AFFECTIVE PSYCHOSIS. <i>Schizophrenia Bulletin</i> , 2018, 44, S184-S184. | 4.3 | 0 |
| 79 | F55INFLUENCE OF SCHIZOPHRENIA POLYGENIC RISK SCORES ON BRAIN SURFACE: AN EXPLORATORY STUDY IN FIRST EPISODE OF PSYCHOSIS. <i>European Neuropsychopharmacology</i> , 2019, 29, S1139. | 0.7 | 0 |
| 80 | SA114POLYGENIC RISK SCORE AND METABOLIC SYNDROME PREDICTION IN PATIENTS WITH A FIRST EPISODE OF NON-AFFECTIVE PSYCHOSIS. <i>European Neuropsychopharmacology</i> , 2019, 29, S1251. | 0.7 | 0 |
| 81 | M97. PREDICTORS OF TRANSITION TO SCHIZOPHRENIA IN BRIEF PSYCHOTIC DISORDER: FINDINGS FROM A 3-YEAR LONGITUDINAL STUDY IN THE PAFIP COHORT. <i>Schizophrenia Bulletin</i> , 2020, 46, S171-S172. | 4.3 | 0 |
| 82 | Data regarding active psychosis and functional outcome, among other clinical variables, during early phases of the illness in first-episode psychosis in the PAFIP 10-year follow-up program. <i>Data in Brief</i> , 2020, 30, 105599. | 1.0 | 0 |
| 83 | Predictive value of prolactin in first episode psychosis at ten years follow-up. <i>Revista De Psiquiatría Y Salud Mental (English Edition)</i> , 2021, 14, 179-180. | 0.3 | 0 |