

gabriella Fibbi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2696221/publications.pdf>

Version: 2024-02-01

83
papers

2,452
citations

159585

30
h-index

233421

45
g-index

84
all docs

84
docs citations

84
times ranked

2930
citing authors

#	ARTICLE	IF	CITATIONS
1	Matrix metalloproteinase 12-dependent cleavage of urokinase receptor in systemic sclerosis microvascular endothelial cells results in impaired angiogenesis. <i>Arthritis and Rheumatism</i> , 2004, 50, 3275-3285.	6.7	118
2	Interaction of urokinase with specific receptors stimulates mobilization of bovine adrenal capillary endothelial cells*1. <i>Experimental Cell Research</i> , 1988, 179, 385-395.	2.6	102
3	The Urokinase Receptor System, A Key Regulator at the Intersection between Inflammation, Immunity, and Coagulation. <i>Current Pharmaceutical Design</i> , 2011, 17, 1924-1943.	1.9	99
4	Multiple pathways of cell invasion are regulated by multiple families of serine proteases. <i>Clinical and Experimental Metastasis</i> , 2002, 19, 193-207.	3.3	94
5	The plasminogen activation system in inflammation. <i>Frontiers in Bioscience - Landmark</i> , 2008, Volume, 4667.	3.0	83
6	Domain 1 of the urokinase-type plasminogen activator receptor is required for its morphologic and functional, β 2 integrin-mediated connection with actin cytoskeleton in human microvascular endothelial cells: Failure of association in systemic sclerosis endothelial cells. <i>Arthritis and Rheumatism</i> , 2006, 54, 3926-3938.	6.7	77
7	Endothelial progenitor cell-dependent angiogenesis requires localization of the full-length form of uPAR in caveolae. <i>Blood</i> , 2011, 118, 3743-3755.	1.4	70
8	Effects of blocking urokinase receptor signaling by antisense oligonucleotides in a mouse model of experimental prostate cancer bone metastases. <i>Gene Therapy</i> , 2005, 12, 702-714.	4.5	67
9	Role of Specific Membrane Receptors in Urokinase-Dependent Migration of Human Keratinocytes. <i>Journal of Investigative Dermatology</i> , 1990, 94, 310-316.	0.7	63
10	bcl-2 Induction of Urokinase Plasminogen Activator Receptor Expression in Human Cancer Cells through Sp1 Activation. <i>Journal of Biological Chemistry</i> , 2004, 279, 6737-6745.	3.4	60
11	A model of anti-angiogenesis: differential transcriptosome profiling of microvascular endothelial cells from diffuse systemic sclerosis patients. <i>Arthritis Research and Therapy</i> , 2006, 8, R115.	3.5	56
12	The antiangiogenic tissue kallikrein pattern of endothelial cells in systemic sclerosis. <i>Arthritis and Rheumatism</i> , 2005, 52, 3618-3628.	6.7	55
13	Production of Second Messengers Following Chemotactic and Mitogenic Urokinase-Receptor Interaction in Human Fibroblasts and Mouse Fibroblasts Transfected with Human Urokinase Receptor. <i>Experimental Cell Research</i> , 1994, 213, 438-448.	2.6	53
14	Functions of the fibrinolytic system in human ito cells and its control by basic fibroblast and platelet-derived growth factor. <i>Hepatology</i> , 1999, 29, 868-878.	7.3	50
15	Cell Invasion Is Affected by Differential Expression of the Urokinase Plasminogen Activator/Urokinase Plasminogen Activator Receptor System in Muscle Satellite Cells from Normal and Dystrophic Patients. <i>Laboratory Investigation</i> , 2001, 81, 27-39.	3.7	48
16	TGF β 1 antagonistic peptides inhibit TGF β 1-dependent angiogenesis. <i>Biochemical Pharmacology</i> , 2009, 77, 813-825.	4.4	48
17	Extracellular acidity strengthens mesenchymal stem cells to promote melanoma progression. <i>Cell Cycle</i> , 2015, 14, 3088-3100.	2.6	47
18	Systemic sclerosis endothelial cells recruit and activate dermal fibroblasts by induction of a connective tissue growth factor (CCN2)/transforming growth factor β -dependent mesenchymal-to-mesenchymal transition. <i>Arthritis and Rheumatism</i> , 2013, 65, 258-269.	6.7	46

#	ARTICLE	IF	CITATIONS
19	One pot environmental friendly synthesis of gold nanoparticles using Punica Granatum Juice: A novel antioxidant agent for future dermatological and cosmetic applications. <i>Journal of Colloid and Interface Science</i> , 2018, 521, 50-61.	9.4	45
20	Antisense oligodeoxynucleotides for urokinase-plasminogen activator receptor have anti-invasive and anti-proliferative effects in vitro and inhibit spontaneous metastases of human melanoma in mice. <i>International Journal of Cancer</i> , 2004, 110, 125-133.	5.1	42
21	Modulation of the angiogenic phenotype of normal and systemic sclerosis endothelial cells by gain and loss of function of pentraxin 3 and matrix metalloproteinase 12. <i>Arthritis and Rheumatism</i> , 2010, 62, 2488-2498.	6.7	42
22	The receptor for urokinase-plasminogen activator (uPAR) controls plasticity of cancer cell movement in mesenchymal and amoeboid migration style. <i>Oncotarget</i> , 2014, 5, 1538-1553.	1.8	42
23	uPA/uPAR system activation drives a glycolytic phenotype in melanoma cells. <i>International Journal of Cancer</i> , 2017, 141, 1190-1200.	5.1	40
24	Inhibition of uPAR-TGF β 2 crosstalk blocks MSC-dependent EMT in melanoma cells. <i>Journal of Molecular Medicine</i> , 2015, 93, 783-794.	3.9	39
25	uPAR-expressing melanoma exosomes promote angiogenesis by VE-Cadherin, EGFR and uPAR overexpression and rise of ERK1,2 signaling in endothelial cells. <i>Cellular and Molecular Life Sciences</i> , 2021, 78, 3057-3072.	5.4	38
26	EphA2-mediated mesenchymal to amoeboid transition induced by endothelial progenitor cells enhances metastatic spread due to cancer-associated fibroblasts. <i>Journal of Molecular Medicine</i> , 2013, 91, 103-115.	3.9	37
27	Melanoma cell therapy: Endothelial progenitor cells as shuttle of the MMP12 uPAR-degrading enzyme. <i>Oncotarget</i> , 2014, 5, 3711-3727.	1.8	37
28	Everolimus selectively targets vemurafenib resistant BRAFV600E melanoma cells adapted to low pH. <i>Cancer Letters</i> , 2017, 408, 43-54.	7.2	36
29	The Mr 17 500 region of the A chain of urokinase is required for interaction with a specific receptor in A431 cells. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 1986, 885, 301-308.	4.1	33
30	Systemic Sclerosis-Endothelial Cell Antiangiogenic Pentraxin 3 and Matrix Metalloprotease 12 Control Human Breast Cancer Tumor Vascularization and Development in Mice. <i>Neoplasia</i> , 2009, 11, 1106-1115.	5.3	32
31	GDF5 Regulates TGF β -Dependent Angiogenesis in Breast Carcinoma MCF-7 Cells: In Vitro and In Vivo Control by Anti-TGF β Peptides. <i>PLoS ONE</i> , 2012, 7, e50342.	2.5	31
32	EGFR/uPAR interaction as druggable target to overcome vemurafenib acquired resistance in melanoma cells. <i>EBioMedicine</i> , 2019, 39, 194-206.	6.1	31
33	Reduction of in vitro invasion and in vivo cartilage degradation in a SCID mouse model by loss of function of the fibrinolytic system of rheumatoid arthritis synovial fibroblasts. <i>Arthritis and Rheumatism</i> , 2011, 63, 2584-2594.	6.7	30
34	Chronic Resveratrol Treatment Inhibits MRC5 Fibroblast SASP-Related Protumoral Effects on Melanoma Cells. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2017, 72, 1187-1195.	3.6	29
35	Interaction of urokinase a chain with the cellular receptor induces both urokinase autocriny and cell movement. <i>Fibrinolysis</i> , 1989, 3, 1.	0.5	27
36	Desmoglein-2-Integrin Beta-8 Interaction Regulates Actin Assembly in Endothelial Cells: Deregulation in Systemic Sclerosis. <i>PLoS ONE</i> , 2013, 8, e68117.	2.5	27

#	ARTICLE	IF	CITATIONS
37	Lipid rafts: integrated platforms for vascular organization offering therapeutic opportunities. Cellular and Molecular Life Sciences, 2015, 72, 1537-1557.	5.4	25
38	Adhesion-dependent heparin production by platelets. Nature, 1982, 296, 352-353.	27.8	23
39	Transforming Growth Factor Beta-1 Stimulates Invasivity of Hepatic Stellate Cells by Engagement of the Cell-associated Fibrinolytic System. Growth Factors, 2001, 19, 87-100.	1.7	23
40	Growth Factor-Dependent Proliferation and Invasion of Muscle Satellite Cells Require the Cell-Associated Fibrinolytic System. Biological Chemistry, 2002, 383, 127-36.	2.5	22
41	Mature and progenitor endothelial cells perform angiogenesis also under protease inhibition: the amoeboid angiogenesis. Journal of Experimental and Clinical Cancer Research, 2018, 37, 74.	8.6	21
42	Selective localization of receptors for urokinase amino-terminal fragment at substratum contact sites of an in vitro-established line of human epidermal cells. Experimental Cell Research, 1992, 203, 427-434.	2.6	20
43	Tumor-tropic endothelial colony forming cells (ECFCs) loaded with near-infrared sensitive Au nanoparticles: A cellular stove approach to the photoablation of melanoma. Oncotarget, 2016, 7, 39846-39860.	1.8	20
44	Cell surface glycosaminoglycans in normal and leukemic leukocytes. Cell Differentiation, 1980, 9, 71-81.	0.4	19
45	Interaction of urokinase a chain with the receptor of human keratinocytes stimulates release of urokinase-like plasminogen activator. Experimental Cell Research, 1990, 187, 33-38.	2.6	19
46	Differential u PAR recruitment in caveolar lipid rafts by GM 1 and GM 3 gangliosides regulates endothelial progenitor cells angiogenesis. Journal of Cellular and Molecular Medicine, 2015, 19, 113-123.	3.6	19
47	Interaction of Urokinase-Type Plasminogen Activator with Its Receptor Rapidly Induces Activation of Glucose Transporters. Biochemistry, 1997, 36, 3076-3083.	2.5	18
48	Endothelial sphingosine kinase/SPNS2 axis is critical for vessel-like formation by human mesoangioblasts. Journal of Molecular Medicine, 2015, 93, 1145-1157.	3.9	18
49	Endothelial Progenitor Cells as Shuttle of Anticancer Agents. Human Gene Therapy, 2016, 27, 784-791.	2.7	18
50	Glycosaminoglycan changes involved in polymorphonuclear leukocyte activation in vitro. Journal of Cellular Physiology, 1982, 111, 149-154.	4.1	16
51	Glycolysis-derived acidic microenvironment as a driver of endothelial dysfunction in systemic sclerosis. Rheumatology, 2021, 60, 4508-4519.	1.9	16
52	Involvement of chondroitin sulphate in preventing adhesive cellular interactions. Biochimica Et Biophysica Acta - Molecular Cell Research, 1983, 762, 512-518.	4.1	15
53	uPAR Knockout Results in a Deep Glycolytic and OXPHOS Reprogramming in Melanoma and Colon Carcinoma Cell Lines. Cells, 2020, 9, 308.	4.1	15
54	Interaction of urokinase with specific receptors abolishes the time of commitment to terminal differentiation of murine erythroleukaemia (Friend) cells. British Journal of Haematology, 1987, 66, 289-294.	2.5	15

#	ARTICLE	IF	CITATIONS
55	Modulation of surface-associated urokinase: Binding, interiorization, delivery to lysosomes, and degradation in human keratinocytes. <i>Experimental Cell Research</i> , 1991, 193, 346-355.	2.6	14
56	Chronic Resveratrol Treatment Reduces the Pro-angiogenic Effect of Human Fibroblast Senescent-Associated Secretory Phenotype on Endothelial Colony-Forming Cells: The Role of IL8. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2019, 74, 625-633.	3.6	14
57	Oleuropein aglycone attenuates the pro-angiogenic phenotype of senescent fibroblasts: A functional study in endothelial cells. <i>Journal of Functional Foods</i> , 2019, 53, 219-226.	3.4	14
58	Piascledine modulates the production of VEGF and TIMP-1 and reduces the invasiveness of rheumatoid arthritis synoviocytes. <i>Scandinavian Journal of Rheumatology</i> , 2006, 35, 346-350.	1.1	12
59	Plasminogen activators and inhibitor type-1 in alveolar osteitis. <i>European Journal of Oral Sciences</i> , 2006, 114, 500-503.	1.5	12
60	Parvovirus B19 activates in vitro normal human dermal fibroblasts: a possible implication in skin fibrosis and systemic sclerosis. <i>Rheumatology</i> , 2020, 59, 3526-3532.	1.9	12
61	Enhanced Antitumoral Activity and Photoacoustic Imaging Properties of AuNP-Enriched Endothelial Colony Forming Cells on Melanoma. <i>Advanced Science</i> , 2021, 8, 2001175.	11.2	12
62	Effects of hyaluronate and heparan sulphate on collagen-fibronectin interactions. <i>International Journal of Biological Macromolecules</i> , 1982, 4, 67-72.	7.5	11
63	Regulation of Urokinase/Urokinase Receptor Interaction by Heparin-like Glycosaminoglycans. <i>Journal of Biological Chemistry</i> , 2001, 276, 4756-4765.	3.4	11
64	Prep1 regulates angiogenesis through a PGC-1 β -mediated mechanism. <i>FASEB Journal</i> , 2019, 33, 13893-13904.	0.5	11
65	CRISPR/Cas9 uPAR Gene Knockout Results in Tumor Growth Inhibition, EGFR Downregulation and Induction of Stemness Markers in Melanoma and Colon Carcinoma Cell Lines. <i>Frontiers in Oncology</i> , 2021, 11, 663225.	2.8	11
66	uPAR Controls Vasculogenic Mimicry Ability Expressed by Drug-Resistant Melanoma Cells. <i>Oncology Research</i> , 2021, 28, 873-884.	1.5	10
67	Proteomic Identification of VEGF-dependent Protein Enrichment to Membrane Caveolar-raft Microdomains in Endothelial Progenitor Cells. <i>Molecular and Cellular Proteomics</i> , 2013, 12, 1926-1938.	3.8	9
68	Synthesis and characterization of modified magnetic nanoparticles as theranostic agents: in vitro safety assessment in healthy cells. <i>Toxicology in Vitro</i> , 2021, 72, 105094.	2.4	9
69	Plasminogen activator: Morphological evidence of binding, internalization and delivery to lysosomes in 3T3 mouse fibroblasts. <i>The Histochemical Journal</i> , 1985, 17, 333-341.	0.6	8
70	Proteases and extracellular environment. <i>Thrombosis and Haemostasis</i> , 2005, 93, 190-191.	3.4	8
71	Th17 lymphocyte-dependent degradation of joint cartilage by synovial fibroblasts in a humanized mouse model of arthritis and reversal by secukinumab. <i>European Journal of Immunology</i> , 2021, 51, 220-230.	2.9	8
72	Interaction of urokinase with specific receptors abolishes the time of commitment to terminal differentiation of murine erythroleukaemia (Friend) cells. <i>British Journal of Haematology</i> , 1987, 66, 289-294.	2.5	7

#	ARTICLE	IF	CITATIONS
73	Cell-Mediated Release of Nanoparticles as a Preferential Option for Future Treatment of Melanoma. <i>Cancers</i> , 2020, 12, 1771.	3.7	6
74	Non-Enzymatic Activities of Proteases: From Scepticism to Reality. <i>Biological Chemistry</i> , 2002, 383, 1-4.	2.5	5
75	Parvovirus B19 induces cellular senescence in human dermal fibroblasts: putative role in systemic sclerosis-associated fibrosis. <i>Rheumatology</i> , 2021, , .	1.9	5
76	Urokinase and its receptor in follicular and inflammatory cysts of the jaws. <i>Oral Diseases</i> , 2010, 16, 753-759.	3.0	4
77	Altered clot formation and lysis are associated with increased fibrinolytic activity in ascites in patients with advanced cirrhosis. <i>Internal and Emergency Medicine</i> , 2021, 16, 339-347.	2.0	4
78	A Possible Role for PAI-1 Blockade in Melanoma Immunotherapy. <i>Journal of Investigative Dermatology</i> , 2021, 141, 2566-2568.	0.7	3
79	Electrophoretic Characterization of Surface Heparan Sulphates in Normal and Virus Transformed 3T3 Cells. <i>Caryologia</i> , 1980, 33, 441-448.	0.3	2
80	Modulation of Surface-Associated Urokinase in Different Cell Lines: Evidence for Urokinase Interiorization and Degradation. <i>Seminars in Thrombosis and Hemostasis</i> , 1991, 17, 262-267.	2.7	2
81	STUDIES ON GLYCOSAMINOGLYCAN-DEPENDENT PROTEASE INHIBITORS. , 1982, , 353-359.		2
82	Role of urokinase receptors of human keratinocytes and dermal fibroblasts. <i>Fibrinolysis</i> , 1989, 3, 1-2.	0.5	1
83	Cell-Type-Independent Accumulation of Phosphatidic Acid Induced by Trifluoperazine in Stimulated Human Platelets, Leukocytes, and Fibroblasts. , 1984, , 75-79.		0