

# Andrea Nistri

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/2700159/andrea-nistri-publications-by-year.pdf>

**Version:** 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

144  
papers

4,659  
citations

35  
h-index

61  
g-index

146  
ext. papers

5,028  
ext. citations

4.4  
avg, IF

5.57  
L-index

#	Paper	IF	Citations
144	GABAergic Mechanisms Can Redress the Tilted Balance between Excitation and Inhibition in Damaged Spinal Networks. <i>Molecular Neurobiology</i> , <b>2021</b> , 58, 3769-3786	6.2	2
143	Overexpressed Na 1.7 Channels Confer Hyperexcitability to Trigeminal Sensory Neurons of Ca 2.1 Mutant Hemiplegic Migraine Mice. <i>Frontiers in Cellular Neuroscience</i> , <b>2021</b> , 15, 640709	6.1	0
142	Modeling a Nociceptive Neuro-Immune Synapse Activated by ATP and 5-HT in Meninges: Novel Clues on Transduction of Chemical Signals Into Persistent or Rhythmic Neuronal Firing. <i>Frontiers in Cellular Neuroscience</i> , <b>2020</b> , 14, 135	6.1	9
141	Modulation of extrasynaptic GABAergic receptor activity influences glutamate release and neuronal survival following excitotoxic damage to mouse spinal cord neurons. <i>Neurochemistry International</i> , <b>2019</b> , 128, 175-185	4.4	8
140	Investigation on 2S3S-Substituted ATP Derivatives and Analogs as Novel P2X3 Receptor Antagonists. <i>ACS Medicinal Chemistry Letters</i> , <b>2019</b> , 10, 493-498	4.3	6
139	Loss of inhibitory synapses causes locomotor network dysfunction of the rat spinal cord during prolonged maintenance in vitro. <i>Brain Research</i> , <b>2019</b> , 1710, 8-21	3.7	1
138	Long-term application of cannabinoids leads to dissociation between changes in cAMP and modulation of GABA receptors of mouse trigeminal sensory neurons. <i>Neurochemistry International</i> , <b>2019</b> , 126, 74-85	4.4	3
137	Pharmacological induction of Heat Shock Protein 70 by celastrol protects motoneurons from excitotoxicity in rat spinal cord in vitro. <i>European Journal of Neuroscience</i> , <b>2019</b> , 49, 215-231	3.5	7
136	Mechanism of Neuroprotection Against Experimental Spinal Cord Injury by Riluzole or Methylprednisolone. <i>Neurochemical Research</i> , <b>2019</b> , 44, 200-213	4.6	28
135	Functional up-regulation of the M-current by retigabine contrasts hyperexcitability and excitotoxicity on rat hypoglossal motoneurons. <i>Journal of Physiology</i> , <b>2018</b> , 596, 2611-2629	3.9	5
134	Differential neuromodulatory role of endocannabinoids in the rodent trigeminal sensory ganglion and cerebral cortex relevant to pain processing. <i>Neuropharmacology</i> , <b>2018</b> , 131, 39-50	5.5	4
133	Nicotine-mediated neuroprotection of rat spinal networks against excitotoxicity. <i>European Journal of Neuroscience</i> , <b>2018</b> , 47, 1353-1374	3.5	7
132	Electrophysiological characterization of the M-current in rat hypoglossal motoneurons. <i>Neuroscience</i> , <b>2017</b> , 340, 62-75	3.9	9
131	Ceftriaxone-mediated upregulation of the glutamate transporter GLT-1 contrasts neurotoxicity evoked by kainate in rat organotypic spinal cord cultures. <i>NeuroToxicology</i> , <b>2017</b> , 60, 34-41	4.4	7
130	Nicotine protects rat hypoglossal motoneurons from excitotoxic death via downregulation of connexin 36. <i>Cell Death and Disease</i> , <b>2017</b> , 8, e2881	9.8	6
129	Hyperpolarization-activated current I in mouse trigeminal sensory neurons in a transgenic mouse model of familial hemiplegic migraine type-1. <i>Neuroscience</i> , <b>2017</b> , 351, 47-64	3.9	5
128	ASIC channel inhibition enhances excitotoxic neuronal death in an in vitro model of spinal cord injury. <i>Neuroscience</i> , <b>2017</b> , 343, 398-410	3.9	13

127	Nicotinic receptors modulate the onset of reactive oxygen species production and mitochondrial dysfunction evoked by glutamate uptake block in the rat hypoglossal nucleus. <i>Neuroscience Letters</i> , <b>2017</b> , 639, 43-48	3.3	12
126	Propofol Protects Rat Hypoglossal Motoneurons in an In Vitro Model of Excitotoxicity by Boosting GABAergic Inhibition and Reducing Oxidative Stress. <i>Neuroscience</i> , <b>2017</b> , 367, 15-33	3.9	6
125	In situ imaging reveals properties of purinergic signalling in trigeminal sensory ganglia in vitro. <i>Purinergic Signalling</i> , <b>2017</b> , 13, 511-520	3.8	13
124	Neurotoxicity of propofol on rat hypoglossal motoneurons in vitro. <i>Neuroscience Letters</i> , <b>2017</b> , 655, 95-100	3.9	5
123	2S3SO-Substituted ATP derivatives as potent antagonists of purinergic P2X3 receptors and potential analgesic agents. <i>Purinergic Signalling</i> , <b>2017</b> , 13, 61-74	3.8	6
122	Neuroprotective effect of propofol against excitotoxic injury to locomotor networks of the rat spinal cord in vitro. <i>European Journal of Neuroscience</i> , <b>2016</b> , 44, 2418-2430	3.5	14
121	A study of cannabinoid-1 receptors during the early phase of excitotoxic damage to rat spinal locomotor networks in vitro. <i>Neuroscience</i> , <b>2016</b> , 333, 214-28	3.9	2
120	Expression and function of calcitonin gene-related peptide (CGRP) receptors in trigeminal ganglia of R192Q Cacna1a knock-in mice. <i>Neuroscience Letters</i> , <b>2016</b> , 620, 104-10	3.3	7
119	Delayed application of the anesthetic propofol contrasts the neurotoxic effects of kainate on rat organotypic spinal slice cultures. <i>NeuroToxicology</i> , <b>2016</b> , 54, 1-10	4.4	6
118	A study of methylprednisolone neuroprotection against acute injury to the rat spinal cord in vitro. <i>Neuroscience</i> , <b>2016</b> , 315, 136-49	3.9	13
117	Inefficient constitutive inhibition of P2X3 receptors by brain natriuretic peptide system contributes to sensitization of trigeminal sensory neurons in a genetic mouse model of familial hemiplegic migraine. <i>Molecular Pain</i> , <b>2016</b> , 12,	3.4	17
116	Loss of inhibition by brain natriuretic peptide over P2X3 receptors contributes to enhanced spike firing of trigeminal ganglion neurons in a mouse model of familial hemiplegic migraine type-1. <i>Neuroscience</i> , <b>2016</b> , 331, 197-205	3.9	13
115	Nicotinic receptor activation contrasts pathophysiological bursting and neurodegeneration evoked by glutamate uptake block on rat hypoglossal motoneurons. <i>Journal of Physiology</i> , <b>2016</b> , 594, 6777-6798	3.9	10
114	Modulatory effects by CB1 receptors on rat spinal locomotor networks after sustained application of agonists or antagonists. <i>Neuroscience</i> , <b>2015</b> , 303, 16-33	3.9	7
113	The volatile anesthetic methoxyflurane protects motoneurons against excitotoxicity in an in vitro model of rat spinal cord injury. <i>Neuroscience</i> , <b>2015</b> , 285, 269-80	3.9	11
112	Brain natriuretic peptide constitutively downregulates P2X3 receptors by controlling their phosphorylation state and membrane localization. <i>Molecular Pain</i> , <b>2015</b> , 11, 71	3.4	13
111	Role of HSP70 in motoneuron survival after excitotoxic stress in a rat spinal cord injury model in vitro. <i>European Journal of Neuroscience</i> , <b>2015</b> , 42, 3054-65	3.5	13
110	Role of calcitonin gene-related peptide and brain natriuretic peptide to modulate the excitability state of trigeminal neurons: relevance to migraine pathology and treatment. <i>Journal of Receptor, Ligand and Channel Research</i> , <b>2015</b> , 31		1

109	Dynamic expression of ATF3 as a novel tool to study activation and migration of endogenous spinal stem cells and their role in neural repair. <i>Neural Regeneration Research</i> , <b>2015</b> , 10, 713-4	4.5	1
108	S100 $\beta$ as an early biomarker of excitotoxic damage in spinal cord organotypic cultures. <i>Journal of Neurochemistry</i> , <b>2014</b> , 130, 598-604	6	22
107	A hyperexcitability phenotype in mouse trigeminal sensory neurons expressing the R192Q Cacna1a missense mutation of familial hemiplegic migraine type-1. <i>Neuroscience</i> , <b>2014</b> , 266, 244-54	3.9	17
106	ATF3 is a novel nuclear marker for migrating ependymal stem cells in the rat spinal cord. <i>Stem Cell Research</i> , <b>2014</b> , 12, 815-27	1.6	23
105	ATP-gated P2X receptors in health and disease. <i>Frontiers in Cellular Neuroscience</i> , <b>2014</b> , 8, 204	6.1	20
104	Calcium/calmodulin-dependent serine protein kinase (CASK) is a new intracellular modulator of P2X3 receptors. <i>Journal of Neurochemistry</i> , <b>2013</b> , 126, 102-12	6	16
103	Mutated CaV2.1 channels dysregulate CASK/P2X3 signaling in mouse trigeminal sensory neurons of R192Q Cacna1a knock-in mice. <i>Molecular Pain</i> , <b>2013</b> , 9, 62	3.4	4
102	Effects of LPS on P2X3 receptors of trigeminal sensory neurons and macrophages from mice expressing the R192Q Cacna1a gene mutation of familial hemiplegic migraine-1. <i>Purinergic Signalling</i> , <b>2013</b> , 9, 7-13	3.8	16
101	Acute Spinal Cord Injury In Vitro: Insight into Basic Mechanisms. <i>Neuromethods</i> , <b>2013</b> , 39-62	0.4	3
100	Evaluation of adenine as scaffold for the development of novel P2X3 receptor antagonists. <i>European Journal of Medicinal Chemistry</i> , <b>2013</b> , 65, 41-50	6.8	4
99	Riluzole: what it does to spinal and brainstem neurons and how it does it. <i>Neuroscientist</i> , <b>2013</b> , 19, 137-44	4.6	35
98	B-type natriuretic peptide-induced delayed modulation of TRPV1 and P2X3 receptors of mouse trigeminal sensory neurons. <i>PLoS ONE</i> , <b>2013</b> , 8, e81138	3.7	22
97	Desensitization properties of P2X3 receptors shaping pain signaling. <i>Frontiers in Cellular Neuroscience</i> , <b>2013</b> , 7, 245	6.1	28
96	Microelectrode arrays in combination with in vitro models of spinal cord injury as tools to investigate pathological changes in network activity: facts and promises. <i>Frontiers in Neuroengineering</i> , <b>2013</b> , 6, 2		4
95	TNF $\alpha$ levels and macrophages expression reflect an inflammatory potential of trigeminal ganglia in a mouse model of familial hemiplegic migraine. <i>PLoS ONE</i> , <b>2013</b> , 8, e52394	3.7	57
94	The mechanism of functional up-regulation of P2X3 receptors of trigeminal sensory neurons in a genetic mouse model of familial hemiplegic migraine type 1 (FHM-1). <i>PLoS ONE</i> , <b>2013</b> , 8, e60677	3.7	30
93	Unusual increase in lumbar network excitability of the rat spinal cord evoked by the PARP-1 inhibitor PJ-34 through inhibition of glutamate uptake. <i>Neuropharmacology</i> , <b>2012</b> , 63, 415-26	5.5	13
92	Functional differences between ATP-gated human and rat P2X3 receptors are caused by critical residues of the intracellular C-terminal domain. <i>Journal of Neurochemistry</i> , <b>2012</b> , 122, 557-67	6	13

91	Functional crosstalk in culture between macrophages and trigeminal sensory neurons of a mouse genetic model of migraine. <i>BMC Neuroscience</i> , <b>2012</b> , 13, 143	3.2	25
90	Postnatal developmental profile of neurons and glia in motor nuclei of the brainstem and spinal cord, and its comparison with organotypic slice cultures. <i>Developmental Neurobiology</i> , <b>2012</b> , 72, 1140-60 <sup>3,2</sup>		35
89	Regulation of P2X3 receptor structure and function. <i>CNS and Neurological Disorders - Drug Targets</i> , <b>2012</b> , 11, 687-98	2.6	18
88	Calcitonin gene-related peptide-mediated enhancement of purinergic neuron/glia communication by the algogenic factor bradykinin in mouse trigeminal ganglia from wild-type and R192Q Cav2.1 Knock-in mice: implications for basic mechanisms of migraine pain. <i>Journal of Neuroscience</i> , <b>2011</b> , 31, 3638-49	6.6	101
87	Molecular Mechanisms Underlying Cell Death in Spinal Networks in Relation to Locomotor Activity After Acute Injury in vitro. <i>Frontiers in Cellular Neuroscience</i> , <b>2011</b> , 5, 9	6.1	30
86	Riluzole is a potent drug to protect neonatal rat hypoglossal motoneurons in vitro from excitotoxicity due to glutamate uptake block. <i>European Journal of Neuroscience</i> , <b>2011</b> , 33, 899-913	3.5	22
85	Studies of locomotor network neuroprotection by the selective poly(ADP-ribose) polymerase-1 inhibitor PJ-34 against excitotoxic injury to the rat spinal cord in vitro. <i>European Journal of Neuroscience</i> , <b>2011</b> , 33, 2216-27	3.5	16
84	Respiratory motoneurons and pathological conditions: lessons from hypoglossal motoneurons challenged by excitotoxic or oxidative stress. <i>Respiratory Physiology and Neurobiology</i> , <b>2011</b> , 179, 89-96	2.8	14
83	Effect of the PARP-1 inhibitor PJ 34 on excitotoxic damage evoked by kainate on rat spinal cord organotypic slices. <i>Cellular and Molecular Neurobiology</i> , <b>2011</b> , 31, 469-78	4.6	18
82	Effects of 6(5H)-phenanthridinone, an inhibitor of poly(ADP-ribose)polymerase-1 activity (PARP-1), on locomotor networks of the rat isolated spinal cord. <i>Cellular and Molecular Neurobiology</i> , <b>2011</b> , 31, 503-8	4.6	21
81	Lipid rafts control P2X3 receptor distribution and function in trigeminal sensory neurons of a transgenic migraine mouse model. <i>Molecular Pain</i> , <b>2011</b> , 7, 77	3.4	32
80	Cystic fibrosis transmembrane conductance regulator modulates synaptic chloride homeostasis in motoneurons of the rat spinal cord during neonatal development. <i>Developmental Neurobiology</i> , <b>2011</b> , 71, 253-68	3.2	20
79	Electrochemical detection of endogenous glutamate release from rat spinal cord organotypic slices as a real-time method to monitor excitotoxicity. <i>Journal of Neuroscience Methods</i> , <b>2011</b> , 197, 128-32	3	19
78	Dynamics of early locomotor network dysfunction following a focal lesion in an in vitro model of spinal injury. <i>European Journal of Neuroscience</i> , <b>2010</b> , 31, 60-78	3.5	21
77	Transient oxidative stress evokes early changes in the functional properties of neonatal rat hypoglossal motoneurons in vitro. <i>European Journal of Neuroscience</i> , <b>2010</b> , 31, 951-66	3.5	25
76	Deconstructing locomotor networks with experimental injury to define their membership. <i>Annals of the New York Academy of Sciences</i> , <b>2010</b> , 1198, 242-51	6.5	7
75	The Cdk5 kinase downregulates ATP-gated ionotropic P2X3 receptor function via serine phosphorylation. <i>Cellular and Molecular Neurobiology</i> , <b>2010</b> , 30, 505-9	4.6	23
74	Kainate-mediated excitotoxicity induces neuronal death in the rat spinal cord in vitro via a PARP-1 dependent cell death pathway (Parthanatos). <i>Cellular and Molecular Neurobiology</i> , <b>2010</b> , 30, 1001-12	4.6	49

73	Familial hemiplegic migraine Ca(v)2.1 channel mutation R192Q enhances ATP-gated P2X3 receptor activity of mouse sensory ganglion neurons mediating trigeminal pain. <i>Molecular Pain</i> , <b>2010</b> , 6, 48	3.4	54
72	The C-terminal Src inhibitory kinase (Csk)-mediated tyrosine phosphorylation is a novel molecular mechanism to limit P2X3 receptor function in mouse sensory neurons. <i>Journal of Biological Chemistry</i> , <b>2009</b> , 284, 21393-401	5.4	33
71	A repertoire of rhythmic bursting produced by hypoglossal motoneurons in physiological and pathological conditions. <i>Philosophical Transactions of the Royal Society B: Biological Sciences</i> , <b>2009</b> , 364, 2493-500	5.8	19
70	The patterns of spontaneous Ca <sup>2+</sup> signals generated by ventral spinal neurons in vitro show time-dependent refinement. <i>European Journal of Neuroscience</i> , <b>2009</b> , 29, 1543-59	3.5	10
69	Adenine-based acyclic nucleotides as novel P2X3 receptor ligands. <i>Journal of Medicinal Chemistry</i> , <b>2009</b> , 52, 4596-603	8.3	21
68	Spinal cord injury: there is nothing permanent except change (Heraclitus, 540-480 BC). <i>Brain Research Bulletin</i> , <b>2009</b> , 78, 2-3	3.9	0
67	Riluzole blocks persistent Na <sup>+</sup> and Ca <sup>2+</sup> currents and modulates release of glutamate via presynaptic NMDA receptors on neonatal rat hypoglossal motoneurons in vitro. <i>European Journal of Neuroscience</i> , <b>2008</b> , 27, 2501-14	3.5	115
66	N-methyl-D-aspartate triggers neonatal rat hypoglossal motoneurons in vitro to express rhythmic bursting with unusual Mg <sup>2+</sup> sensitivity. <i>Neuroscience</i> , <b>2008</b> , 154, 804-20	3.9	13
65	Kainate and metabolic perturbation mimicking spinal injury differentially contribute to early damage of locomotor networks in the in vitro neonatal rat spinal cord. <i>Neuroscience</i> , <b>2008</b> , 155, 538-55	3.9	52
64	Molecular mechanisms of sensitization of pain-transducing P2X3 receptors by the migraine mediators CGRP and NGF. <i>Molecular Neurobiology</i> , <b>2008</b> , 37, 83-90	6.2	108
63	The effects induced by the sulphonylurea glibenclamide on the neonatal rat spinal cord indicate a novel mechanism to control neuronal excitability and inhibitory neurotransmission. <i>British Journal of Pharmacology</i> , <b>2007</b> , 150, 47-57	8.6	15
62	Activity-independent intracellular Ca <sup>2+</sup> oscillations are spontaneously generated by ventral spinal neurons during development in vitro. <i>Cell Calcium</i> , <b>2007</b> , 41, 317-29	4	23
61	Exocytotic release of ATP from cultured astrocytes. <i>Journal of Biological Chemistry</i> , <b>2007</b> , 282, 28749-28758	5.8	200
60	Neutralization of nerve growth factor induces plasticity of ATP-sensitive P2X3 receptors of nociceptive trigeminal ganglion neurons. <i>Journal of Neuroscience</i> , <b>2007</b> , 27, 8190-201	6.6	72
59	ERG conductance expression modulates the excitability of ventral horn GABAergic interneurons that control rhythmic oscillations in the developing mouse spinal cord. <i>Journal of Neuroscience</i> , <b>2007</b> , 27, 919-28	6.6	46
58	Delayed upregulation of ATP P2X3 receptors of trigeminal sensory neurons by calcitonin gene-related peptide. <i>Journal of Neuroscience</i> , <b>2006</b> , 26, 6163-71	6.6	138
57	Comparison of P2X and TRPV1 receptors in ganglia or primary culture of trigeminal neurons and their modulation by NGF or serotonin. <i>Molecular Pain</i> , <b>2006</b> , 2, 11	3.4	87
56	Experimental and modeling studies of desensitization of P2X3 receptors. <i>Molecular Pharmacology</i> , <b>2006</b> , 70, 373-82	4.3	57

55	Persistent rhythmic oscillations induced by nicotine on neonatal rat hypoglossal motoneurons in vitro. <i>European Journal of Neuroscience</i> , <b>2006</b> , 24, 2543-56	3.5	10
54	Glutamate uptake block triggers deadly rhythmic bursting of neonatal rat hypoglossal motoneurons. <i>Journal of Physiology</i> , <b>2006</b> , 572, 407-23	3.9	27
53	Tuning and playing a motor rhythm: how metabotropic glutamate receptors orchestrate generation of motor patterns in the mammalian central nervous system. <i>Journal of Physiology</i> , <b>2006</b> , 572, 323-34	3.9	48
52	Oscillatory circuits underlying locomotor networks in the rat spinal cord. <i>Critical Reviews in Neurobiology</i> , <b>2006</b> , 18, 25-36		18
51	Desensitization of nicotinic ACh receptors: shaping cholinergic signaling. <i>Trends in Neurosciences</i> , <b>2005</b> , 28, 371-8	13.3	280
50	An electron microscopy study of changes in dense core vesicles of PC12 cells following pulse depolarization. <i>NeuroReport</i> , <b>2005</b> , 16, 381-5	1.7	2
49	Activation and desensitization of neuronal nicotinic receptors modulate glutamatergic transmission on neonatal rat hypoglossal motoneurons. <i>European Journal of Neuroscience</i> , <b>2005</b> , 22, 2723-34	3.5	45
48	Desensitization of neuronal nicotinic receptors of human neuroblastoma SH-SY5Y cells during short or long exposure to nicotine. <i>British Journal of Pharmacology</i> , <b>2005</b> , 146, 1087-95	8.6	23
47	Long-term exposure to the new nicotinic antagonist 1,2-bisN-cytisinylethane upregulates nicotinic receptor subtypes of SH-SY5Y human neuroblastoma cells. <i>British Journal of Pharmacology</i> , <b>2005</b> , 146, 1096-109	8.6	22
46	Metabotropic glutamate receptor activity induces a novel oscillatory pattern in neonatal rat hypoglossal motoneurons. <i>Journal of Physiology</i> , <b>2005</b> , 563, 139-59	3.9	29
45	Activation of group I metabotropic glutamate receptors depresses recurrent inhibition of motoneurons in the neonatal rat spinal cord in vitro. <i>Experimental Brain Research</i> , <b>2005</b> , 164, 406-10	2.3	7
44	Identification of negative residues in the P2X3 ATP receptor ectodomain as structural determinants for desensitization and the Ca <sup>2+</sup> -sensing modulatory sites. <i>Journal of Biological Chemistry</i> , <b>2004</b> , 279, 53109-15	5.4	45
43	Activation of group I metabotropic glutamate receptors enhances efficacy of glutamatergic inputs to neonatal rat hypoglossal motoneurons in vitro. <i>European Journal of Neuroscience</i> , <b>2004</b> , 20, 1245-54	3.5	15
42	Modulation of rhythmic patterns and cumulative depolarization by group I metabotropic glutamate receptors in the neonatal rat spinal cord in vitro. <i>European Journal of Neuroscience</i> , <b>2004</b> , 19, 533-41	3.5	29
41	Agonist-dependence of recovery from desensitization of P2X(3) receptors provides a novel and sensitive approach for their rapid up or downregulation. <i>British Journal of Pharmacology</i> , <b>2004</b> , 141, 1048-58	8.6	46
40	Chronic NGF treatment of rat nociceptive DRG neurons in culture facilitates desensitization and deactivation of GABAA receptor-mediated currents. <i>British Journal of Pharmacology</i> , <b>2004</b> , 142, 425-34	8.6	9
39	Quantal release of ATP from clusters of PC12 cells. <i>Journal of Physiology</i> , <b>2004</b> , 560, 505-17	3.9	32
38	Role of group II and III metabotropic glutamate receptors in rhythmic patterns of the neonatal rat spinal cord in vitro. <i>Experimental Brain Research</i> , <b>2004</b> , 156, 495-504	2.3	14

37	Effect of metabotropic glutamate receptor activity on rhythmic discharges of the neonatal rat spinal cord in vitro. <i>Experimental Brain Research</i> , <b>2003</b> , 153, 388-93	2.3	17
36	Molecular biology and electrophysiology of neuronal nicotinic receptors of rat chromaffin cells. <i>European Journal of Neuroscience</i> , <b>2003</b> , 17, 2313-22	3.5	55
35	Distinct subtypes of group I metabotropic glutamate receptors on rat spinal neurons mediate complex facilitatory and inhibitory effects. <i>European Journal of Neuroscience</i> , <b>2003</b> , 18, 1873-83	3.5	23
34	Modulation of neuronal nicotinic receptor function by the neuropeptides CGRP and substance P on autonomic nerve cells. <i>British Journal of Pharmacology</i> , <b>2003</b> , 139, 1061-73	8.6	37
33	A structural model of agonist binding to the alpha3beta4 neuronal nicotinic receptor. <i>British Journal of Pharmacology</i> , <b>2003</b> , 140, 921-31	8.6	47
32	Pre and postsynaptic effects of metabotropic glutamate receptor activation on neonatal rat hypoglossal motoneurons. <i>Neuroscience Letters</i> , <b>2003</b> , 338, 9-9	3.3	
31	Modulation of P2X3 receptors by Mg <sup>2+</sup> on rat DRG neurons in culture. <i>Neuropharmacology</i> , <b>2003</b> , 44, 132-40	5.5	25
30	The ATP-mediated fast current of rat dorsal root ganglion neurons is a novel effector for GABA(B) receptor activation. <i>Neuroscience Letters</i> , <b>2003</b> , 338, 181-4	3.3	22
29	Effects of caffeine on the excitability and intracellular Ca <sup>2+</sup> transients of neonatal rat hypoglossal motoneurons in vitro. <i>Neuroscience Letters</i> , <b>2003</b> , 346, 177-81	3.3	7
28	Bimodal action of protons on ATP currents of rat PC12 cells. <i>Journal of General Physiology</i> , <b>2003</b> , 122, 33-44	3.4	10
27	Experimental and modeling studies of novel bursts induced by blocking Na <sup>+</sup> pump and synaptic inhibition in the rat spinal cord. <i>Journal of Neurophysiology</i> , <b>2002</b> , 88, 676-91	3.2	28
26	Inhibition of spinal or hypoglossal motoneurons of the newborn rat by glycine or GABA. <i>European Journal of Neuroscience</i> , <b>2002</b> , 15, 975-83	3.5	53
25	Expression and dendritic mRNA localization of GABAC receptor rho1 and rho2 subunits in developing rat brain and spinal cord. <i>European Journal of Neuroscience</i> , <b>2002</b> , 15, 1747-58	3.5	77
24	Enhancement of neuronal nicotinic receptor activity of rat chromaffin cells by a novel class of peptides. <i>Annals of the New York Academy of Sciences</i> , <b>2002</b> , 971, 100-7	6.5	7
23	A novel class of peptides with facilitating action on neuronal nicotinic receptors of rat chromaffin cells in vitro: functional and molecular dynamics studies. <i>Molecular Pharmacology</i> , <b>2002</b> , 61, 43-54	4.3	17
22	Calibration of agonist concentrations applied by pressure pulses or via rapid solution exchanger. <i>Journal of Neuroscience Methods</i> , <b>2001</b> , 110, 155-61	3	26
21	Characteristics of fast Na <sup>+</sup> current of hypoglossal motoneurons in a rat brainstem slice preparation. <i>European Journal of Neuroscience</i> , <b>2001</b> , 13, 763-72	3.5	10
20	Alternating rhythmic activity induced by dorsal root stimulation in the neonatal rat spinal cord in vitro. <i>Journal of Physiology</i> , <b>2001</b> , 530, 105-12	3.9	83



19	Evidence for increased extracellular K(+) as an important mechanism for dorsal root induced alternating rhythmic activity in the neonatal rat spinal cord in vitro. <i>Neuroscience Letters</i> , <b>2001</b> , 304, 77-80 <sup>33</sup>		17
18	Depression of windup of spinal neurons in the neonatal rat spinal cord in vitro by an NK3 tachykinin receptor antagonist. <i>Journal of Neurophysiology</i> , <b>2001</b> , 85, 1502-11	3.2	31
17	Negative cross talk between anionic GABAA and cationic P2X ionotropic receptors of rat dorsal root ganglion neurons. <i>Journal of Neuroscience</i> , <b>2001</b> , 21, 4958-68	6.6	98
16	Differential short-term changes in GABAergic or glycinergic synaptic efficacy on rat hypoglossal motoneurons. <i>Journal of Neurophysiology</i> , <b>2001</b> , 86, 565-74	3.2	11
15	Facilitation of cholinergic transmission by substance P methyl ester in the mouse hippocampal slice preparation. <i>European Journal of Neuroscience</i> , <b>2000</b> , 12, 585-94	3.5	13
14	Antagonism of nicotinic receptors of rat chromaffin cells by N,N, N-trimethyl-1-(4-trans-stilbenoxy)-2-propylammonium iodide: a patch clamp and ligand binding study. <i>British Journal of Pharmacology</i> , <b>2000</b> , 129, 1771-9	8.6	17
13	Opposite changes in synaptic activity of organotypic rat spinal cord cultures after chronic block of AMPA/kainate or glycine and GABAA receptors. <i>Journal of Physiology</i> , <b>2000</b> , 523 Pt 3, 639-51	3.9	52
12	Current and voltage clamp studies of the spike medium afterhyperpolarization of hypoglossal motoneurons in a rat brain stem slice preparation. <i>Journal of Neurophysiology</i> , <b>2000</b> , 83, 2987-95	3.2	63
11	Relative contribution by GABA or glycine to Cl(-)-mediated synaptic transmission on rat hypoglossal motoneurons in vitro. <i>Journal of Neurophysiology</i> , <b>2000</b> , 84, 2715-24	3.2	78
10	Rapid relief of block by mecamylamine of neuronal nicotinic acetylcholine receptors of rat chromaffin cells in vitro: an electrophysiological and modeling study. <i>Molecular Pharmacology</i> , <b>2000</b> , 58, 778-87	4.3	40
9	Voltage-activated K+ currents of hypoglossal motoneurons in a brain stem slice preparation from the neonatal rat. <i>Journal of Neurophysiology</i> , <b>1999</b> , 81, 140-8	3.2	21
8	Generation of rhythmic patterns of activity by ventral interneurons in rat organotypic spinal slice culture. <i>Journal of Physiology</i> , <b>1999</b> , 517 ( Pt 2), 459-75	3.9	51
7	Modulation by substance P of synaptic transmission in the mouse hippocampal slice. <i>European Journal of Neuroscience</i> , <b>1998</b> , 10, 3076-84	3.5	27
6	Pharmacological block of the electrogenic sodium pump disrupts rhythmic bursting induced by strychnine and bicuculline in the neonatal rat spinal cord. <i>Journal of Neurophysiology</i> , <b>1997</b> , 77, 17-23	3.2	62
5	Localization of rhythmogenic networks responsible for spontaneous bursts induced by strychnine and bicuculline in the rat isolated spinal cord. <i>Journal of Neuroscience</i> , <b>1996</b> , 16, 7063-76	6.6	128
4	Effects of thyrotropin-releasing hormone on GABAergic synaptic transmission of the rat hippocampus. <i>European Journal of Neuroscience</i> , <b>1996</b> , 8, 1299-305	3.5	10
3	Membrane potential oscillations of neonatal rat spinal motoneurons evoked by electrical stimulation of dorsal root fibres. <i>European Journal of Neuroscience</i> , <b>1995</b> , 7, 2403-8	3.5	11
2	Desensitization of AMPA receptors limits the amplitude of EPSPs and the excitability of motoneurons of the rat isolated spinal cord. <i>European Journal of Neuroscience</i> , <b>1995</b> , 7, 1229-34	3.5	34

- 1 GABA receptor mechanisms in the central nervous system. *Progress in Neurobiology*, **1991**, 36, 35-92 10.9 471