
Xulin Xu

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2700200/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Pharmacologic activation of cholinergic alpha7 nicotinic receptors mitigates depressive-like behavior in a mouse model of chronic stress. Journal of Neuroinflammation, 2017, 14, 234.	7.2	56
2	Flupirtine attenuates chronic restraint stress-induced cognitive deficits and hippocampal apoptosis in male mice. Behavioural Brain Research, 2015, 288, 1-10.	2.2	51
3	Long-lasting spatial learning and memory impairments caused by chronic cerebral hypoperfusion associate with a dynamic change of HCN1/HCN2 expression in hippocampal CA1 region. Neurobiology of Learning and Memory, 2015, 123, 72-83.	1.9	29
4	Baclofen ameliorates spatial working memory impairments induced by chronic cerebral hypoperfusion via up-regulation of HCN2 expression in the PFC in rats. Behavioural Brain Research, 2016, 308, 6-13.	2.2	23
5	Anti-thrombotic effects mediated by dihydromyricetin involve both platelet inhibition and endothelial protection. Pharmacological Research, 2021, 167, 105540.	7.1	21
6	Pathophysiological roles of cell surface and extracellular protein disulfide isomerase and their molecular mechanisms. British Journal of Pharmacology, 2021, 178, 2911-2930.	5.4	20
7	Clonidine ameliorates cognitive impairment induced by chronic cerebral hypoperfusion via up-regulation of the GABABR1 and GAD67 in hippocampal CA1 in rats. Pharmacology Biochemistry and Behavior, 2015, 132, 96-102.	2.9	17
8	Activity-dependent downregulation of M-Type (Kv7) K+ channels surface expression requires the activation of iGluRs/Ca2+/PKC signaling pathway in hippocampal neuron. Neuropharmacology, 2015, 95, 154-167.	4.1	9
9	Fluoxetine ameliorates cognitive impairments induced by chronic cerebral hypoperfusion via down-regulation of HCN2 surface expression in the hippocampal CA1 area in rats. Pharmacology Biochemistry and Behavior, 2016, 140, 1-7.	2.9	8