

# Haiyong Wang

## List of Publications by Citations

**Source:** <https://exaly.com/author-pdf/2710024/haiyong-wang-publications-by-citations.pdf>

**Version:** 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

48  
papers

370  
citations

10  
h-index

17  
g-index

52  
ext. papers

498  
ext. citations

3.6  
avg, IF

3.96  
L-index

#	Paper	IF	Citations
48	The prognosis analysis of different metastasis pattern in patients with different breast cancer subtypes: a SEER based study. <i>Oncotarget</i> , <b>2017</b> , 8, 26368-26379	3.3	40
47	Screening differential circular RNA expression profiles reveal that hsa_circ_0128298 is a biomarker in the diagnosis and prognosis of hepatocellular carcinoma. <i>Cancer Management and Research</i> , <b>2018</b> , 10, 1275-1283	3.6	40
46	Significant efficacy and well safety of apatinib in an advanced liver cancer patient: a case report and literature review. <i>Oncotarget</i> , <b>2017</b> , 8, 20510-20515	3.3	34
45	Circular RNA Hsa_Circ_0091579 Serves as a Diagnostic and Prognostic Marker for Hepatocellular Carcinoma. <i>Cellular Physiology and Biochemistry</i> , <b>2018</b> , 51, 290-300	3.9	30
44	Circular RNA hsa_circ_0078602 may have potential as a prognostic biomarker for patients with hepatocellular carcinoma. <i>Oncology Letters</i> , <b>2019</b> , 17, 2091-2098	2.6	20
43	The prognostic analysis of different metastatic patterns in extensive-stage small-cell lung cancer patients: a large population-based study. <i>Future Oncology</i> , <b>2018</b> , 14, 1397-1407	3.6	19
42	Risk factors of brain metastasis during the course of EGFR-TKIs therapy for patients with EGFR-mutated advanced lung adenocarcinoma. <i>Oncotarget</i> , <b>2016</b> , 7, 81906-81917	3.3	17
41	Total lymphocyte count, neutrophil-lymphocyte ratio, and platelet-lymphocyte ratio as prognostic factors in advanced non-small cell lung cancer with chemoradiotherapy. <i>Cancer Management and Research</i> , <b>2018</b> , 10, 6677-6683	3.6	17
40	CD44, a marker of cancer stem cells, is positively correlated with PD-L1 expression and immune cells infiltration in lung adenocarcinoma. <i>Cancer Cell International</i> , <b>2020</b> , 20, 583	6.4	11
39	Preoperative radiation may improve the outcomes of resectable IIIA/N2 non-small-cell lung cancer patients: A propensity score matching-based analysis from surveillance, epidemiology, and end results database. <i>Cancer Medicine</i> , <b>2018</b> , 7, 4354-4360	4.8	11
38	Better cancer specific survival in young small cell lung cancer patients especially with AJCC stage III. <i>Oncotarget</i> , <b>2017</b> , 8, 34923-34934	3.3	10
37	TERT mutations correlate with higher TMB value and unique tumor microenvironment and may be a potential biomarker for anti-CTLA4 treatment. <i>Cancer Medicine</i> , <b>2020</b> , 9, 7151-7160	4.8	9
36	KEAP1-NFE2L2-Mutant NSCLC and Immune Checkpoint Inhibitors: A Large Database Analysis. <i>Journal of Thoracic Oncology</i> , <b>2020</b> , 15, e85-e86	8.9	8
35	Better survival in PMRT of female breast cancer patients with >5 negative lymph nodes: A population-based study. <i>Medicine (United States)</i> , <b>2017</b> , 96, e5998	1.8	6
34	Thoracic radiation therapy could give survival benefit to elderly patients with extensive-stage small-cell lung cancer. <i>Future Oncology</i> , <b>2017</b> , 13, 1149-1158	3.6	6
33	A 5-Gene Signature Is Closely Related to Tumor Immune Microenvironment and Predicts the Prognosis of Patients with Non-Small Cell Lung Cancer. <i>BioMed Research International</i> , <b>2020</b> , 2020, 2147397	3.97	6
32	Should all breast cancer patients with four or more positive lymph nodes who underwent modified radical mastectomy be treated with postoperative radiotherapy? A population-based study. <i>Oncotarget</i> , <b>2016</b> , 7, 75492-75502	3.3	6

31	Efficacy of cisplatin/pemetrexed with bevacizumab to treat advanced lung adenocarcinoma with different drive genes: case report and literature review. <i>OncoTargets and Therapy</i> , <b>2016</b> , 9, 4639-44	4.4	6
30	Male patients with resected IIIA-N2 non-small-cell lung cancer may benefit from postoperative radiotherapy: a population-based survival analysis. <i>Future Oncology</i> , <b>2018</b> , 14, 2371-2381	3.6	6
29	Including positive lymph node count in the AJCC N staging may be a better predictor of the prognosis of NSCLC patients, especially stage III patients: a large population-based study. <i>International Journal of Clinical Oncology</i> , <b>2019</b> , 24, 1359-1366	4.2	5
28	PDL1 high expression without TP53, KEAP1 and EPHA5 mutations could better predict survival for patients with NSCLC receiving atezolizumab. <i>Lung Cancer</i> , <b>2021</b> , 151, 76-83	5.9	5
27	PLNR0% may be a benefit from PORT for patients with IIIA-N2 NSCLC: a large population-based study. <i>Cancer Management and Research</i> , <b>2018</b> , 10, 3561-3567	3.6	5
26	Postmastectomy radiation therapy for breast cancer patients with one to three positive lymph nodes: a propensity score matching analysis. <i>Future Oncology</i> , <b>2017</b> , 13, 1395-1404	3.6	4
25	Greater efficacy of intracavitary infusion of bevacizumab compared to traditional local treatments for patients with malignant cavity serous effusion. <i>Oncotarget</i> , <b>2017</b> , 8, 35262-35271	3.3	4
24	Tumor Primary Location May Affect Metastasis Pattern for Patients with Stage IV NSCLC: A Population-Based Study. <i>Journal of Oncology</i> , <b>2020</b> , 2020, 4784701	4.5	4
23	Poor survival of non-small-cell lung cancer patients with main bronchus tumor: a large population-based study. <i>Future Oncology</i> , <b>2019</b> , 15, 2819-2827	3.6	4
22	Previous Radiotherapy Increases the Efficacy of IL-2 in Malignant Pleural Effusion: Potential Evidence of a Radio-Memory Effect?. <i>Frontiers in Immunology</i> , <b>2018</b> , 9, 2916	8.4	4
21	Patients With -Mutant NSCLC May Not Benefit From Immune Checkpoint Inhibitors: A Population-Based Study. <i>JTO Clinical and Research Reports</i> , <b>2020</b> , 1, 100006	1.4	3
20	A Novel Nomogram including AJCC Stages Could Better Predict Survival for NSCLC Patients Who Underwent Surgery: A Large Population-Based Study. <i>Journal of Oncology</i> , <b>2020</b> , 2020, 7863984	4.5	3
19	Great efficacy of bevacizumab plus erlotinib for leptomeningeal metastases from non-small cell lung cancer with initially positive EGFR mutation: a case report. <i>Cancer Biology and Therapy</i> , <b>2018</b> , 19, 1073-1077	4.6	3
18	Application of prophylactic cranial irradiation in limited-stage small-cell lung cancer: which patients could benefit?. <i>Future Oncology</i> , <b>2019</b> , 15, 3237-3245	3.6	3
17	The appropriate number of ELNs for lymph node negative breast cancer patients underwent MRM: a population-based study. <i>Oncotarget</i> , <b>2017</b> , 8, 65668-65676	3.3	3
16	High level of programmed death ligand 1 (PD-L1) predicts longer survival in patients with resectable small cell lung cancer. <i>International Journal of Clinical and Experimental Pathology</i> , <b>2018</b> , 11, 2675-2682	1.4	3
15	Optimizing intrapleural bevacizumab dosing in non-small-cell lung cancer-mediated malignant pleural effusion: less is more. <i>Future Oncology</i> , <b>2018</b> , 14, 2131-2138	3.6	2
14	Radiotherapy may improve survival of ES-SCLC with distant metastasis only for patients with one metastatic site: A population-based study. <i>Oncology Letters</i> , <b>2020</b> , 19, 139-146	2.6	2

13	Duration of immunotherapy in patients with advanced lung adenocarcinoma with negative driver genes: case report and literature review. <i>Thoracic Cancer</i> , <b>2020</b> , 11, 3001-3006	3.2	2
12	Surgical intervention may be a therapeutic option for NSCLC patients with AJCC stage IV: a large population-based study. <i>Cancer Management and Research</i> , <b>2018</b> , 10, 3219-3226	3.6	2
11	Incorporating the Number of PLN into the AJCC Stage Could Better Predict the Survival for Patients with NSCLC: A Large Population-Based Study. <i>Journal of Oncology</i> , <b>2020</b> , 2020, 1087237	4.5	1
10	A 5-Genomic Mutation Signature Can Predict the Survival for Patients With NSCLC Receiving Atezolizumab. <i>Frontiers in Immunology</i> , <b>2021</b> , 12, 606027	8.4	1
9	Young male breast cancer, a small crowd, the survival, and prognosis?: A population-based study. <i>Medicine (United States)</i> , <b>2018</b> , 97, e12686	1.8	1
8	Characteristics and survival difference of clinical tumor size 0 extensive-stage small cell lung cancer with different metastasis pattern. <i>Journal of Thoracic Disease</i> , <b>2018</b> , 10, 5414-5420	2.6	1
7	Better efficacy of intrapleural infusion of bevacizumab with pemetrexed for malignant pleural effusion mediated from nonsquamous non-small cell lung cancer. <i>OncoTargets and Therapy</i> , <b>2018</b> , 11, 8421-8426	4.4	1
6	How many ELNs are optimal for breast cancer patients with more than three PLNs who underwent MRM? A large population-based study. <i>OncoTargets and Therapy</i> , <b>2018</b> , 11, 1005-1011	4.4	1
5	A clinical variable-based nomogram could predict the survival for advanced NSCLC patients receiving second-line atezolizumab. <i>Cancer Medicine</i> , <b>2021</b> , 10, 6218-6226	4.8	1
4	Clinicopathological Difference and Survival Impact of Patients with c-SCLC and SCLC. <i>International Journal of General Medicine</i> , <b>2021</b> , 14, 6899-6906	2.3	0
3	Genomic Variations and Immune-Related Features of TMB, PD-L1 Expression and CD8 T Cell Infiltration in Chinese Pulmonary Sarcomatoid Carcinoma.. <i>International Journal of General Medicine</i> , <b>2022</b> , 15, 4209-4220	2.3	0
2	Kinetic change of serum carcinoembryonic antigen can early predict progression in patients with metastatic non-small cell lung cancer during maintenance therapy with bevacizumab plus pemetrexed. <i>Oncotarget</i> , <b>2017</b> , 8, 74910-74916	3.3	
1	A prognostic score model to determine which breast cancer patients with 1-3 positive lymph nodes after modified radical mastectomy should receive radiotherapy. <i>Oncotarget</i> , <b>2018</b> , 9, 385-393	3.3	