

# Krzysztof Hryniewicz

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2712918/publications.pdf>

Version: 2024-02-01

21  
papers

335  
citations

840776

11  
h-index

940533

16  
g-index

21  
all docs

21  
docs citations

21  
times ranked

277  
citing authors

#	ARTICLE	IF	CITATIONS
1	Formation, diagenesis and fauna of cold seep carbonates from the Miocene Taishu Group of Tsushima (Japan). <i>Geological Magazine</i> , 2021, 158, 964-984.	1.5	6
2	Taphonomy and palaeoecology of deep-water chemosymbiotic bivalves from the Eocene of Outer Eastern Carpathians, Ukraine. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2020, 553, 109782.	2.3	2
3	Mesozoic marine reptiles from Spitsbergen and their ecosystems. <i>Geology Today</i> , 2019, 35, 20-25.	0.9	2
4	Early Devonian bivalves from Hamar Laghdad, Morocco. <i>Neues Jahrbuch Fur Geologie Und Palaontologie - Abhandlungen</i> , 2018, 290, 191-202.	0.4	1
5	New bivalves from a Middle Devonian methane seep in Morocco: the oldest record of repetitive shell morphologies among some seep bivalve molluscs. <i>Journal of Systematic Palaeontology</i> , 2017, 15, 19-41.	1.5	19
6	Extensive Early Cretaceous (Albian) methane seepage on Ellef Ringnes Island, Canadian High Arctic. <i>Bulletin of the Geological Society of America</i> , 2017, 129, 788-805.	3.3	17
7	Mass occurrence of seep-specific bivalves in the oldest-known cold seep metazoan community. <i>Scientific Reports</i> , 2017, 7, 14292.	3.3	22
8	Gastropods from the Late Jurassic–Early Cretaceous seep deposits in Spitsbergen, Svalbard. <i>Zootaxa</i> , 2017, 4329, 351.	0.5	6
9	Paleocene methane seep and wood-fall marine environments from Spitsbergen, Svalbard. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2016, 462, 41-56.	2.3	14
10	Late Jurassic–Early Cretaceous hydrocarbon seep boulders from Novaya Zemlya and their faunas. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2015, 436, 231-244.	2.3	22
11	The palaeoecology of the latest Jurassic–earliest Cretaceous hydrocarbon seep carbonates from Spitsbergen, Svalbard. <i>Lethaia</i> , 2015, 48, 353-374.	1.4	31
12	Brachiopods from Late Jurassic–Early Cretaceous hydrocarbon seep deposits, central Spitsbergen, Svalbard. <i>Zootaxa</i> , 2014, 3884, 501-32.	0.5	10
13	&lt;p&gt;&lt;strong&gt;Bivalves from the latest Jurassic-earliest Cretaceous hydrocarbon seep carbonates from central Spitsbergen, Svalbard&lt;/strong&gt;&lt;/p&gt;. <i>Zootaxa</i> , 2014, 3859, 1.	0.5	21
14	A Boreal serpulid fauna from Volgian-Ryazanian (latest Jurassic-earliest Cretaceous) shelf sediments and hydrocarbon seeps from Svalbard. <i>Geodiversitas</i> , 2014, 36, 527-540.	0.8	16
15	Shallow water methane-derived authigenic carbonate mounds at the Codling Fault Zone, western Irish Sea. <i>Marine Geology</i> , 2014, 357, 139-150.	2.1	23
16	Hydrocarbon seeps from close to the Jurassic–Cretaceous boundary, Svalbard. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2011, 306, 15-26.	2.3	71
17	Ammonites from hydrocarbon seep carbonate bodies from the uppermost Jurassic - lowermost Cretaceous of Spitsbergen and their biostratigraphical importance. <i>Neues Jahrbuch Fur Geologie Und Palaontologie - Abhandlungen</i> , 2011, 262, 267-288.	0.4	24
18	Large onychites (cephalopod hooks) from the Upper Jurassic of the Boreal Realm. <i>Acta Palaeontologica Polonica</i> , 0, , .	0.4	2

#	ARTICLE	IF	CITATIONS
19	A new sediment-dwelling pholadid bivalve from Oligocene glaciomarine sediments of King George Island, West Antarctica. <i>Acta Palaeontologica Polonica</i> , 0, 61, .	0.4	0
20	Thyasirid bivalves from Cretaceous and Paleogene cold seeps. <i>Acta Palaeontologica Polonica</i> , 0, 62, .	0.4	13
21	A late Paleocene fauna from shallow-water chemosynthesis-based ecosystems, Spitsbergen, Svalbard. <i>Acta Palaeontologica Polonica</i> , 0, 64, .	0.4	13