

Nikolai Albert

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2723235/publications.pdf>

Version: 2024-02-01

28
papers

882
citations

687363

13
h-index

552781

26
g-index

29
all docs

29
docs citations

29
times ranked

1416
citing authors

#	ARTICLE	IF	CITATIONS
1	Discontinuation of antipsychotics in individuals with first-episode schizophrenia and its association to functional outcomes, hospitalization and death: a register-based nationwide follow-up study. <i>Psychological Medicine</i> , 2023, 53, 5033-5041.	4.5	2
2	Associations between duration of untreated psychosis and domains of positive and negative symptoms persist after 10 years of follow-up: A secondary analysis from the OPUS trial. <i>Schizophrenia Research</i> , 2021, 228, 575-580.	2.0	4
3	Reversibility of Antipsychotic-Induced Weight Gain: A Systematic Review and Meta-Analysis. <i>Frontiers in Endocrinology</i> , 2021, 12, 577919.	3.5	9
4	The Danish OPUS Early Intervention Services for First-Episode Psychosis: A Phase 4 Prospective Cohort Study With Comparison of Randomized Trial and Real-World Data. <i>American Journal of Psychiatry</i> , 2021, 178, 941-951.	7.2	14
5	Repetitive transcranial magnetic stimulation and transcranial direct current stimulation for auditory hallucinations in schizophrenia: Systematic review and meta-analysis. <i>Journal of Psychiatric Research</i> , 2021, 143, 163-175.	3.1	16
6	Discontinuation of antipsychotic medication – time to rethink trial design. <i>Lancet Psychiatry</i> , 2020, 7, 841-842.	7.4	7
7	Future directions in risk research. , 2020, , 417-430.		0
8	Cognitive functioning following discontinuation of antipsychotic medication. A naturalistic sub-group analysis from the OPUS II trial. <i>Psychological Medicine</i> , 2019, 49, 1138-1147.	4.5	20
9	The outcome of early intervention in first episode psychosis. <i>International Review of Psychiatry</i> , 2019, 31, 413-424.	2.8	36
10	Development and validation of multivariable prediction models of remission, recovery, and quality of life outcomes in people with first episode psychosis: a machine learning approach. <i>The Lancet Digital Health</i> , 2019, 1, e261-e270.	12.3	33
11	Emotion recognition latency, but not accuracy, relates to real life functioning in individuals at ultra-high risk for psychosis. <i>Schizophrenia Research</i> , 2019, 210, 197-202.	2.0	13
12	Development of a fidelity scale for Danish specialized early interventions service. <i>Microbial Biotechnology</i> , 2019, 13, 568-573.	1.7	7
13	Programme fidelity of specialized early intervention in Denmark. <i>Microbial Biotechnology</i> , 2019, 13, 627-632.	1.7	9
14	Attenuated Psychosis Syndromes Among Danish Youth and Young Adults: Early Identification and Intervention. , 2019, , 333-348.		0
15	Early Intervention Service for Young People With Psychosis. <i>JAMA Psychiatry</i> , 2018, 75, 427.	11.0	9
16	Association of Substance Use Disorders With Conversion From Schizotypal Disorder to Schizophrenia. <i>JAMA Psychiatry</i> , 2018, 75, 733.	11.0	21
17	Social functioning in patients with first-episode psychosis. <i>Lancet Psychiatry</i> , 2018, 5, 3-4.	7.4	2
18	T54. TAILOR – TAPERED DISCONTINUATION VERSUS MAINTENANCE THERAPY OF ANTIPSYCHOTIC MEDICATION IN PATIENTS WITH NEWLY DIAGNOSED SCHIZOPHRENIA SPECTRUM DISORDERS IN REMISSION OF PSYCHOTIC SYMPTOMS. <i>Schizophrenia Bulletin</i> , 2018, 44, S134-S135.	4.3	2

#	ARTICLE	IF	CITATIONS
19	Stability and development of psychotic symptoms and the use of antipsychotic medication – long-term follow-up. <i>Psychological Medicine</i> , 2017, 47, 2118-2129.	4.5	22
20	The effect of duration of untreated psychosis and treatment delay on the outcomes of prolonged early intervention in psychotic disorders. <i>NPJ Schizophrenia</i> , 2017, 3, 34.	3.6	61
21	Antipsychotic medication and remission of psychotic symptoms 10 years after a first-episode psychosis. <i>Schizophrenia Research</i> , 2017, 182, 42-48.	2.0	80
22	Course of illness in a sample of patients diagnosed with a schizotypal disorder and treated in a specialized early intervention setting. Findings from the 3.5 year follow-up of the OPUS II study. <i>Schizophrenia Research</i> , 2017, 182, 24-30.	2.0	16
23	Early intervention services are effective and must be defended. <i>World Psychiatry</i> , 2017, 16, 272-274.	10.4	3
24	Five years of specialised early intervention versus two years of specialised early intervention followed by three years of standard treatment for patients with a first episode psychosis: randomised, superiority, parallel group trial in Denmark (OPUS II). <i>BMJ: British Medical Journal</i> , 2017, 356, i6681.	2.3	79
25	TAILOR – tapered discontinuation versus maintenance therapy of antipsychotic medication in patients with newly diagnosed schizophrenia or persistent delusional disorder in remission of psychotic symptoms: study protocol for a randomized clinical trial. <i>Trials</i> , 2017, 18, 445.	1.6	26
26	Gender differences in first-episode psychosis at 5-year follow-up – two different courses of disease? Results from the OPUS study at 5-year follow-up. <i>European Psychiatry</i> , 2014, 29, 44-51.	0.2	102
27	Predictors of recovery in first episode psychosis: The OPUS cohort at 10year follow-up. <i>Schizophrenia Research</i> , 2013, 150, 163-168.	2.0	144
28	Predictors of recovery from psychosis. <i>Schizophrenia Research</i> , 2011, 125, 257-266.	2.0	143