

Anthony W Addison

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2733175/publications.pdf>

Version: 2024-02-01

104
papers

13,506
citations

109137

35
h-index

35952

97
g-index

108
all docs

108
docs citations

108
times ranked

11804
citing authors

#	ARTICLE	IF	CITATIONS
1	Synthesis, structure, and spectroscopic properties of copper(II) compounds containing nitrogen-sulphur donor ligands; the crystal and molecular structure of aqua[1,7-bis(N-methylbenzimidazol-2-yl)-2,6-dithiaheptane]copper(II) perchlorate. <i>Journal of the Chemical Society Dalton Transactions</i> , 1984, , 1349-1356.	1.1	8,412
2	Conversion constants for redox potentials measured versus different reference electrodes in acetonitrile solutions at 25°C. <i>Inorganica Chimica Acta</i> , 2000, 298, 97-102.	1.2	1,281
3	Spectroscopic and redox studies of some copper(II) complexes with biomimetic donor atoms: implications for protein copper centres. <i>Journal of the Chemical Society Dalton Transactions</i> , 1979, , 600.	1.1	459
4	Synthesis of some imidazole- and pyrazole-derived chelating agents. <i>Journal of Heterocyclic Chemistry</i> , 1981, 18, 803-805.	1.4	320
5	Copper complexes of the "tripod" ligand tris(2-benzimidazolymethyl)amine: five- and six-coordinate copper(II) derivatives and some copper(I) derivatives. <i>Inorganic Chemistry</i> , 1981, 20, 103-110.	1.9	154
6	Iridium Dihydroxybipyridine Complexes Show That Ligand Deprotonation Dramatically Speeds Rates of Catalytic Water Oxidation. <i>Inorganic Chemistry</i> , 2013, 52, 9175-9183.	1.9	142
7	Dipicolylamine Complexes of Copper(II): Two Different Coordination Geometries in the Same Unit Cell of Cu(Dipica) ₂ (BF ₄) ₂ . <i>Inorganic Chemistry</i> , 1996, 35, 467-471.	1.9	110
8	Nitrosyliron(III) hemoglobin: autoreduction and spectroscopy. <i>Biochemistry</i> , 1986, 25, 4104-4113.	1.2	105
9	Synthesis of some benzimidazole- and benzothiazole-derived ligand systems and their precursory diacids. <i>Journal of Heterocyclic Chemistry</i> , 1983, 20, 1481-1484.	1.4	100
10	New iron(II) spin-crossover complexes with heterocyclic amine-derived ligands and STEPS experiments on photogenerated metastable high-spin states. <i>Journal of the Chemical Society Dalton Transactions</i> , 1987, , 2621.	1.1	97
11	Spectroscopy and structure of thiolate and thioether complexes of copper(II) and the relationship of their redox chemistry to that of certain copper proteins. <i>Inorganic Chemistry</i> , 1984, 23, 1957-1967.	1.9	96
12	Small molecule analogs for the specific iron-binding site of lactoferrin: a single-crystal x-ray structure of bis(methanol)bis[2-(5-methylpyrazol-3-yl)phenolato]iron(III) nitrate-methanol and spectroscopic studies on iron(III) phenolate complexes. <i>Inorganic Chemistry</i> , 1980, 19, 3655-3663.	1.9	95
13	Pentacoordinate copper complexes of nitrogen-sulfur donors: structural chemistry of two complexes of bis(2-(2-benzimidazolyl)ethyl) sulfide with the sulfur alternatively in equatorial and axial coordination modes. <i>Inorganic Chemistry</i> , 1983, 22, 3645-3653.	1.9	87
14	Is ligand topology an influence on the redox potentials of copper complexes?. <i>Inorganica Chimica Acta</i> , 1989, 162, 217-220.	1.2	85
15	Copper(II) and Nickel(II) Complexes of Dianionic and Tetraanionic Dinucleating Macrocycles. <i>Inorganic Chemistry</i> , 1998, 37, 1028-1036.	1.9	84
16	A stable bis(thiolate) of copper(II) with long axial copper-sulfur linkages: crystal and molecular structure of trans-[Cu(cyclam)(SC ₆ F ₅) ₂]. <i>Inorganic Chemistry</i> , 1983, 22, 1225-1228.	1.9	65
17	Synthesis and proton transfer-linked redox tuning of ruthenium(II) complexes with tridentate 2,6-bis(benzimidazol-2-yl)pyridine ligands. <i>Journal of the Chemical Society Dalton Transactions</i> , 1993, , 2477.	1.1	65
18	Complexes of structural analogues of terpyridyl with iron and zinc; the x-ray crystal structure of bis[2,6-bis(benzimidazol-2-yl)pyridine]iron(II) trifluoromethylsulphonate bis-ethanol solvate. <i>Polyhedron</i> , 1992, 11, 635-646.	1.0	62

#	ARTICLE	IF	CITATIONS
19	Mono- and Trinuclear Nickel(II) Complexes with Sulfur-Containing Oxime Ligands: An Uncommon Templated Coupling of Oxime with Nitrile. <i>Inorganic Chemistry</i> , 1999, 38, 1759-1766.	1.9	61
20	Copper(II) and zinc(II) co-ordination compounds of tridentate bis(benzimidazole)pyridine ligands. Crystal and molecular structures of bis[2,6-bis(1-methylbenzimidazol-2-yl)pyridine]copper(II) diperchlorate monohydrate and (acetonitrile)[2,6-bis(benzimidazol-2-yl)pyridine](perchlorato)copper(II) perchlorate. <i>Journal of the Chemical Society Dalton Transactions</i> , 1988, , 1429-1435.	1.1	54
21	A Tetrameric Nickel(II) μ_3 -Chair with both Antiferromagnetic Internal Coupling and Ferromagnetic Spin Alignment. <i>Angewandte Chemie - International Edition</i> , 2001, 40, 4734-4737.	7.2	53
22	A Triple-Decker Heptadecanuclear (Cu ^{II}) ₁₅ (Cr ^{III}) ₂ Complex Assembled from Pentanuclear Metallacrowns. <i>European Journal of Inorganic Chemistry</i> , 2010, 2010, 4851-4858.	1.0	51
23	Structure of trimeric haemerythrin. <i>Nature</i> , 1983, 303, 86-88.	13.7	50
24	Comparison of hemerythrins from four species of sipunculids by optical absorption, circular dichroism, fluorescence emission, and resonance Raman spectroscopy. <i>Biochemistry</i> , 1977, 16, 1743-1749.	1.2	49
25	Structural Demonstration of the Role of Ligand Framework Conformability in Copper(II)/Copper(I) Redox Potentials. <i>Inorganic Chemistry</i> , 1997, 36, 134-135.	1.9	49
26	Formation of Coordination Polymers or Discrete Adducts via Reactions of Gadolinium(III)–Copper(II) 15-Metallacrown-5 Complexes with Polycarboxylates: Synthesis, Structures and Magnetic Properties. <i>Inorganic Chemistry</i> , 2014, 53, 1320-1330.	1.9	49
27	Nickel(II) complexes with dithiadiminoamine and dithiabis(thiosemicarbazone) ligands. <i>Dalton Transactions RSC</i> , 2000, , 335-341.	2.3	48
28	Manifestation of π - π Stacking Interactions in Luminescence Properties and Energy Transfer in Aromatically-Derived Tb, Eu and Gd Tris(pyrazolyl)borate Complexes. <i>Inorganic Chemistry</i> , 2015, 54, 3125-3133.	1.9	48
29	Mononuclear and Mixed-Valence Binuclear Oxovanadium Complexes with Benzimidazole-Derived Chelating Agents. <i>Inorganic Chemistry</i> , 2002, 41, 2243-2249.	1.9	47
30	Magnetic and Sorption Properties of Supramolecular Systems Based on Pentanuclear Copper(II) 12-Metallacrown-4 Complexes and Isomeric Phthalates: Structural Modeling of the Different Stages of Alcohol Sorption. <i>European Journal of Inorganic Chemistry</i> , 2011, 2011, 4826-4836.	1.0	47
31	Structural aspects of the bis(2,2-dipicolylamine)iron(II) cation. <i>Inorganica Chimica Acta</i> , 1989, 158, 211-215.	1.2	43
32	Structural, magnetic and related attributes of some oximate-bridged tetranuclear nickel(ii) rhombs and a dinuclear congener. Electronic supplementary information (ESI) available: mass spectra, χ_T vs. T, response of magnetic properties, low-lying spin levels and UV-VIS data. See http://www.rsc.org/suppdata/dt/b3/b300539a/ . <i>Dalton Transactions</i> , 2003, , 1587-1595.	1.6	43
33	Helical Antiferromagnetic Copper(II) Chains with a Collagen Structural Motif. <i>Inorganic Chemistry</i> , 1996, 35, 5966-5967.	1.9	42
34	Pentadentate thioether-oxime macrocyclic and quasi-macrocyclic complexes of copper(II) and nickel(II). <i>Inorganica Chimica Acta</i> , 2000, 300-302, 992-1003.	1.2	41
35	Analogs for the specific iron-binding site in the transferrins: molecular structure of a ternary iron(III) model complex and spectroscopic, redox and reactivity properties of related compounds. <i>Inorganic Chemistry</i> , 1990, 29, 3425-3433.	1.9	37
36	Copper(II) complexes of tetradentate thioether-oxime ligands. <i>Inorganica Chimica Acta</i> , 2005, 358, 3449-3456.	1.2	37

#	ARTICLE	IF	CITATIONS
37	Iron ligand recognition by monomeric hemoglobins. BBA - Proteins and Proteomics, 1996, 1295, 209-221. Structure and electrochemical properties of antiferromagnetically coupled binuclear hydroxo-bridged copper(II) complexes with pyridazine and phthalazine ligands. Crystal and molecular structures of	2.1	35
38	[.mu.-3,6-bis(1-pyrazolyl)pyridazine-N2,.mu.-N3,.mu.-N4,N5](.mu.-hydroxo)trichloroaquodicopper-(II)-0.8-water, [.mu.-3,6-bis(1-pyrazolyl)pyridazine-N2,.mu.-N3,.mu.-N4,N5](.+-.-hydroxo)tribromoquodicopper-(II)-0.6-water, and [.mu.-1,4-bis(1-methyl-2-imidazolyl)phthalazine](.mu.-hydroxo)tribromoquodicopper-(II)-wa.	1.9	34
39	Inorga Synthesis, structure and magnetic properties of Nd ³⁺ and Pr ³⁺ 2D polymers with tetrafluoro-p-phthalate. Dalton Transactions, 2011, 40, 10989.	1.6	32
40	Structure, magnetic and luminescence properties of the lanthanide complexes Ln ₂ (Salphen) ₃ ·H ₂ O (Ln=Pr, Nd, Sm, Eu, Gd, Tb, Dy; H ₂ Salphen=N,N-ε ² -bis(salicylidene)-1,2-phenylenediamine). Inorganica Chimica Acta, 2014, 414, 97-104.	1.2	31
41	Crystal and molecular structure of dipolar spin-coupled dimers of an irregularly pentacoordinate copper(II) complex, [Cu(5-Melin)(DBM)]. Inorganic Chemistry, 1982, 21, 60-63.	1.9	29
42	Medium effects on the redox properties of tris(2,2-ε ² -bipyridyl)ruthenium complexes. Inorganica Chimica Acta, 1993, 204, 141-146.	1.2	29
43	Copper co-ordination chemistry of some quadridentate pyridazine and phthalazine (N ₄) thioether ligands. Binuclear copper(II) complexes exhibiting two-electron reduction at positive potentials. Journal of the Chemical Society Dalton Transactions, 1986, , 2381.	1.1	28
44	Efficient mechanochemical synthesis of tris(pyrazolylborate) complexes of manganese(II), cobalt(II) and nickel(II). Inorganic Chemistry Communication, 2004, 7, 485-488.	1.8	28
45	Some high-potential trithioether chelates of copper. Inorganica Chimica Acta, 1992, 196, 97-103.	1.2	26
46	Copper(II) complexes of new unsymmetrical NSN thioether ligands. Inorganica Chimica Acta, 2001, 324, 123-130.	1.2	26
47	Ruthenium(II) complexes with phenanthroline-, benzimidazole-, benzothiazole-, and pyridine-derived bidentate and tridentate ligands: reactivity and spectroscopic and electrochemical characterization. Transition Metal Chemistry, 1993, 18, 197-204.	0.7	25
48	Heme proteins and metalloporphyrins: Redox chemistry and oxygen binding. International Journal of Quantum Chemistry, 1979, 16, 311-329.	1.0	23
49	Some nickel(II) complexes with pentadentate and tridentate heterocyclic N- and S-donor ligands. Inorganica Chimica Acta, 2000, 308, 22-30.	1.2	23
50	A new class of macrocyclic complexes formed via nickel-promoted macrocyclisation of dioxime with dinitrile. Chemical Communications, 2002, , 468-469.	2.2	23
51	Dinuclear Copper(II) Complexes with Bis-thiocarbohydrazone Ligands. European Journal of Inorganic Chemistry, 2008, 2008, 2530-2536. Structural trends in a series of isostructural lanthanide-ε ² -copper metallacrown sulfates (Ln ^{III} = Pr, Nd, Sm, Eu, Gd, Dy and Ho):	1.0	23
52	hexaaquapentakis[1/4₃-glycinehydroxamato(2-)]sulfatopentacopper(II)lanthanide(III) heptaaquapentakis[1/4₃-glycinehydroxamato(2-)]sulfatopentacopper(II)lanthanide(III) sulfate hexahydrate. Acta Crystallographica Section C: Crystal Structure Communications, 2011, 67, m255-m265.	0.4	23
53	Triethylammonium benzene-1,3,5-tricarboxylato(pyridine)zinc(II): a two-dimensional undulating mesh network. Inorganic Chemistry Communication, 2003, 6, 402-404.	1.8	22
54	The 1, 8-bis(2-ε ² -pyridyl)-3, 6-dithiaoctane complex of nickel(II): X-ray crystal structure and borohydride adduct formation. Inorganica Chimica Acta, 1998, 278, 217-222.	1.2	20

#	ARTICLE	IF	CITATIONS
55	A nickel(II) dithia dioxime derived macrocycle. <i>Inorganic Chemistry</i> , 1994, 33, 397-399.	1.9	19
56	Crystal structures and intense luminescence of tris(3-(2-pyridyl)-pyrazolyl)borate Tb ³⁺ and Eu ³⁺ complexes with carboxylate co-ligands. <i>Dalton Transactions</i> , 2017, 46, 3457-3469.	1.6	19
57	High Nuclearity Assemblies and One-Dimensional (1D) Coordination Polymers Based on Lanthanide-Copper 15-Metallacrown-5 Complexes (Ln ³⁺ = Pr, Nd, Sm, Eu). <i>Inorganic Chemistry</i> , 2017, 56, 13152-13165.	1.9	19
58	Oxo-bridged complexes of iron(III) derived from 2-(2-hydroxyphenyl)-benzothiazole and 2-(2-hydroxyphenyl)benzimidazole ligands. <i>Inorganica Chimica Acta</i> , 1989, 166, 59-69.	1.2	18
59	A novel paradigm for metal-induced ring flipping in the copper(II) complex of 1,2-bis(N) Tj ETQq1 1 0.784314 rgBT/Overlock_10 Tf 505	1.8	18
60	Combination of single-molecule magnet behaviour and luminescence properties in a new series of lanthanide complexes with tris(pyrazolyl)borate and oligo(1 ² -diketonate) ligands. <i>Dalton Transactions</i> , 2020, 49, 7774-7789.	1.6	17
61	The Hexakis(thiocyanato)ferrate(III) Ion: a Coordination Chemistry Classic Reveals an Interesting Geometry Pattern for the Thiocyanate Ligands. <i>European Journal of Inorganic Chemistry</i> , 2005, 2005, 2404-2408.	1.0	16
62	Ni(II), Co(II) and Mn(II) tris(pyrazolyl)borate complexes with 2,6-di-tert-butyl-4-carboxy-phenol: Formation of coordinated phenoxyl radical. <i>Inorganic Chemistry Communication</i> , 2005, 8, 932-935.	1.8	14
63	Autoxidation of Reduced Horse Heart Cytochrome <i>c</i> Catalyzed by Cardiolipin-Containing Membranes. <i>Journal of Physical Chemistry B</i> , 2016, 120, 12219-12231.	1.2	14
64	The metamorphosis of heterometallic trinuclear antiferromagnetic complexes into nano-sized superparamagnetic spinels. <i>Materials Chemistry and Physics</i> , 2010, 121, 47-52.	2.0	13
65	Supramolecular Maleate Adducts of Copper(II) 12-Metallacrown-4: Magnetism, EPR, and Alcohol Sorption Properties. <i>European Journal of Inorganic Chemistry</i> , 2017, 2017, 4866-4878.	1.0	13
66	Magnetic Properties of Ln ³⁺ -Cu ^{II} 15-Metallacrown-5 Dimers with Terephthalate (Ln ³⁺ = Pr, Nd, Sm, Eu). <i>European Journal of Inorganic Chemistry</i> , 2018, 2018, 3504-3511.	1.0	13
67	Synthesis of some benzimidazole-, pyridine- and imidazole-derived chelating agents. <i>Journal of Heterocyclic Chemistry</i> , 1989, 26, 541-543.	1.4	12
68	A novel copper(II) complex of a tripodal ligand with phenolate-phenol interligand, intramolecular hydrogen bonding. <i>Dalton Transactions</i> , 2009, , 8111.	1.6	11
69	New Pyrazole- and Benzimidazole-derived Ligand Systems. <i>Journal of Heterocyclic Chemistry</i> , 2018, 55, 1291-1307.	1.4	11
70	Synthesis, structure and magnetic properties of oligometallic systems derived from di- and trinuclear copper(ii) amido-oximate complexes. <i>Dalton Transactions</i> , 2008, , 3007.	1.6	10
71	A tetranuclear copper(II) complex with bis(o-aminobenzaldehyde)thiocarbohydrazone. <i>Inorganica Chimica Acta</i> , 2010, 363, 2065-2070.	1.2	10
72	The role of the bridging group in exchange coupling in dinuclear homo- and heterometallic Ni(ii) and Co(ii) complexes with oxalate, oxamidate and dithiooxamidate bridges. <i>Dalton Transactions</i> , 2012, 41, 11319.	1.6	10

#	ARTICLE	IF	CITATIONS
73	Chemical liquid deposition process for microstructure fabrication. <i>Materials & Design</i> , 2005, 26, 670-679.	5.1	9
74	Effect of Zero-Field Splitting on the Magnetic Susceptibility of Binuclear Complexes of Iron(III). <i>Theoretical and Experimental Chemistry</i> , 2005, 41, 229-234.	0.2	9
75	Synthesis, spectroscopic and redox behaviour of copper(II) complexes with quinquedentate thiaaza ligands. <i>Inorganica Chimica Acta</i> , 1993, 203, 29-35.	1.2	8
76	Some iron(III) complexes with polydentate Schiff base ligands. <i>Inorganica Chimica Acta</i> , 1988, 147, 61-64.	1.2	7
77	Magnetic Characteristics of Trinuclear Complexes $[M_3O(CH_3COO)_6(pz)_3]^+$ (M = Fe, Cr; pz = Pyrazine). <i>Theoretical and Experimental Chemistry</i> , 2004, 40, 214-219.	0.2	7
78	Sorption discrimination between secondary alcohol enantiomers by chiral alkyl-dicarboxylate MOFs. <i>RSC Advances</i> , 2016, 6, 93707-93714.	1.7	7
79	The 1,8-Bis(2- π -pyridyl)-3,6-dithiooctane Complex of Rhodium(III). <i>Journal of Chemical Crystallography</i> , 2012, 42, 295-298.	0.5	6
80	Thermochromism of heme adducts of Glycera hemoglobin and some other monomeric heme proteins. <i>Journal of Inorganic Biochemistry</i> , 1990, 39, 351-369.	1.5	5
81	Synthesis of some benzimidazole-, benzothiazole- and pyridine-derived chelating agents. <i>Journal of Heterocyclic Chemistry</i> , 2002, 39, 399-404.	1.4	5
82	Ru(II) thioether complexes with dangling pyridine ligands. <i>Polyhedron</i> , 2014, 68, 70-75.	1.0	5
83	Absorption- and Excitation-Modulated Luminescence of Pr^{3+} , Nd^{3+} , and Lu^{3+} Compounds with Dianions of Tetrafluoroterephthalic and Camphoric Acids. <i>ACS Omega</i> , 2019, 4, 2669-2675.	1.6	5
84	Heme rotational isomerism is not required for the production of Q-band splitting in the spectra of iron-porphyrin proteins. <i>Inorganic Chemistry</i> , 1991, 30, 1151-1153.	1.9	4
85	The crystal structures of $\{LnCu_5\}^{3+}$ ($Ln = Gd, Dy$ and Ho) 15-metallacrown-5 complexes and a reevaluation of the isotypic Eu^{III} analogue. <i>Acta Crystallographica Section E: Crystallographic Communications</i> , 2019, 75, 1215-1223.	0.2	3
86	The coelomic hemerythrin of siphonosome <i>cumanense</i> . <i>Comparative Biochemistry and Physiology Part B: Comparative Biochemistry</i> , 1982, 72, 433-438.	0.2	2
87	Spectroscopic and kinetic aspects of <i>Elephas maximus</i> hemoglobin. <i>FEBS Journal</i> , 1990, 189, 185-191.	0.2	2
88	Electron-Counting Rules for Transition Metal-Nitric Oxide Complexes. <i>Journal of Chemical Education</i> , 1997, 74, 1354.	1.1	2
89	Triangular kinetic schemes applied to the stability of a Heme-Globin complex. <i>Journal of Inorganic Biochemistry</i> , 1997, 66, 83-98.	1.5	2
90	Copper(II) Schiff-base complexes and apoglobin stability. <i>Journal of Inorganic Biochemistry</i> , 1999, 73, 137-144.	1.5	2

#	ARTICLE	IF	CITATIONS
91	5,7,12,14-Tetrahydro-5,14:7,12-bis([1,2]benzeno)pentacene-6,13-diol dimethylformamide disolvate. IUCrData, 2016, 1, .	0.1	2
92	Chiroptical and physicochemical properties of the extracellular haemoglobin from <i>Cirriformia tentaculata</i> . <i>Comparative Biochemistry and Physiology Part B: Comparative Biochemistry</i> , 1990, 97, 391-399.	0.2	1
93	Ruthenium(II) complexes of some simple classic amine ligands. <i>Inorganica Chimica Acta</i> , 2015, 432, 185-191.	1.2	1
94	New homodinuclear tris(3-alkylpyrazolyl)borate complexes of Colland Nillwith a tetraacetylene dianion as a bridging ligand. <i>Acta Crystallographica Section C, Structural Chemistry</i> , 2016, 72, 777-785.	0.2	1
95	Crystal structure of 5,7,12,14-tetrahydro-5,14:7,12-bis([1,2]benzeno)pentacene-6,13-dione. <i>Acta Crystallographica Section E: Crystallographic Communications</i> , 2016, 72, 1734-1738.	0.2	1
96	Lanthanide Complexes with 4,4'-bis(2-sulfonatostyryl)-biphenyl: Crystal Structures and Luminescence Properties. <i>European Journal of Inorganic Chemistry</i> , 2022, 2022, .	1.0	1
97	Crystal structure of (2,2'-bipyridyl)[2,6-bis(1-butyl-1H-benzimidazol-2-yl)pyridine]chloridoiridium(III) trifluoromethanesulfonate. <i>Acta Crystallographica Section E: Crystallographic Communications</i> , 2017, 73, 127-132.	0.2	0
98	Crystal structure of bis(pivaloylhydroxamato) ²⁻ copper(II). <i>Acta Crystallographica Section E: Crystallographic Communications</i> , 2018, 74, 1384-1387.	0.2	0
99	Ruthenium(II) complexes of the tetradentate polypyridyl thioether 1,2-bis[3-(2-pyridyl)-1-thiopropyl]benzene. <i>Polyhedron</i> , 2020, 179, 114367.	1.0	0
100	Crystal structure of a Tb ^{III} –Cu ^{II} glycinehydroxamate 15-metallacrown-5 sulfate complex. <i>Acta Crystallographica Section E: Crystallographic Communications</i> , 2021, 77, 1197-1202.	0.2	0
101	1,3-Bis(N-Methylbenzimidazol-2-yl)-2-Phenylpropanedichloridocopper(II). <i>Journal of Chemical Crystallography</i> , 0, , 1.	0.5	0
102	Two Bifunctionalised Derivatives of 1,2,4-Triazole. <i>Journal of Chemical Crystallography</i> , 0, , 1.	0.5	0
103	Bis[3-(anthracen-9-yl)pentane-2,4-dionato] ²⁻ (N,N'-dimethylformamide) ¹⁻ [tris(pyrazol-1-yl)N] Tj ETQq1 1 0.784314 rgBT /Over Communications, 2022, 78, 103-107.	0.2	0
104	Chlorocobalt complexes with pyridylethyl-derived diazacycloalkanes. <i>Acta Crystallographica Section E: Crystallographic Communications</i> , 2022, 78, 235-243.	0.2	0