

# Jir Klimes

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/2733842/jiri-klimes-publications-by-year.pdf>

**Version:** 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

71  
papers

730  
citations

15  
h-index

20  
g-index

73  
ext. papers

776  
ext. citations

2.9  
avg, IF

3.39  
L-index

#	Paper	IF	Citations
71	Indacaterol/Glycopyrronium versus Salmeterol/Fluticasone in Patients with COPD-A Cost-Effectiveness Analysis in the Czech Republic. <i>Value in Health Regional Issues</i> , <b>2018</b> , 16, 112-118	1.6	1
70	Cost-of-illness analysis and regression modeling in cystic fibrosis: a retrospective prevalence-based study. <i>European Journal of Health Economics</i> , <b>2017</b> , 18, 73-82	3.6	7
69	A model of natural degradation of 17- $\beta$ -ethinylestradiol in surface water and identification of degradation products by GC-MS. <i>Environmental Science and Pollution Research</i> , <b>2017</b> , 24, 23196-23206	5.1	2
68	Solid-Phase Extraction of Ibuprofen from Pharmaceuticals via Ligand Exchange Using Zirconium Dioxide. <i>Current Analytical Chemistry</i> , <b>2016</b> , 12, 523-528	1.7	3
67	Placental passage of olomoucine II, but not purvalanol A, is affected by p-glycoprotein (ABCB1), breast cancer resistance protein (ABCG2) and multidrug resistance-associated proteins (ABCCs). <i>Xenobiotica</i> , <b>2016</b> , 46, 416-23	2	1
66	Dinaciclib, a cyclin-dependent kinase inhibitor, is a substrate of human ABCB1 and ABCG2 and an inhibitor of human ABCC1 in vitro. <i>Biochemical Pharmacology</i> , <b>2015</b> , 98, 465-72	6	23
65	LC-UV/MS methods for the analysis of prochelator-boronyl salicylaldehyde isonicotinoyl hydrazone (BSIH) and its active chelator salicylaldehyde isonicotinoyl hydrazone (SIH). <i>Journal of Pharmaceutical and Biomedical Analysis</i> , <b>2015</b> , 105, 55-63	3.5	9
64	Simultaneous determination of the novel thiosemicarbazone anti-cancer agent, Bp4eT, and its main phase I metabolites in plasma: application to a pilot pharmacokinetic study in rats. <i>Biomedical Chromatography</i> , <b>2014</b> , 28, 621-9	1.7	7
63	Medical and Productivity Costs of Rheumatoid Arthritis in The Czech Republic: Cost-of-Illness Study Based on Disease Severity. <i>Value in Health Regional Issues</i> , <b>2014</b> , 4, 75-81	1.6	12
62	Work Productivity and Costs Related to Patients with Ankylosing Spondylitis, Rheumatoid Arthritis, and Psoriasis. <i>Value in Health Regional Issues</i> , <b>2014</b> , 4, 100-106	1.6	10
61	Zirconia--a stationary phase capable of the separation of polar markers of myocardial metabolism in hydrophilic interaction chromatography. <i>Journal of Separation Science</i> , <b>2014</b> , 37, 1089-93	3.4	4
60	Identification of in vitro metabolites of the novel anti-tumor thiosemicarbazone, DpC, using ultra-high performance liquid chromatography-quadrupole-time-of-flight mass spectrometry. <i>Analytical and Bioanalytical Chemistry</i> , <b>2013</b> , 405, 1651-61	4.4	5
59	Development of LC-MS/MS method for the simultaneous analysis of the cardioprotective drug dexrazoxane and its metabolite ADR-925 in isolated cardiomyocytes and cell culture medium. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , <b>2013</b> , 76, 243-51	3.5	9
58	Olomoucine II, but not purvalanol A, is transported by breast cancer resistance protein (ABCG2) and P-glycoprotein (ABCB1). <i>PLoS ONE</i> , <b>2013</b> , 8, e75520	3.7	6
57	Identification of Pavinane Alkaloids in the Genera Argemone and Eschscholzia by GC-MS. <i>Natural Product Communications</i> , <b>2012</b> , 7, 1934578X1200701	0.9	1
56	Alkaloids from Some Amaryllidaceae Species and Their Cholinesterase Activity. <i>Natural Product Communications</i> , <b>2012</b> , 7, 1934578X1200700	0.9	5
55	LC-MS/MS identification of the principal in vitro and in vivo phase I metabolites of the novel thiosemicarbazone anti-cancer drug, Bp4eT. <i>Analytical and Bioanalytical Chemistry</i> , <b>2012</b> , 403, 309-21	4.4	13

54	The influence of a carbon layer deposited on a zirconia surface on the retention of polar analytes in an organic rich mobile phase. <i>Journal of Chromatography A</i> , <b>2012</b> , 1232, 242-7	4.5	6
53	The Retention Behavior of Acidic, Basic and Neutral Pharmaceuticals on the Deactivated Polybutadiene Zirconia Phase. <i>Current Analytical Chemistry</i> , <b>2012</b> , 8, 574-582	1.7	1
52	The retention behaviour of polar compounds on zirconia based stationary phases under hydrophilic interaction liquid chromatography conditions. <i>Journal of Chromatography A</i> , <b>2011</b> , 1218, 6981-6	4.5	23
51	Use of different stationary phases for separation of isoniazid, its metabolites and vitamin B6 forms. <i>Journal of Separation Science</i> , <b>2011</b> , 34, 1357-65	3.4	12
50	Hydrophilic interaction liquid chromatography in the separation of a moderately lipophilic drug from its highly polar metabolites--the cardioprotectant dexrazoxane as a model case. <i>Journal of Chromatography A</i> , <b>2011</b> , 1218, 416-26	4.5	16
49	HPLC analysis of a syrup containing nimesulide and its hydrolytic degradation product. <i>Chemical Papers</i> , <b>2010</b> , 64,	1.9	2
48	Development of an LC-MS/MS method for analysis of interconvertible Z/E isomers of the novel anticancer agent, Bp4eT. <i>Analytical and Bioanalytical Chemistry</i> , <b>2010</b> , 397, 161-171	4.4	8
47	An innovative approach to the analysis of 3-[4-(2-methylpropyl)phenyl]propanoic acid as an impurity of ibuprofen on a carbon-coated zirconia stationary phase. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , <b>2009</b> , 49, 1150-6	3.5	6
46	Development and Validation of an LCESI-MS Ion-Trap Method for Analysis of Impurities in Transkarbam 12, a Novel Transdermal Accelerant. <i>Chromatographia</i> , <b>2009</b> , 69, 977-983	2.1	1
45	RP-ZrO <sub>2</sub> Stationary Phase as an Alternative to Separate of Doxazosin Impurities. <i>Chromatographia</i> , <b>2009</b> , 70, 185-189	2.1	2
44	HPLC methods for determination of two novel thiosemicarbazone anti-cancer drugs (N4mT and Dp44mT) in plasma and their application to in vitro plasma stability of these agents. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , <b>2009</b> , 877, 316-22	3.2	18
43	Isolation and identification of amphetamines in urine by thin-layer chromatography. <i>Journal of Planar Chromatography - Modern TLC</i> , <b>2008</b> , 21, 465-468	0.9	0
42	HPLC-DAD and MS/MS analysis of novel drug candidates from the group of aromatic hydrazones revealing the presence of geometric isomers. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , <b>2008</b> , 48, 295-302	3.5	21
41	Determination of lipophilicity of novel potential antituberculosic agents using HPLC on monolithic stationary phase and theoretical calculations. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , <b>2008</b> , 48, 310-4	3.5	14
40	Investigation of the stability of aromatic hydrazones in plasma and related biological material. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , <b>2008</b> , 47, 360-70	3.5	30
39	Titania-based stationary phase in separation of ondansetron and its related compounds. <i>Journal of Chromatography A</i> , <b>2008</b> , 1189, 83-91	4.5	15
38	Development and validation of HPLC-DAD methods for the analysis of two novel iron chelators with potent anti-cancer activity. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , <b>2007</b> , 43, 1343-51	3.5	11
37	Potential employment of non-silica-based stationary phases in pharmaceutical analysis. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , <b>2007</b> , 44, 1048-55	3.5	11

36	High-performance liquid chromatographic method with UV photodiode-array, fluorescence and mass spectrometric detection for simultaneous determination of galantamine and its phase I metabolites in biological samples. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , <b>2007</b> , 853, 265-74	3.2	30
35	Using of HPLC coupled with coulometric detector for the determination of biotin in pharmaceuticals. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , <b>2007</b> , 45, 730-5	3.5	11
34	Preliminary pharmaceutico-analytical evaluation of transkarbam 12 using liquid chromatography. <i>Journal of Separation Science</i> , <b>2006</b> , 29, 1595-9	3.4	3
33	Comparison of different stationary phases for bioanalytical studies of biologically active compounds. <i>Journal of Separation Science</i> , <b>2006</b> , 29, 2126-35	3.4	3
32	HPLC study on stability of pyridoxal isonicotinoyl hydrazone. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , <b>2006</b> , 40, 105-12	3.5	17
31	Optimization of HPLC chromatographic conditions for determination of Transkarbam 12 and its degradation products. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , <b>2006</b> , 42, 136-42	3.5	5
30	HPLC determination of a novel aroylhydrazone iron chelator (o-108) in rabbit plasma and its application to a pilot pharmacokinetic study. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , <b>2006</b> , 838, 107-12	3.2	14
29	Determination of the lipophilicity of potential antituberculosic compounds by RP-TLC. <i>Journal of Planar Chromatography - Modern TLC</i> , <b>2006</b> , 19, 422-426	0.9	10
28	Disposition study of a new potential antineoplastic agent dimefluron in rats using high-performance liquid chromatography with ultraviolet and mass spectrometric detection. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , <b>2005</b> , 37, 1059-71	3.5	9
27	Use of chiral liquid chromatography for the evaluation of stereospecificity in the carbonyl reduction of potential benzo[c]fluorene antineoplastics benfluron and dimefluron in various species. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , <b>2005</b> , 37, 1049-57	3.5	6
26	Stability of ramipril in the solvents of different pH. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , <b>2005</b> , 37, 1179-83	3.5	15
25	Use of the zirconia-based stationary phase for separation of ibuprofen and its impurities. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , <b>2005</b> , 38, 609-18	3.5	13
24	Utilization of zirconia stationary phase as a tool in drug control. <i>Journal of Separation Science</i> , <b>2005</b> , 28, 1307-14	3.4	9
23	Development of high-performance liquid chromatographic determination of salicylaldehyde isonicotinoyl hydrazone in rabbit plasma and application of this method to an in vivo study. <i>Journal of Separation Science</i> , <b>2005</b> , 28, 1300-6	3.4	30
22	Reversed-phase thin-layer chromatographic determination of the lipophilicity of potential antituberculosic compounds. <i>Journal of Planar Chromatography - Modern TLC</i> , <b>2005</b> , 18, 450-454	0.9	7
21	Chromatographic methods for the separation of biocompatible iron chelators from their synthetic precursors and iron chelates. <i>Journal of Separation Science</i> , <b>2004</b> , 27, 1503-10	3.4	11
20	HPLC study of glimepiride under hydrolytic stress conditions. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , <b>2004</b> , 36, 205-9	3.5	32
19	Reversed-Phase High Performance Liquid Chromatographic Determination of Lipophilicity of Potential Antituberculosis Compounds. <i>Journal of Liquid Chromatography and Related Technologies</i> , <b>2004</b> , 27, 2539-2545	1.3	5

18	Photochemical stability of nimesulide. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , <b>2003</b> , 31, 827-325	3.5	15
17	A study of the conditions of the supercritical fluid extraction in the analysis of selected anti-inflammatory drugs in plasma. <i>Il Farmaco</i> , <b>2002</b> , 57, 117-22		7
16	LIPOPHILICITY CHARACTERIZATION BY REVERSED-PHASE HPLC OF POTENTIAL ANTITUBERCULOTICS. <i>Journal of Liquid Chromatography and Related Technologies</i> , <b>2002</b> , 25, 2849-2856	1.3	4
15	Study of the lipophilicity of potential antituberculosic compounds by reversed-phase thin-layer chromatography. <i>Journal of Planar Chromatography - Modern TLC</i> , <b>2002</b> , 15, 200-203	0.9	10
14	HPLC evaluation of diclofenac in transdermal therapeutic preparations. <i>International Journal of Pharmaceutics</i> , <b>2001</b> , 217, 153-60	6.5	19
13	DETERMINATION OF LIPOPHILICITY OF POTENTIAL ANTITUBERCULOUS DRUGS BY REVERSED-PHASE HIGH PERFORMANCE LIQUID CHROMATOGRAPHY. <i>Journal of Liquid Chromatography and Related Technologies</i> , <b>2001</b> , 24, 2257-2265	1.3	4
12	Reversed-phase thin-layer chromatographic determination of the lipophilicity of potential antituberculosic compounds. <i>Journal of Planar Chromatography - Modern TLC</i> , <b>2001</b> , 14, 291-295	0.9	7
11	Combination of molecular modeling and quantitative structure-activity relationship analysis in the study of antimycobacterial activity of pyridine derivatives. <i>International Journal of Pharmaceutics</i> , <b>2000</b> , 207, 1-6	6.5	16
10	HPLC ANALYSIS OF TIAPROFENIC ACID IN THE SAMPLES OF WHOLE BLOOD USING L-L AND S-L EXTRACTIONS. <i>Journal of Liquid Chromatography and Related Technologies</i> , <b>2000</b> , 23, 3191-3201	1.3	1
9	Microcolumn high-performance liquid chromatographic assay for doxycycline in isolated alveolar macrophages. <i>Journal of Chromatography A</i> , <b>1999</b> , 846, 181-4	4.5	10
8	Relationship between structure and reversed-phase thin-layer chromatographic lipophilicity parameters in a group of piperazine derivatives. <i>Journal of Chromatography A</i> , <b>1997</b> , 766, 165-170	4.5	12
7	High-performance liquid chromatographic determination of terguride in solid dosage forms and plasma. <i>Biomedical Applications</i> , <b>1995</b> , 663, 309-13		3
6	Determination of ibuprofen in erythrocytes and plasma by high performance liquid chromatography. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , <b>1995</b> , 13, 899-903	3.5	33
5	High-Performance Liquid Chromatographic Analysis of Kebuzone and Its Metabolites in the Samples of Erythrocytes, Plasma, and Whole Blood. <i>Journal of Liquid Chromatography and Related Technologies</i> , <b>1995</b> , 18, 2147-2166		
4	High-performance liquid chromatographic assay for ibuprofen in whole blood using solid-phase extraction. <i>Biomedical Applications</i> , <b>1994</b> , 654, 282-6		11
3	Relations between Structure and Antituberculosic Activity of 4-Alkoxybenzoic Acids. <i>Collection of Czechoslovak Chemical Communications</i> , <b>1993</b> , 58, 191-196		6
2	Chromatographic behaviour of dipyrildylsulphides Relationship between log k' values and structure by reversed-phase high-performance liquid chromatography. <i>Journal of Chromatography A</i> , <b>1992</b> , 595, 334-336	4.5	8
1	Simultaneous high-performance liquid chromatographic determination of salicylates in whole blood, plasma and isolated erythrocytes. <i>Biomedical Applications</i> , <b>1992</b> , 584, 221-228		18

