

# Alan Tennant

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2743506/publications.pdf>

Version: 2024-02-01

186  
papers

11,391  
citations

34016

52  
h-index

32761

100  
g-index

190  
all docs

190  
docs citations

190  
times ranked

11931  
citing authors

#	ARTICLE	IF	CITATIONS
1	Scale Banking for <scp>Patient-Reported</scp> Outcome Measures That Measure Functioning in Rheumatoid Arthritis: A Daily Activities Metric. <i>Arthritis Care and Research</i> , 2022, 74, 579-587.	1.5	6
2	Fatigue and anxiety mediate the effect of dyspnea on quality of life in amyotrophic lateral sclerosis. <i>Amyotrophic Lateral Sclerosis and Frontotemporal Degeneration</i> , 2022, 23, 390-398.	1.1	4
3	The four self-efficacy trajectories among people with multiple sclerosis: Clinical associations and implications. <i>Journal of the Neurological Sciences</i> , 2022, 436, 120188.	0.3	3
4	Cross-cultural adaptation and psychometric testing of the Dutch and German versions of the Evaluation of Daily Activity Questionnaire in people with rheumatoid arthritis. <i>Rheumatology International</i> , 2021, 41, 951-964.	1.5	2
5	Measuring coping in people with amyotrophic lateral sclerosis using the Coping Index-ALS: A patient derived, Rasch compliant scale. <i>Journal of the Neurological Sciences</i> , 2021, 421, 117285.	0.3	0
6	Measuring Quality of Life in Carers of People With Dementia: Development and Psychometric Evaluation of Scales measuring the Impact of DEmentia on CARers (SIDE CAR). <i>Gerontologist</i> , The, 2021, 61, e1-e11.	2.3	8
7	Quality of life in multiple sclerosis is dominated by fatigue, disability and self-efficacy. <i>Journal of the Neurological Sciences</i> , 2021, 426, 117437.	0.3	19
8	Use and detailed metric properties of patient-reported outcome measures for rheumatoid arthritis: a systematic review covering two decades. <i>RMD Open</i> , 2021, 7, e001707.	1.8	8
9	Measuring Burnout in Pediatric Oncology Staff: Should We Be Using the Maslach Burnout Inventory?. <i>Journal of Pediatric Oncology Nursing</i> , 2020, 37, 55-64.	1.5	12
10	Validation and assessment of minimally clinically important difference of the unadjusted Health Assessment Questionnaire in a Danish cohort: uncovering ordinal bias. <i>Scandinavian Journal of Rheumatology</i> , 2020, 49, 1-7.	0.6	13
11	Do pain, anxiety and depression influence quality of life for people with amyotrophic lateral sclerosis/motor neuron disease? A national study reconciling previous conflicting literature. <i>Journal of Neurology</i> , 2020, 267, 607-615.	1.8	25
12	Investigating the patient acceptable symptom state cut-offs: longitudinal data from a community cohort using the shoulder pain and disability index. <i>Rheumatology International</i> , 2020, 40, 599-605.	1.5	15
13	The influence and added value of a Standardized Assessment and Reporting System for functioning outcomes upon national rehabilitation quality reports. <i>International Journal for Quality in Health Care</i> , 2020, 32, 379-387.	0.9	4
14	Sexual Morbidity Assessment in Gyne-Oncology Follow-Up: Development of the Sexual Well-Being After Cervical or Endometrial Cancer (SWELL-CE) Patient-Reported Outcome Measure. <i>Journal of Sexual Medicine</i> , 2020, 17, 2005-2015.	0.3	5
15	The Valued Life Activities Scale (VLAs): linguistic validation, cultural adaptation and psychometric testing in people with rheumatic and musculoskeletal diseases in the UK. <i>BMC Musculoskeletal Disorders</i> , 2020, 21, 505.	0.8	2
16	Creating a common metric based on existing activities of daily living tools to enable standardized reporting of functioning outcomes achieved during rehabilitation. <i>Journal of Rehabilitation Medicine</i> , 2020, 52, jrm00085.	0.8	3
17	The WHOQOL-BREF: a modern psychometric evaluation of its internal construct validity in people with multiple sclerosis. <i>Quality of Life Research</i> , 2020, 29, 1961-1972.	1.5	15
18	Measuring quality of life in ALS/MND: validation of the WHOQOL-BREF. <i>Amyotrophic Lateral Sclerosis and Frontotemporal Degeneration</i> , 2020, 21, 364-372.	1.1	5

#	ARTICLE	IF	CITATIONS
19	Measuring Work-Related Functioning Using the Work Rehabilitation Questionnaire (WORQ). International Journal of Environmental Research and Public Health, 2019, 16, 2795.	1.2	13
20	Revisiting the disabilities of the arm, shoulder and hand (DASH) and QuickDASH in rheumatoid arthritis. BMC Musculoskeletal Disorders, 2019, 20, 41.	0.8	23
21	The relationships between symptoms, disability, perceived health and quality of life in amyotrophic lateral sclerosis/motor neuron disease. Amyotrophic Lateral Sclerosis and Frontotemporal Degeneration, 2019, 20, 317-327.	1.1	27
22	Uptake of the OMERACT-OARSI Hip and Knee Osteoarthritis Core Outcome Set: Review of Randomized Controlled Trials from 1997 to 2017. Journal of Rheumatology, 2019, 46, 976-980.	1.0	25
23	The Functional Independence Measure 18-item version can be reported as a unidimensional interval-scaled metric: Internal construct validity revisited. Journal of Rehabilitation Medicine, 2019, 51, 193-200.	0.8	30
24	Supporting construct validity of the Evaluation of Daily Activity Questionnaire using Linear Logistic Test Models. Quality of Life Research, 2019, 28, 1627-1639.	1.5	7
25	The prevalence and impact of self-reported foot and ankle pain in the over 55 age group: a secondary data analysis from a large community sample. Journal of Foot and Ankle Research, 2019, 12, 53.	0.7	13
26	The OMERACT-OARSI Core Domain Set for Measurement in Clinical Trials of Hip and/or Knee Osteoarthritis. Journal of Rheumatology, 2019, 46, 981-989.	1.0	82
27	The measurement of functioning using the International Classification of Functioning, Disability and Health: comparing qualifier ratings with existing health status instruments. Disability and Rehabilitation, 2019, 41, 541-548.	0.9	39
28	Validation of Fatigue Impact Scale with various item sets – a Rasch analysis. Disability and Rehabilitation, 2019, 41, 840-846.	0.9	13
29	Psychological determinants of job retention in multiple sclerosis. Multiple Sclerosis Journal, 2019, 25, 419-426.	1.4	9
30	Development and validation of Spasticity Index-Amyotrophic Lateral Sclerosis. Acta Neurologica Scandinavica, 2018, 138, 47-54.	1.0	7
31	The psychometric properties of the Evaluation of Daily Activity Questionnaire in seven musculoskeletal conditions. Disability and Rehabilitation, 2018, 40, 2070-2080.	0.9	3
32	Quality of life for post-polio syndrome: a patient derived, Rasch standard scale. Disability and Rehabilitation, 2018, 40, 597-602.	0.9	4
33	Effect of Fatigue, Older Age, Higher Body Mass Index, and Female Sex on Disability in Early Rheumatoid Arthritis in the Treatment-Target Era. Arthritis Care and Research, 2018, 70, 361-368.	1.5	12
34	Measuring Physical and Cognitive Fatigue in People With Post-Polio Syndrome: Development of the Neurological Fatigue Index for Post-Polio Syndrome (NFI-PP). PM and R, 2018, 10, 129-136.	0.9	1
35	Risk factors for social withdrawal in amyotrophic lateral sclerosis/motor neurone disease. Amyotrophic Lateral Sclerosis and Frontotemporal Degeneration, 2018, 19, 591-598.	1.1	11
36	Measure of activity performance of the hand (MAP-Hand) questionnaire: linguistic validation, cultural adaptation and psychometric testing in people with rheumatoid arthritis in the UK. BMC Musculoskeletal Disorders, 2018, 19, 275.	0.8	4

#	ARTICLE	IF	CITATIONS
37	Standardized reporting of functioning information on ICF-based common metrics. <i>European Journal of Physical and Rehabilitation Medicine</i> , 2018, 54, 110-117.	1.1	37
38	All metrics are equal, but some metrics are more equal than others: A systematic search and review on the use of the term "metric". <i>PLoS ONE</i> , 2018, 13, e0193861.	1.1	6
39	The Impact of Missing Values and Single Imputation upon Rasch Analysis Outcomes: A Simulation Study. <i>Journal of Applied Measurement</i> , 2018, 19, 1-25.	0.3	4
40	Adaptation of the osteoarthritis-specific quality of life scale (the OAQoL) for use in Germany, Hungary, Italy, Spain and Turkey. <i>Rheumatology International</i> , 2017, 37, 727-734.	1.5	6
41	Sexual functioning in multiple sclerosis: Relationships with depression, fatigue and physical function. <i>Multiple Sclerosis Journal</i> , 2017, 23, 1268-1275.	1.4	24
42	A study of the internal construct and predictive validity of the SMART assessment for emergence from vegetative state. <i>Brain Injury</i> , 2017, 31, 185-192.	0.6	3
43	Integrating patient reported outcome measures and computerized adaptive test estimates on the same common metrics: an example from the assessment of activities in rheumatoid arthritis. <i>International Journal of Rheumatic Diseases</i> , 2017, 20, 1413-1425.	0.9	3
44	Patient-reported Outcomes as Predictors of Change in Disease Activity and Disability in Early Rheumatoid Arthritis: Results from the Yorkshire Early Arthritis Register. <i>Journal of Rheumatology</i> , 2017, 44, 1331-1340.	1.0	6
45	Rasch analysis of SF-36 Health Survey in multiple sclerosis. <i>Neurology and Urodynamics</i> , 2017, 36, 1161-1166.	0.8	14
46	Metric properties of the Spinal Cord Independence Measure - Self report in a community survey. <i>Journal of Rehabilitation Medicine</i> , 2016, 48, 149-164.	0.8	31
47	Multiple sclerosis and employment: Associations of psychological factors and work instability. <i>Journal of Rehabilitation Medicine</i> , 2016, 48, 799-805.	0.8	18
48	Further work on validating the Huntington's disease quality of life questionnaire (HDQoL). <i>Journal of Neurology, Neurosurgery and Psychiatry</i> , 2016, 87, A65.2-A65.	0.9	0
49	Toward a Standardized Reporting of Outcomes in Hand Osteoarthritis: Developing a Common Metric of Outcome Measures Commonly Used to Assess Functioning. <i>Arthritis Care and Research</i> , 2016, 68, 1115-1127.	1.5	9
50	Measurement properties of multidimensional patient-reported outcome measures in neurodisability: a systematic review of evaluation studies. <i>Developmental Medicine and Child Neurology</i> , 2016, 58, 437-451.	1.1	30
51	The Manual Work Instability Scale: development and validation. <i>Occupational Medicine</i> , 2016, 66, 300-304.	0.8	6
52	Harmonizing routinely collected health information for strengthening quality management in health systems: requirements and practice. <i>Journal of Health Services Research and Policy</i> , 2016, 21, 223-228.	0.8	12
53	Minimal Clinically Important Difference as Applied in Rheumatology: An OMERACT Rasch Working Group Systematic Review and Critique. <i>Journal of Rheumatology</i> , 2016, 43, 194-202.	1.0	56
54	Spasticity in multiple sclerosis: Associations with impairments and overall quality of life. <i>Multiple Sclerosis and Related Disorders</i> , 2016, 5, 34-39.	0.9	46

#	ARTICLE	IF	CITATIONS
55	Toward a Clinical Definition of Early Osteoarthritis: Onset of Patient-Reported Knee Pain Begins on Stairs. Data From the Osteoarthritis Initiative. <i>Arthritis Care and Research</i> , 2015, 67, 40-47.	1.5	77
56	PSYCHOLOGICAL ATTRIBUTES AND WORK INSTABILITY IN MS. <i>Journal of Neurology, Neurosurgery and Psychiatry</i> , 2015, 86, e4.29-e4.	0.9	0
57	The reliability and validity of the English version of the Evaluation of Daily Activity Questionnaire for people with rheumatoid arthritis. <i>Rheumatology</i> , 2015, 54, 1605-1615.	0.9	17
58	Validity and reliability of the Persian version of Behrseta's disease quality-of-life (BD-QoL) questionnaire: a cross-cultural adaptation. <i>Rheumatology International</i> , 2015, 35, 677-684.	1.5	4
59	A Systematic Review of Generic Multidimensional Patient-Reported Outcome Measures for Children, Part I: Descriptive Characteristics. <i>Value in Health</i> , 2015, 18, 315-333.	0.1	56
60	A Systematic Review of Generic Multidimensional Patient-Reported Outcome Measures for Children, Part II: Evaluation of Psychometric Performance of English-Language Versions in a General Population. <i>Value in Health</i> , 2015, 18, 334-345.	0.1	37
61	Meaningful health outcomes for paediatric neurodisability: Stakeholder prioritisation and appropriateness of patient reported outcome measures. <i>Health and Quality of Life Outcomes</i> , 2015, 13, 87.	1.0	53
62	The content validity and acceptability of the Evaluation of Daily Activity Questionnaire in musculoskeletal conditions. <i>British Journal of Occupational Therapy</i> , 2015, 78, 144-157.	0.5	3
63	Adaptation and validation of the Ankylosing Spondylitis Work Instability Scale (AS-WIS) for use in Turkey. <i>Rheumatology International</i> , 2015, 35, 125-131.	1.5	1
64	Translation into British Sign Language and validation of the Strengths and Difficulties Questionnaire. <i>Health Services and Delivery Research</i> , 2015, 3, 1-96.	1.4	10
65	OPO083-HPR...Development of the Evaluation of Daily Activities Questionnaire in Musculoskeletal Conditions: Psychometric Testing. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 93.1-93.	0.5	0
66	Is the Pain Visual Analogue Scale Linear and Responsive to Change? An Exploration Using Rasch Analysis. <i>PLoS ONE</i> , 2014, 9, e99485.	1.1	152
67	Linguistic validation and cultural adaptation of an English version of the Evaluation of Daily Activity Questionnaire in rheumatoid arthritis. <i>Health and Quality of Life Outcomes</i> , 2014, 12, 143.	1.0	10
68	Self-perceived health status following aneurysmal subarachnoid haemorrhage: a cohort study. <i>BMJ Open</i> , 2014, 4, e003932.	0.8	25
69	Validation of the educational needs assessment tool as a generic instrument for rheumatic diseases in seven European countries. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 2122-2129.	0.5	37
70	Staff burnout in paediatric oncology: new tools to facilitate the development and evaluation of effective interventions. <i>European Journal of Cancer Care</i> , 2014, 23, 450-461.	0.7	20
71	A Systematic Literature Review on the Application of Rasch Analysis in Musculoskeletal Disease - A Special Interest Group Report of OMERACT 11. <i>Journal of Rheumatology</i> , 2014, 41, 159-164.	1.0	61
72	Construct Validity of the Swedish Version of the Revised Piper Fatigue Scale in an Oncology Sample - A Rasch Analysis. <i>Value in Health</i> , 2014, 17, 360-363.	0.1	8

#	ARTICLE	IF	CITATIONS
73	Development of item bank to measure deliberate self-harm behaviours: Facilitating tailored scales and computer adaptive testing for specific research and clinical purposes. <i>Psychiatry Research</i> , 2014, 217, 240-247.	1.7	10
74	Informing the NHS Outcomes Framework: evaluating meaningful health outcomes for children with neurodisability using multiple methods including systematic review, qualitative research, Delphi survey and consensus meeting. <i>Health Services and Delivery Research</i> , 2014, 2, 1-224.	1.4	40
75	Assessing the risk of self-harm in an adult offender population: an incidence cohort study. <i>Health Technology Assessment</i> , 2014, 18, 1-152.	1.3	10
76	Development and evaluation of tools and an intervention to improve patient- and carer-centred outcomes in Longer-Term Stroke care and exploration of adjustment post stroke: the LoTS care research programme. <i>Programme Grants for Applied Research</i> , 2014, 2, 1-224.	0.4	11
77	Construct validity of the psychological general well being index (PGWBI) in a sample of patients undergoing treatment for stress-related exhaustion: a rasch analysis. <i>Health and Quality of Life Outcomes</i> , 2013, 11, 2.	1.0	95
78	Measuring engagement in deliberate self-harm behaviours: psychometric evaluation of six scales. <i>BMC Psychiatry</i> , 2013, 13, 4.	1.1	42
79	Epidemiology of neurologically disabling disorders. <i>Handbook of Clinical Neurology / Edited By P J Vinken and G W Bruyn</i> , 2013, 110, 77-92.	1.0	8
80	Assessing social isolation in motor neurone disease: A Rasch analysis of the MND Social Withdrawal Scale. <i>Journal of the Neurological Sciences</i> , 2013, 334, 112-118.	0.3	8
81	Reliability, Construct Validity and Measurement Potential of the International Classification of Functioning, Disability and Health Comprehensive Core Set for Chronic Widespread Pain. <i>Journal of Musculoskeletal Pain</i> , 2013, 21, 231-243.	0.3	1
82	An introduction to Rasch analysis for Psychiatric practice and research. <i>Journal of Psychiatric Research</i> , 2013, 47, 141-148.	1.5	78
83	Fatigue and daytime sleepiness scale in myotonic dystrophy type 1. <i>Muscle and Nerve</i> , 2013, 47, 89-95.	1.0	52
84	Adaptive screening for depression – Recalibration of an item bank for the assessment of depression in persons with mental and somatic diseases and evaluation in a simulated computer-adaptive test environment. <i>Journal of Psychosomatic Research</i> , 2013, 75, 437-443.	1.2	23
85	Can We Scientifically and Reliably Measure the Level of Consciousness in Vegetative and Minimally Conscious States? Rasch Analysis of the Coma Recovery Scale-Revised. <i>Archives of Physical Medicine and Rehabilitation</i> , 2013, 94, 527-535.e1.	0.5	55
86	Development of a Self-Report Scale of Spasticity. <i>Topics in Stroke Rehabilitation</i> , 2013, 20, 485-492.	1.0	7
87	Validation of the Longer-term Unmet Needs after Stroke (LUNS) monitoring tool: a multicentre study. <i>Clinical Rehabilitation</i> , 2013, 27, 1020-1028.	1.0	37
88	Perceived changes and minimum clinically important difference of the Neurological Fatigue Index for multiple sclerosis (NFI-MS). <i>Multiple Sclerosis Journal</i> , 2013, 19, 502-505.	1.4	19
89	The reliability and validity of the World Health Organization Disability Assessment Schedule (WHODAS-II) in stroke. <i>Disability and Rehabilitation</i> , 2013, 35, 214-220.	0.9	55
90	The impact of fatigue and psychosocial variables on quality of life for patients with motor neuron disease. <i>Amyotrophic Lateral Sclerosis and Frontotemporal Degeneration</i> , 2013, 14, 537-545.	1.1	31

#	ARTICLE	IF	CITATIONS
91	FRI0485-HPRâ€¦Development of the united kingdom evaluation of daily activities questionnaire in rheumatoid arthritis: Psychometric testing. <i>Annals of the Rheumatic Diseases</i> , 2013, 71, 753.1-753.	0.5	0
92	Is the Epworth Sleepiness Scale Suitable for Use in Stroke?. <i>Topics in Stroke Rehabilitation</i> , 2013, 20, 493-499.	1.0	13
93	Rasch analysis of the WHOQOL-BREF in post polio syndrome. <i>Journal of Rehabilitation Medicine</i> , 2013, 45, 873-880.	0.8	14
94	Screening for the risk of job loss in multiple sclerosis (MS): development of an MS-specific Work Instability Scale (MS-WIS). <i>Multiple Sclerosis Journal</i> , 2012, 18, 862-870.	1.4	42
95	The unidimensional self-efficacy scale for MS (USE-MS): developing a patient based and patient reported outcome. <i>Multiple Sclerosis Journal</i> , 2012, 18, 1326-1333.	1.4	25
96	Modifying the Medical Research Council grading system through Rasch analyses. <i>Brain</i> , 2012, 135, 1639-1649.	3.7	224
97	The use of the Visual Analogue Scale (VAS) in rehabilitation outcomes. <i>Journal of Rehabilitation Medicine</i> , 2012, 44, 609-610.	0.8	93
98	ACTIVLIM-Stroke: A Crosscultural Rasch-Built Scale of Activity Limitations in Patients With Stroke. <i>Stroke</i> , 2012, 43, 815-823.	1.0	41
99	Co-calibration of deliberate self harm (DSH) behaviours: Towards a common measurement metric. <i>Psychiatry Research</i> , 2012, 200, 26-34.	1.7	9
100	Is the Berg Balance Scale an Internally Valid and Reliable Measure of Balance Across Different Etiologies in Neurorehabilitation? A Revisited Rasch Analysis Study. <i>Archives of Physical Medicine and Rehabilitation</i> , 2012, 93, 1209-1216.	0.5	91
101	The use of raw scores from ordinal scales: Time to end malpractice?. <i>Journal of Rehabilitation Medicine</i> , 2012, 44, 97-98.	0.8	149
102	Depression and anxiety in patients with rheumatoid arthritis: prevalence rates based on a comparison of the Depression, Anxiety and Stress Scale (DASS) and the hospital, Anxiety and Depression Scale (HADS). <i>BMC Psychiatry</i> , 2012, 12, 6.	1.1	208
103	Validation of the Neurological Fatigue Index for stroke (NFI-Stroke). <i>Health and Quality of Life Outcomes</i> , 2012, 10, 51.	1.0	24
104	Reliability, construct validity and measurement potential of the ICF comprehensive core set for osteoarthritis. <i>BMC Musculoskeletal Disorders</i> , 2011, 12, 255.	0.8	34
105	Strategies for assessment and outcome measurement in Physical and Rehabilitation Medicine: An educational review. <i>Journal of Rehabilitation Medicine</i> , 2011, 43, 661-672.	0.8	61
106	Unified Balance Scale: Classic psychometric and clinical properties. <i>Journal of Rehabilitation Medicine</i> , 2011, 43, 445-453.	0.8	12
107	Unified Balance Scale: An activity-based, bed to community, and aetiology-independent measure of balance calibrated with Rasch analysis. <i>Journal of Rehabilitation Medicine</i> , 2011, 43, 435-444.	0.8	34
108	Past and present issues in Rasch analysis: The Functional Independence Measure (FIM) revisited. <i>Journal of Rehabilitation Medicine</i> , 2011, 43, 884-891.	0.8	111

#	ARTICLE	IF	CITATIONS
109	Can Neuropathic Screening Tools Be Used As Outcome Measures?. Pain Medicine, 2011, 12, 276-281.	0.9	9
110	Development of a patient reported outcome measure for fatigue in motor neurone disease: the Neurological Fatigue Index (NFI-MND). Health and Quality of Life Outcomes, 2011, 9, 101.	1.0	31
111	Validation of the World Health Organization disability assessment schedule II (WHODAS-II) in patients with osteoarthritis. Rheumatology International, 2011, 31, 339-346.	1.5	65
112	Cross-cultural validation of the Educational Needs Assessment Tool in RA in 7 European countries. BMC Musculoskeletal Disorders, 2011, 12, 110.	0.8	49
113	Rasch analysis of the hospital anxiety and depression scale (hads) for use in motor neurone disease. Health and Quality of Life Outcomes, 2011, 9, 82.	1.0	96
114	The interval scaling properties of the London Handicap Scale: an example from the adaptation of the scale for use in Turkey. Clinical Rehabilitation, 2011, 25, 248-255.	1.0	6
115	The Warwick-Edinburgh Mental Well-being Scale (WEMWBS): a valid and reliable tool for measuring mental well-being in diverse populations and projects. Journal of Epidemiology and Community Health, 2011, 65, A38-A39.	2.0	164
116	Construct Validity of the Holistic Complementary and Alternative Medicines Questionnaire (HCAMQ) – An Investigation Using Modern Psychometric Approaches. Evidence-based Complementary and Alternative Medicine, 2011, 2011, 1-8.	0.5	9
117	The Visual Analogue WOMAC 3.0 scale - internal validity and responsiveness of the VAS version. BMC Musculoskeletal Disorders, 2010, 11, 80.	0.8	65
118	Development of a patient reported outcome scale for fatigue in multiple sclerosis: The Neurological Fatigue Index (NFI-MS). Health and Quality of Life Outcomes, 2010, 8, 22.	1.0	88
119	Rasch analysis of the Modified Fatigue Impact Scale (MFIS) in multiple sclerosis. Journal of Neurology, Neurosurgery and Psychiatry, 2010, 81, 1049-1051.	0.9	76
120	Rasch analysis of the Beck Depression Inventory-II in a neurological rehabilitation sample. Disability and Rehabilitation, 2010, 32, 8-17.	0.9	61
121	Psychometric Properties of the Health Assessment Questionnaire Disability Index (HAQ-DI) and the Modified Health Assessment Questionnaire (MHAQ) in Patients with Knee Osteoarthritis. Turkish Journal of Rheumatology, 2010, 25, 147-155.	0.2	7
122	An application of computerised adaptive testing for measuring health status in patients with knee osteoarthritis. Disability and Rehabilitation, 2010, 32, 1928-1938.	0.9	10
123	Rasch analysis of the Fatigue Severity Scale in multiple sclerosis. Multiple Sclerosis Journal, 2009, 15, 81-87.	1.4	95
124	Adaptation and cross-cultural validation of the rheumatoid arthritis work instability scale (RA-WIS). Annals of the Rheumatic Diseases, 2009, 68, 1686-1690.	0.5	33
125	Modification and validation of the Lysholm Knee Scale to assess articular cartilage damage. Osteoarthritis and Cartilage, 2009, 17, 53-58.	0.6	60
126	Variability in depression prevalence in early rheumatoid arthritis: a comparison of the CES-D and HAD-D Scales. BMC Musculoskeletal Disorders, 2009, 10, 18.	0.8	38



#	ARTICLE	IF	CITATIONS
127	Reducing work disability in Ankylosing Spondylitis – development of a work instability scale for AS. BMC Musculoskeletal Disorders, 2009, 10, 68.	0.8	35
128	Rasch analysis provides new insights into the measurement properties of the neck disability index. Arthritis and Rheumatism, 2009, 61, 544-551.	6.7	88
129	Rasch model analysis of the Depression, Anxiety and Stress Scales (DASS). BMC Psychiatry, 2009, 9, 21.	1.1	133
130	Psychometric analysis of the Self-Harm Inventory using Rasch modelling. BMC Psychiatry, 2009, 9, 53.	1.1	28
131	Internal construct validity of the Warwick-Edinburgh Mental Well-being Scale (WEMWBS): a Rasch analysis using data from the Scottish Health Education Population Survey. Health and Quality of Life Outcomes, 2009, 7, 15.	1.0	700
132	Measuring the impact and distress of osteoarthritis from the patients' perspective. Health and Quality of Life Outcomes, 2009, 7, 37.	1.0	17
133	The development of the L-QoL: a quality-of-life instrument specific to systemic lupus erythematosus. Annals of the Rheumatic Diseases, 2009, 68, 196-200.	0.5	134
134	Development and validation of a needs-based quality of life instrument for osteoarthritis. Arthritis and Rheumatism, 2008, 59, 841-848.	6.7	53
135	An initial application of computerized adaptive testing (CAT) for measuring disability in patients with low back pain. BMC Musculoskeletal Disorders, 2008, 9, 166.	0.8	45
136	The development of a short measure of physical function for hip OA HOOS-Physical Function Shortform (HOOS-PS): an OARSI/OMERACT initiative. Osteoarthritis and Cartilage, 2008, 16, 551-559.	0.6	156
137	Measuring pain: issues of interpretation. Lancet, The, 2008, 372, 443.	6.3	3
138	Assessing Participation in Daily Living and the Effectiveness of Rehabilitation in Age Related Macular Degeneration Patients Using the Impact of Vision Impairment Scale. Ophthalmic Epidemiology, 2008, 15, 105-113.	0.8	74
139	The Foot Posture Index: Rasch Analysis of a Novel, Foot-Specific Outcome Measure. Archives of Physical Medicine and Rehabilitation, 2007, 88, 88-93.	0.5	184
140	Goal attainment scaling: Current methodological challenges. Disability and Rehabilitation, 2007, 29, 1583-1588.	0.9	93
141	Validation of the Middlesex Elderly Assessment of Mental State (MEAMS) as a cognitive screening test in patients with acquired brain injury in Turkey. Disability and Rehabilitation, 2007, 29, 315-321.	0.9	14
142	Quality of life in infants and children with atopic dermatitis: Addressing issues of differential item functioning across countries in multinational clinical trials. Health and Quality of Life Outcomes, 2007, 5, 45.	1.0	33
143	Internal construct validity of the Oxford Knee Scale: Evidence from Rasch measurement. Arthritis and Rheumatism, 2007, 57, 1363-1367.	6.7	57
144	The Rasch measurement model in rheumatology: What is it and why use it? When should it be applied, and what should one look for in a Rasch paper?. Arthritis and Rheumatism, 2007, 57, 1358-1362.	6.7	1,178

#	ARTICLE	IF	CITATIONS
145	Keeping nurses working: development and psychometric testing of the Nurse-Work Instability Scale (Nurse-WIS). <i>Journal of Advanced Nursing</i> , 2007, 57, 543-551.	1.5	26
146	An introduction to the Rasch measurement model: An example using the Hospital Anxiety and Depression Scale (HADS). <i>British Journal of Clinical Psychology</i> , 2007, 46, 1-18.	1.7	768
147	Assessing normative cut points through differential item functioning analysis: an example from the adaptation of the Middlesex Elderly Assessment of Mental State (MEAMS) for use as a cognitive screening test in Turkey. <i>Health and Quality of Life Outcomes</i> , 2006, 4, 18.	1.0	7
148	Cross-diagnostic validity in a generic instrument: an example from the Functional Independence Measure in Scandinavia. <i>Health and Quality of Life Outcomes</i> , 2006, 4, 55.	1.0	32
149	Cross-cultural validity of FIM in spinal cord injury. <i>Spinal Cord</i> , 2006, 44, 746-752.	0.9	44
150	Evaluation of the Edinburgh Post Natal Depression Scale using Rasch analysis. <i>BMC Psychiatry</i> , 2006, 6, 28.	1.1	62
151	Identifying patients at risk of nursing home admission: The Leeds Elderly Assessment Dependency Screening tool (LEADS). <i>BMC Health Services Research</i> , 2006, 6, 31.	0.9	20
152	Screening for job loss: Development of a work instability scale for traumatic brain injury. <i>Brain Injury</i> , 2006, 20, 835-843.	0.6	35
153	Admission to hospital following head injury in England: Incidence and socio-economic associations. <i>BMC Public Health</i> , 2005, 5, 21.	1.2	46
154	Development of a foot impact scale for rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2005, 53, 418-422.	6.7	103
155	CROSS-CULTURAL VALIDITY OF FUNCTIONAL INDEPENDENCE MEASURE ITEMS IN STROKE: A STUDY USING RASCH ANALYSIS. <i>Journal of Rehabilitation Medicine</i> , 2005, 37, 23-31.	0.8	57
156	Psychometric properties of the Mini-Mental State Examination in patients with acquired brain injury in Turkey. <i>Journal of Rehabilitation Medicine</i> , 2005, 37, 306-311.	0.8	36
157	Construct validity and reliability of the Rivermead Post-Concussion Symptoms Questionnaire. <i>Clinical Rehabilitation</i> , 2005, 19, 878-887.	1.0	231
158	Development of the PsAQoL: a quality of life instrument specific to psoriatic arthritis. <i>Annals of the Rheumatic Diseases</i> , 2004, 63, 162-169.	0.5	218
159	Application of Rasch Analysis in the Development and Application of Quality of Life Instruments. <i>Value in Health</i> , 2004, 7, S22-S26.	0.1	350
160	Issues in cross-cultural validity: Example from the adaptation, reliability, and validity testing of a Turkish version of the Stanford Health Assessment Questionnaire. <i>Arthritis and Rheumatism</i> , 2004, 51, 14-19.	6.7	210
161	Assessing and Adjusting for Cross-Cultural Validity of Impairment and Activity Limitation Scales Through Differential Item Functioning Within the Framework of the Rasch Model. <i>Medical Care</i> , 2004, 42, 37.	1.1	256
162	Predicting outcome for older people in a hospital setting: which scales are appropriate?. <i>International Journal of Therapy and Rehabilitation</i> , 2004, 11, 25-30.	0.1	6

#	ARTICLE	IF	CITATIONS
163	Development of the BD-QoL: a quality of life measure specific to Behçet's disease. <i>Journal of Rheumatology</i> , 2004, 31, 931-7.	1.0	45
164	Adaptation and validation of the Turkish version of the Rheumatoid Arthritis Quality of Life Scale. <i>Rheumatology International</i> , 2003, 23, 21-26.	1.5	45
165	Development of the ASQoL: a quality of life instrument specific to ankylosing spondylitis. <i>Annals of the Rheumatic Diseases</i> , 2003, 62, 20-26.	0.5	480
166	Assessing comparability of dressing disability in different countries by response conversion. <i>European Journal of Public Health</i> , 2003, 13, 15-19.	0.1	9
167	Assessing comparability of dressing disability in different countries by response conversion. <i>European Journal of Public Health</i> , 2003, 13, 15-19.	0.1	8
168	Team approach versus ad hoc health services for young people with physical disabilities: a retrospective cohort study. <i>Lancet</i> , 2002, 360, 1280-1286.	6.3	107
169	Developing a disease-specific quality of life measure for people with multiple sclerosis. <i>Clinical Rehabilitation</i> , 2001, 15, 247-258.	1.0	117
170	Adaptation of the Functional Independence Measure for use in Turkey. <i>Clinical Rehabilitation</i> , 2001, 15, 311-319.	1.0	182
171	THE USE OF OUTCOME MEASURES IN PHYSICAL MEDICINE AND REHABILITATION WITHIN EUROPE. <i>Journal of Rehabilitation Medicine</i> , 2001, 33, 273-278.	0.8	128
172	Measuring the function of children with juvenile arthritis. <i>British Journal of Rheumatology</i> , 2001, 40, 1274-1278.	2.5	22
173	Measuring outcome. <i>British Medical Bulletin</i> , 2000, 56, 287-295.	2.7	6
174	The Leeds assessment scale of handicap: its operationalisation, reliability, validity and responsiveness in in-patient rehabilitation. <i>Disability and Rehabilitation</i> , 2000, 22, 529-538.	0.9	3
175	Evaluation of a community-based neuropsychological rehabilitation service for people with traumatic brain injury. <i>NeuroRehabilitation</i> , 1999, 13, 147-155.	0.5	14
176	The persistence of mood disorders following traumatic brain injury: a 1 year follow-up. <i>Brain Injury</i> , 1999, 13, 547-553.	0.6	30
177	Mood disorders following traumatic brain injury: identifying the extent of the problem and the people at risk. <i>Brain Injury</i> , 1998, 12, 177-190.	0.6	95
178	Models of disability: A critical perspective. <i>Disability and Rehabilitation</i> , 1997, 19, 478-479.	0.9	13
179	Prevalence of hip problems in the population aged 55 years and over: access to specialist care and future demand for hip arthroplasty. <i>Rheumatology</i> , 1997, 36, 74-76.	0.9	47
180	Outcome following stroke. <i>Disability and Rehabilitation</i> , 1997, 19, 278-284.	0.9	63

#	ARTICLE	IF	CITATIONS
181	The Barthel Index: an ordinal score or interval level measure?. <i>Clinical Rehabilitation</i> , 1996, 10, 301-308.	1.0	44
182	ARE WE MAKING THE MOST OF THE STANFORD HEALTH ASSESSMENT QUESTIONNAIRE?. <i>Rheumatology</i> , 1996, 35, 574-578.	0.9	60
183	Prevalence of self reported stroke in a population in northern England.. <i>Journal of Epidemiology and Community Health</i> , 1996, 50, 140-143.	2.0	83
184	Quality of life--a measure too far?. <i>Annals of the Rheumatic Diseases</i> , 1995, 54, 439-440.	0.5	23
185	The long-term outcome of head injury: Implications for service planning. <i>Brain Injury</i> , 1995, 9, 595-605.	0.6	50
186	Prevalence of knee problems in the population aged 55 years and over: identifying the need for knee arthroplasty. <i>BMJ: British Medical Journal</i> , 1995, 310, 1291-1293.	2.4	87