Ilana R Cliffer

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2749535/publications.pdf

Version: 2024-02-01

1937685 1720034 9 52 4 7 citations h-index g-index papers 9 9 9 61 docs citations times ranked citing authors all docs

#	Article	IF	Citations
1	Monthly measurement of child lengths between 6 and 27 months of age in Burkina Faso reveals both chronic and episodic growth faltering. American Journal of Clinical Nutrition, 2022, 115, 94-104.	4.7	3
2	Linear Growth Spurts are Preceded by Higher Weight Gain Velocity and Followed by Weight Slowdowns Among Rural Children in Burkina Faso: A Longitudinal Study. Journal of Nutrition, 2022, 152, 1963-1973.	2.9	4
3	Seasonality of Child Growth: High Temperatures Coincide with Growth Faltering among Young Children in Burkina Faso. Current Developments in Nutrition, 2022, 6, 74.	0.3	0
4	Fortified blended flour supplements displace plain cereals in feeding of young children. Maternal and Child Nutrition, 2021, 17, e13089.	3.0	3
5	Effectiveness and cost-effectiveness of 4 supplementary foods for treating moderate acute malnutrition: results from a cluster-randomized intervention trial in Sierra Leone. American Journal of Clinical Nutrition, 2021, 114, 973-985.	4.7	12
6	Advancing Nutrition in the International Food Assistance Agenda: Progress and Future Directions Identified at the 2018 Food Assistance for Nutrition Evidence Summit. Food and Nutrition Bulletin, 2020, 41, 8-17.	1.4	6
7	Impact of stakeholder perspectives on cost-effectiveness estimates of four specialized nutritious foods for preventing stunting and wasting in children 6–23 months in Burkina Faso. Nutrition Journal, 2020, 19, 20.	3.4	9
8	Cost-Effectiveness of 4 Specialized Nutritious Foods in the Prevention of Stunting and Wasting in Children Aged 6–23 Months in Burkina Faso: A Geographically Randomized Trial. Current Developments in Nutrition, 2020, 4, nzaa006.	0.3	10
9	Factors that May Influence the Effectiveness of 4 Specialized Nutritious Foods in the Prevention of Stunting and Wasting in Children Aged 6–23 Months in Burkina Faso. Current Developments in Nutrition, 2020, 4, nzaa002.	0.3	5