## Adelaida M M Celaya

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2757527/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Mutations in L-type amino acid transporter-2 support SLC7A8 as a novel gene involved in age-related hearing loss. ELife, 2018, 7, .	6.0	38
2	<i>G6PD</i> overexpression protects from oxidative stress and ageâ€related hearing loss. Aging Cell, 2020, 19, e13275.	6.7	37
3	Transforming growth factor β1 inhibition protects from noise-induced hearing loss. Frontiers in Aging Neuroscience, 2015, 7, 32.	3.4	34
4	MPZL2, Encoding the Epithelial Junctional Protein Myelin Protein Zero-like 2, Is Essential for Hearing in Man and Mouse. American Journal of Human Genetics, 2018, 103, 74-88.	6.2	34
5	Deficit of mitogen-activated protein kinase phosphatase 1 (DUSP1) accelerates progressive hearing loss. ELife, 2019, 8, .	6.0	21
6	A Comparative Study of Drug Delivery Methods Targeted to the Mouse Inner Ear: Bullostomy <em>Versus</em> Transtympanic Injection. Journal of Visualized Experiments, 2017, , .	0.3	12
7	IGF-1 Haploinsufficiency Causes Age-Related Chronic Cochlear Inflammation and Increases Noise-Induced Hearing Loss. Cells, 2021, 10, 1686.	4.1	12
8	Dual-Specificity Phosphatase 1 (DUSP1) Has a Central Role in Redox Homeostasis and Inflammation in the Mouse Cochlea. Antioxidants, 2021, 10, 1351.	5.1	11
9	Betaineâ€homocysteine <i>S</i> â€methyltransferase deficiency causes increased susceptibility to noiseâ€induced hearing loss associated with plasma hyperhomocysteinemia. FASEB Journal, 2019, 33, 5942-5956.	0.5	7
10	Therapeutic efficiency of the APAFâ€1 antagonist LPT99 in a rat model of cisplatinâ€induced hearing loss. Clinical and Translational Medicine, 2021, 11, e363.	4.0	6