Stephane Campeau

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2758337/publications.pdf

Version: 2024-02-01

24 papers 738 citations

687363 13 h-index 610901 24 g-index

25 all docs

25 docs citations

25 times ranked

798 citing authors

#	Article	IF	CITATIONS
1	Diversity of diatoms, benthic macroinvertebrates, and fish varies in response to different environmental correlates in Arctic rivers across North America. Freshwater Biology, 2022, 67, 95-115.	2.4	15
2	Responses of soil erosion to warming and wetting in a cold Canadian agricultural catchment. Catena, 2021, 201, 105184.	5.0	7
3	Impacts of climate change on the hydrology of northern midlatitude cold regions. Progress in Physical Geography, 2020, 44, 338-375.	3.2	34
4	Shifting Hydrological Processes in a Canadian Agroforested Catchment due to a Warmer and Wetter Climate. Water (Switzerland), 2020, 12, 739.	2.7	12
5	Local and regional drivers of taxonomic homogenization in stream communities along a land use gradient. Global Ecology and Biogeography, 2019, 28, 1597-1609.	5.8	21
6	Three-dimensional hydrostratigraphical modelling of the regional aquifer system of the St. Maurice Delta Complex (St. Lawrence Lowlands, Canada). Canadian Water Resources Journal, 2018, 43, 92-112.	1.2	7
7	<i>Gomphonema caperatum</i> sp. nov., <i>G. obstipum</i> sp. nov. and similar taxa from rivers of North America. Diatom Research, 2017, 32, 59-73.	1.2	9
8	Landscape diversity and forest edge density regulate stream water quality in agricultural catchments. Ecological Indicators, 2017, 72, 627-639.	6.3	92
9	Assemblage diversity, cell density and within-slide variability: Implications for quality assurance/quality control and uncertainty assessment in diatom-based monitoring. Ecological Indicators, 2016, 69, 415-421.	6.3	8
10	Using diatoms to monitor stream biological integrity in Eastern Canada: An overview of 10 years of index development and ongoing challenges. Science of the Total Environment, 2014, 475, 187-200.	8.0	62
11	The Impact of "Man-Made Hydrological Drought―on Plant Species Abundance in the Low-Flow Channel Downstream from the Matawin Dam, Quebec. Water (Switzerland), 2013, 5, 875-892.	2.7	6
12	Modeling the response time of diatom assemblages to simulated water quality improvement and degradation in running waters. Canadian Journal of Fisheries and Aquatic Sciences, 2011, 68, 487-497.	1.4	16
13	Fishing for diatoms: fish gut analysis reveals water quality changes over a 75-year period. Journal of Paleolimnology, 2010, 43, 121-130.	1.6	10
14	Algae-based Biomonitoring: Predicting Diatom Reference Communities in Unpolluted Streams using Classification Trees, Random Forests, and Artificial Neural Networks. Water Quality Research Journal of Canada, 2010, 45, 413-425.	2.7	6
15	The Eastern Canadian Diatom Index (IDEC) Version 2.0: Including Meaningful Ecological Classes and an Expanded Coverage Area that Encompasses Additional Geological Characteristics. Water Quality Research Journal of Canada, 2010, 45, 463-477.	2.7	11
16	Defining ecological thresholds to determine class boundaries in a bioassessment tool: The case of the Eastern Canadian Diatom Index (IDEC). Ecological Indicators, 2010, 10, 980-989.	6.3	33
17	The effect of excluding diatom taxa and reducing taxonomic resolution on multivariate analyses and stream bioassessment. Ecological Indicators, 2009, 9, 213-225.	6.3	71
18	Are diatoms good integrators of temporal variability in stream water quality?. Freshwater Biology, 2008, 53, 827-841.	2.4	87

#	Article	IF	CITATION
19	Diatom reference communities in Québec (Canada) streams based on Kohonen self-organizing maps and multivariate analyses. Canadian Journal of Fisheries and Aquatic Sciences, 2006, 63, 2087-2106.	1.4	37
20	A diatom-based index for the biological assessment of eastern Canadian rivers: an application of correspondence analysis (CA). Canadian Journal of Fisheries and Aquatic Sciences, 2006, 63, 1793-1811.	1.4	78
21	Diatoms and biomonitoring: should cell size be accounted for?. Hydrobiologia, 2006, 573, 1-16.	2.0	30
22	A large-scale stream benthic diatom database. Hydrobiologia, 2005, 542, 151-163.	2.0	7
23	Late Holocene diatom biostratigraphy and sea-level changes in the southeastern Beaufort Sea. Canadian Journal of Earth Sciences, 2000, 37, 63-80.	1.3	20
24	Diatoms as quantitative paleodepth indicators in coastal areas of the southeastern Beaufort Sea, Arctic Ocean. Palaeogeography, Palaeoclimatology, Palaeoecology, 1999, 146, 67-97.	2.3	27