

Isao Oze

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2759703/publications.pdf>

Version: 2024-02-01

110
papers

2,758
citations

172207

29
h-index

223531

46
g-index

112
all docs

112
docs citations

112
times ranked

5170
citing authors

#	ARTICLE	IF	CITATIONS
1	Moderate-to-vigorous Physical Activity and Sedentary Behavior Are Independently Associated With Renal Function: A Cross-sectional Study. <i>Journal of Epidemiology</i> , 2023, 33, 285-293.	1.1	7
2	Coffee and tea consumption and mortality from all causes, cardiovascular disease and cancer: a pooled analysis of prospective studies from the Asia Cohort Consortium. <i>International Journal of Epidemiology</i> , 2022, 51, 626-640.	0.9	37
3	Alcohol intake and stomach cancer risk in Japan: A pooled analysis of six cohort studies. <i>Cancer Science</i> , 2022, 113, 261-276.	1.7	3
4	The association of reproductive history with hypertension and obesity according to menopausal status: the J-MICC Study. <i>Hypertension Research</i> , 2022, 45, 708-714.	1.5	2
5	Body Mass Index and Thyroid Cancer Risk: A Pooled Analysis of Half a Million Men and Women in the Asia Cohort Consortium. <i>Thyroid</i> , 2022, 32, 306-314.	2.4	17
6	Associations of breastfeeding history with metabolic syndrome and cardiovascular risk factors in community-dwelling parous women: The Japan Multi-Institutional Collaborative Cohort Study. <i>PLoS ONE</i> , 2022, 17, e0262252.	1.1	5
7	New insights into the genetic contribution of <i>ALDH2</i> rs671 in pancreatic carcinogenesis: Evaluation by mediation analysis. <i>Cancer Science</i> , 2022, 113, 1441-1450.	1.7	3
8	Trends in the incidence of head and neck cancer by subsite between 1993 and 2015 in Japan. <i>Cancer Medicine</i> , 2022, 11, 1553-1560.	1.3	29
9	Association between body mass index and oesophageal cancer mortality: a pooled analysis of prospective cohort studies with >800,000 individuals in the Asia Cohort Consortium. <i>International Journal of Epidemiology</i> , 2022, 51, 1190-1203.	0.9	8
10	A genome-wide association study on adherence to low-carbohydrate diets in Japanese. <i>European Journal of Clinical Nutrition</i> , 2022, , .	1.3	1
11	Large-scale Integrated Analysis of Genetics and Metabolomic Data Reveals Potential Links Between Lipids and Colorectal Cancer Risk. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2022, 31, 1216-1226.	1.1	3
12	Association between germline pathogenic variants and breast cancer risk in Japanese women: The HERPACC study. <i>Cancer Science</i> , 2022, 113, 1451-1462.	1.7	2
13	Developing and validating polygenic risk scores for colorectal cancer risk prediction in East Asians. <i>International Journal of Cancer</i> , 2022, 151, 1726-1736.	2.3	5
14	Differential Effect of Polymorphisms on Body Mass Index Across the Life Course of Japanese: The Japan Multi-Institutional Collaborative Cohort Study. <i>Journal of Epidemiology</i> , 2021, 31, 172-179.	1.1	5
15	A genome-wide association study in Japanese identified one variant associated with a preference for a Japanese dietary pattern. <i>European Journal of Clinical Nutrition</i> , 2021, 75, 937-945.	1.3	8
16	A genome-wide association study on fish consumption in a Japanese population—the Japan Multi-Institutional Collaborative Cohort study. <i>European Journal of Clinical Nutrition</i> , 2021, 75, 480-488.	1.3	5
17	Impact of <i>PSCA</i> Polymorphisms on the Risk of Duodenal Ulcer. <i>Journal of Epidemiology</i> , 2021, 31, 12-20.	1.1	9
18	Body mass index and colorectal cancer risk: A Mendelian randomization study. <i>Cancer Science</i> , 2021, 112, 1579-1588.	1.7	25

#	ARTICLE	IF	CITATIONS
19	A genome-wide association study on confection consumption in a Japanese population: the Japan Multi-Institutional Collaborative Cohort Study. <i>British Journal of Nutrition</i> , 2021, 126, 1843-1851.	1.2	6
20	Reproducibility and validity of food group intake in a short food frequency questionnaire for the middle-aged Japanese population. <i>Environmental Health and Preventive Medicine</i> , 2021, 26, 28.	1.4	29
21	A Personal Breast Cancer Risk Stratification Model Using Common Variants and Environmental Risk Factors in Japanese Females. <i>Cancers</i> , 2021, 13, 3796.	1.7	4
22	Study Profile of the Japan Multi-institutional Collaborative Cohort (J-MICC) Study. <i>Journal of Epidemiology</i> , 2021, 31, 660-668.	1.1	41
23	A genome-wide association study on meat consumption in a Japanese population: the Japan Multi-Institutional Collaborative Cohort study. <i>Journal of Nutritional Science</i> , 2021, 10, e61.	0.7	3
24	Association of skipping breakfast and short sleep duration with the prevalence of metabolic syndrome in the general Japanese population: Baseline data from the Japan Multi-Institutional Collaborative cohort study. <i>Preventive Medicine Reports</i> , 2021, 24, 101613.	0.8	6
25	Risk Prediction for Gastric Cancer Using GWAS-Identified Polymorphisms, Helicobacter pylori Infection and Lifestyle-Related Risk Factors in a Japanese Population. <i>Cancers</i> , 2021, 13, 5525.	1.7	3
26	Genome-wide association study of serum prostate-specific antigen levels based on 1000 Genomes imputed data in Japanese: the Japan Multi-Institutional Collaborative Cohort Study. <i>Nagoya Journal of Medical Science</i> , 2021, 83, 183-194.	0.6	1
27	Alcohol Drinking and Bladder Cancer Risk From a Pooled Analysis of Ten Cohort Studies in Japan. <i>Journal of Epidemiology</i> , 2020, 30, 309-313.	1.1	2
28	Combined effect of weight gain within normal weight range and parental hypertension on the prevalence of hypertension; from the J-MICC Study. <i>Journal of Human Hypertension</i> , 2020, 34, 125-131.	1.0	0
29	The interaction between ABCA1 polymorphism and physical activity on the HDL-cholesterol levels in a Japanese population. <i>Journal of Lipid Research</i> , 2020, 61, 86-94.	2.0	11
30	Identification of Novel Loci and New Risk Variant in Known Loci for Colorectal Cancer Risk in East Asians. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2020, 29, 477-486.	1.1	25
31	Risk of second primary malignancies after definitive treatment for esophageal cancer: A competing risk analysis. <i>Cancer Medicine</i> , 2020, 9, 394-400.	1.3	12
32	Association between Socioeconomic Status and Digestive Tract Cancers: A Case-Control Study. <i>Cancers</i> , 2020, 12, 3258.	1.7	9
33	Association of Dietary Acid Load with the Prevalence of Metabolic Syndrome among Participants in Baseline Survey of the Japan Multi-Institutional Collaborative Cohort Study. <i>Nutrients</i> , 2020, 12, 1605.	1.7	23
34	Across-Site Differences in the Mechanism of Alcohol-Induced Digestive Tract Carcinogenesis: An Evaluation by Mediation Analysis. <i>Cancer Research</i> , 2020, 80, 1601-1610.	0.4	22
35	Relationship between the strength of craving as assessed by the Tobacco Craving Index and success of quitting smoking in Japanese smoking cessation therapy. <i>PLoS ONE</i> , 2020, 15, e0243374.	1.1	3
36	Revisit of an unanswered question by pooled analysis of eight cohort studies in Japan: Does cigarette smoking and alcohol drinking have interaction for the risk of esophageal cancer?. <i>Cancer Medicine</i> , 2019, 8, 6414-6425.	1.3	22

#	ARTICLE	IF	CITATIONS
37	Association of BMI, Smoking, and Alcohol with Multiple Myeloma Mortality in Asians: A Pooled Analysis of More than 800,000 Participants in the Asia Cohort Consortium. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2019, 28, 1861-1867.	1.1	11
38	Associations of Nutrient Patterns with the Prevalence of Metabolic Syndrome: Results from the Baseline Data of the Japan Multi-Institutional Collaborative Cohort Study. <i>Nutrients</i> , 2019, 11, 990.	1.7	24
39	Networks for early career epidemiologists around the world: the current status and future directions. <i>International Journal of Epidemiology</i> , 2019, 48, 1021-1023.	0.9	1
40	Genome-wide association meta-analysis and Mendelian randomization analysis confirm the influence of ALDH2 on sleep duration in the Japanese population. <i>Sleep</i> , 2019, 42, .	0.6	16
41	Genome-wide meta-analysis identifies multiple novel loci associated with serum uric acid levels in Japanese individuals. <i>Communications Biology</i> , 2019, 2, 115.	2.0	66
42	Large-Scale Genome-Wide Association Study of East Asians Identifies Loci Associated With Risk for Colorectal Cancer. <i>Gastroenterology</i> , 2019, 156, 1455-1466.	0.6	111
43	GWAS analysis reveals a significant contribution of PSCA to the risk of <i>Helicobacter pylori</i> -induced gastric atrophy. <i>Carcinogenesis</i> , 2019, 40, 661-668.	1.3	13
44	Association of genetic risk score and chronic kidney disease in a Japanese population. <i>Nephrology</i> , 2019, 24, 670-673.	0.7	12
45	Trends in Small-Cell Lung Cancer Survival in 1993–2006 Based on Population-Based Cancer Registry Data in Japan. <i>Journal of Epidemiology</i> , 2019, 29, 347-353.	1.1	13
46	Phase 2 Study of Afatinib Alone or Combined With Bevacizumab in Chemonaive Patients With Advanced Non-Small-Cell Lung Cancer Harboring EGFR Mutations: AfaBev-CS Study Protocol. <i>Clinical Lung Cancer</i> , 2019, 20, 134-138.	1.1	19
47	Perceived Barriers to Career Progression Among Early-Career Epidemiologists: Report of a Workshop at the 22nd World Congress of Epidemiology. <i>Journal of Epidemiology</i> , 2019, 29, 38-41.	1.1	10
48	Association between ALDH2 and ADH1B polymorphisms, alcohol drinking and gastric cancer: a replication and mediation analysis. <i>Gastric Cancer</i> , 2018, 21, 936-945.	2.7	36
49	Changes in self-efficacy associated with success in quitting smoking in participants in Japanese smoking cessation therapy. <i>International Journal of Nursing Practice</i> , 2018, 24, e12647.	0.8	13
50	Genome-wide association study identifies seven novel susceptibility loci for primary open-angle glaucoma. <i>Human Molecular Genetics</i> , 2018, 27, 1486-1496.	1.4	111
51	A genome-wide association study in the Japanese population identifies the 12q24 locus for habitual coffee consumption: The J-MICC Study. <i>Scientific Reports</i> , 2018, 8, 1493.	1.6	32
52	Smoking and subsequent risk of acute myeloid leukaemia: A pooled analysis of 9 cohort studies in Japan. <i>Hematological Oncology</i> , 2018, 36, 262-268.	0.8	10
53	Heterogeneous impact of smoking on major salivary gland cancer according to histopathological subtype: A case-control study. <i>Cancer</i> , 2018, 124, 118-124.	2.0	21
54	Establishment and validation of prognostic nomograms in first-line metastatic gastric cancer patients. <i>Journal of Gastrointestinal Oncology</i> , 2018, 9, 52-63.	0.6	26

#	ARTICLE	IF	CITATIONS
55	Genome-wide association study identifies gastric cancer susceptibility loci at 12q24.11 and 20q11.21. <i>Cancer Science</i> , 2018, 109, 4015-4024.	1.7	39
56	Genome-Wide Association Study of Renal Function Traits: Results from the Japan Multi-Institutional Collaborative Cohort Study. <i>American Journal of Nephrology</i> , 2018, 47, 304-316.	1.4	18
57	Improvement in 5-Year Relative Survival in Cancer of the Corpus Uteri From 1993-2000 to 2001-2006 in Japan. <i>Journal of Epidemiology</i> , 2018, 28, 75-80.	1.1	6
58	Cigarette smoking, alcohol drinking, and oral cavity and pharyngeal cancer in the Japanese: a population-based cohort study in Japan. <i>European Journal of Cancer Prevention</i> , 2018, 27, 171-179.	0.6	19
59	Genomewide Association Study of Leisure-Time Exercise Behavior in Japanese Adults. <i>Medicine and Science in Sports and Exercise</i> , 2018, 50, 2433-2441.	0.2	36
60	Development of a prediction model and estimation of cumulative risk for upper aerodigestive tract cancer on the basis of the aldehyde dehydrogenase 2 genotype and alcohol consumption in a Japanese population. <i>European Journal of Cancer Prevention</i> , 2017, 26, 38-47.	0.6	24
61	Changes in trends in colorectal cancer incidence rate by anatomic site between 1978 and 2004 in Japan. <i>European Journal of Cancer Prevention</i> , 2017, 26, 269-276.	0.6	23
62	Cognitive, behavioural and psychosocial factors associated with successful and maintained quit smoking status among patients who received smoking cessation intervention with nurses' counselling. <i>Journal of Advanced Nursing</i> , 2017, 73, 1681-1695.	1.5	13
63	Plasma microRNA-103, microRNA-107, and microRNA-194 levels are not biomarkers for human diffuse gastric cancer. <i>Journal of Cancer Research and Clinical Oncology</i> , 2017, 143, 551-554.	1.2	10
64	The more from East-Asian, the better: risk prediction of colorectal cancer risk by GWAS-identified SNPs among Japanese. <i>Journal of Cancer Research and Clinical Oncology</i> , 2017, 143, 2481-2492.	1.2	12
65	Heterogeneous impact of alcohol consumption according to treatment method on survival in head and neck cancer: A prospective study. <i>Cancer Science</i> , 2017, 108, 91-100.	1.7	38
66	Genetic Variants of <i>RAMP2</i> and <i>CLR</i> are Associated with Stroke. <i>Journal of Atherosclerosis and Thrombosis</i> , 2017, 24, 1267-1281.	0.9	11
67	Japanese Nurses' Perceptions Toward Tobacco Use Intervention for Hospitalized Cancer Patients Who Entered End of Life. <i>Cancer Nursing</i> , 2016, 39, E45-E51.	0.7	3
68	Bone Scan Index predicts skeletal-related events in patients with metastatic breast cancer. <i>SpringerPlus</i> , 2016, 5, 1095.	1.2	11
69	Prognostic Value of Drinking Status and Aldehyde Dehydrogenase 2 Polymorphism in Patients With Head and Neck Squamous Cell Carcinoma. <i>Journal of Epidemiology</i> , 2016, 26, 292-299.	1.1	16
70	A risk prediction model for colorectal cancer using genome-wide association study-identified polymorphisms and established risk factors among Japanese: results from two independent case-control studies. <i>European Journal of Cancer Prevention</i> , 2016, 25, 500-507.	0.6	16
71	Aldehyde dehydrogenase 2 (<i>ALDH2</i>) and alcohol dehydrogenase 1B (<i>ADH1B</i>) polymorphisms exacerbate bladder cancer risk associated with alcohol drinking: gene-environment interaction. <i>Carcinogenesis</i> , 2016, 37, 583-588.	1.3	32
72	Genetic variants of <i>SLC17A1</i> are associated with cholesterol homeostasis and hyperhomocysteinaemia in Japanese men. <i>Scientific Reports</i> , 2015, 5, 15888.	1.6	7

#	ARTICLE	IF	CITATIONS
73	Impact of metallothionein gene polymorphisms on the risk of lung cancer in a Japanese population. <i>Molecular Carcinogenesis</i> , 2015, 54, E122-8.	1.3	12
74	Association between brain-muscle-ARNT-like protein-2 (BMAL2) gene polymorphism and type 2 diabetes mellitus in obese Japanese individuals: A cross-sectional analysis of the Japan Multi-institutional Collaborative Cohort Study. <i>Diabetes Research and Clinical Practice</i> , 2015, 110, 301-308.	1.1	16
75	A phase II study of cisplatin plus S-1 with concurrent thoracic radiotherapy for locally advanced non-small-cell lung cancer: The Okayama Lung Cancer Study Group Trial 0501. <i>Lung Cancer</i> , 2015, 87, 141-147.	0.9	30
76	Lack of Association between the BIM Deletion Polymorphism and the Risk of Lung Cancer with and without EGFR Mutations. <i>Journal of Thoracic Oncology</i> , 2015, 10, 59-66.	0.5	13
77	Phase II Trial of Gefitinib in Combination with Bevacizumab as First-Line Therapy for Advanced Non-Small Cell Lung Cancer with Activating EGFR Gene Mutations: The Okayama Lung Cancer Study Group Trial 1001. <i>Journal of Thoracic Oncology</i> , 2015, 10, 486-491.	0.5	93
78	Polymorphisms in CYP19A1, HSD17B1 and HSD17B2 genes and serum sex hormone level among postmenopausal Japanese women. <i>Maturitas</i> , 2015, 82, 394-401.	1.0	9
79	Clinical Characteristics Associated with Long-term Survival in Metastatic Gastric Cancer after Systemic Chemotherapy. <i>Asian Pacific Journal of Cancer Prevention</i> , 2015, 16, 5433-5438.	0.5	6
80	Burden of Total and Cause-Specific Mortality Related to Tobacco Smoking among Adults Aged ≥45 Years in Asia: A Pooled Analysis of 21 Cohorts. <i>PLoS Medicine</i> , 2014, 11, e1001631.	3.9	98
81	Coffee and green tea consumption is associated with upper aerodigestive tract cancer in Japan. <i>International Journal of Cancer</i> , 2014, 135, 391-400.	2.3	30
82	Varenicline Is More Effective in Attenuating Weight Gain Than Nicotine Patch 12 Months After the End of Smoking Cessation Therapy: An Observational Study in Japan. <i>Nicotine and Tobacco Research</i> , 2014, 16, 1026-1029.	1.4	11
83	Induction of cytotoxic T cells as a novel independent survival factor in malignant melanoma with percutaneous peptide immunization. <i>Journal of Dermatological Science</i> , 2014, 75, 43-48.	1.0	14
84	Methylation Status of Blood Leukocyte DNA and Risk of Gastric Cancer in a High-Risk Chinese Population. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2014, 23, 2019-2026.	1.1	10
85	Long-term survival and conditional survival of cancer patients in Japan using population-based cancer registry data. <i>Cancer Science</i> , 2014, 105, 1480-1486.	1.7	131
86	Polymorphisms in PPAR Genes (PPARD, PPARG, and PPARGC1A) and the Risk of Chronic Kidney Disease in Japanese: Cross-Sectional Data from the J-MICC Study. <i>PPAR Research</i> , 2013, 1-8.	1.1	10
87	Polymorphisms in Base Excision Repair Genes Are Associated With Endometrial Cancer Risk Among Postmenopausal Japanese Women. <i>International Journal of Gynecological Cancer</i> , 2013, 23, 1561-1568.	1.2	17
88	Factors Associated With Weight Gain After Smoking Cessation Therapy in Japan. <i>Nursing Research</i> , 2013, 62, 414-421.	0.8	7
89	The aldehyde dehydrogenase 2 (ALDH2) Glu504Lys polymorphism interacts with alcohol drinking in the risk of stomach cancer. <i>Carcinogenesis</i> , 2013, 34, 1510-1515.	1.3	74
90	Diabetes mellitus and cancer risk: Pooled analysis of eight cohort studies in Japan. <i>Cancer Science</i> , 2013, 104, 1499-1507.	1.7	94

#	ARTICLE	IF	CITATIONS
91	Cigarette Smoking and Pancreatic Cancer Risk: A Revisit with an Assessment of the Nicotine Dependence Phenotype. <i>Asian Pacific Journal of Cancer Prevention</i> , 2013, 14, 4409-4413.	0.5	6
92	Inverse association between yoghurt intake and upper aerodigestive tract cancer risk in a Japanese population. <i>European Journal of Cancer Prevention</i> , 2012, 21, 453-459.	0.6	9
93	Folate, alcohol, and aldehyde dehydrogenase 2 polymorphism and the risk of oral and pharyngeal cancer in Japanese. <i>European Journal of Cancer Prevention</i> , 2012, 21, 193-198.	0.6	31
94	Time to First Cigarette and Upper Aerodigestive Tract Cancer Risk in Japan. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2012, 21, 1986-1992.	1.1	12
95	Favorable Response of Heavily Treated Wilms' Tumor to Paclitaxel and Carboplatin. <i>Onkologie</i> , 2012, 35, 283-286.	1.1	6
96	Cigarette Smoking and Esophageal Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence Among the Japanese Population. <i>Japanese Journal of Clinical Oncology</i> , 2012, 42, 63-73.	0.6	53
97	DNA Methylation in Peripheral Blood: A Potential Biomarker for Cancer Molecular Epidemiology. <i>Journal of Epidemiology</i> , 2012, 22, 384-394.	1.1	121
98	Treatment-Related Death in Patients with Small-Cell Lung Cancer in Phase III Trials over the Last Two Decades. <i>PLoS ONE</i> , 2012, 7, e42798.	1.1	12
99	Insulin-like growth factor 2 hypomethylation of blood leukocyte DNA is associated with gastric cancer risk. <i>International Journal of Cancer</i> , 2012, 131, 2596-2603.	2.3	27
100	Impact of smoking status on clinical outcome in oral cavity cancer patients. <i>Oral Oncology</i> , 2012, 48, 186-191.	0.8	41
101	Perceptions and Practices of Japanese Nurses Regarding Tobacco Intervention for Cancer Patients. <i>Journal of Epidemiology</i> , 2011, 21, 391-397.	1.1	11
102	Meta-analysis of neutropenia or leukopenia as a prognostic factor in patients with malignant disease undergoing chemotherapy. <i>Cancer Chemotherapy and Pharmacology</i> , 2011, 68, 301-307.	1.1	71
103	Inverse association between toothbrushing and upper aerodigestive tract cancer risk in a Japanese population. <i>Head and Neck</i> , 2011, 33, 1628-1637.	0.9	51
104	Alcohol Drinking and Esophageal Cancer Risk: An Evaluation Based on a Systematic Review of Epidemiologic Evidence Among the Japanese Population. <i>Japanese Journal of Clinical Oncology</i> , 2011, 41, 677-692.	0.6	69
105	Comparison between self-reported facial flushing after alcohol consumption and ALDH2 Glu504Lys polymorphism for risk of upper aerodigestive tract cancer in a Japanese population. <i>Cancer Science</i> , 2010, 101, 1875-1880.	1.7	68
106	Response to Yokoyama et al.: Past and current tendency for facial flushing after a small dose of alcohol is a marker for increased risk of upper aerodigestive tract cancer in Japanese drinkers. <i>Cancer Science</i> , 2010, 101, 2499-2500.	1.7	0
107	Impact of smoking on lung cancer risk is stronger in those with the homozygous aldehyde dehydrogenase 2 null allele in a Japanese population. <i>Carcinogenesis</i> , 2010, 31, 660-665.	1.3	38
108	Impact of Multiple Alcohol Dehydrogenase Gene Polymorphisms on Risk of Upper Aerodigestive Tract Cancers in a Japanese Population. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2009, 18, 3097-3102.	1.1	61

#	ARTICLE	IF	CITATIONS
109	Immunohistochemical detection of neuroendocrine differentiation in non-small-cell lung cancer and its clinical implications. <i>Journal of Cancer Research and Clinical Oncology</i> , 2009, 135, 1055-1059.	1.2	32
110	Twenty-Seven Years of Phase III Trials for Patients with Extensive Disease Small-Cell Lung Cancer: Disappointing Results. <i>PLoS ONE</i> , 2009, 4, e7835.	1.1	87