

List of Publications by Citations

Source: <https://exaly.com/author-pdf/2763211/daniel-r-witte-publications-by-citations.pdf>
Version: 2024-03-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

257 papers	16,859 citations	61 h-index	125 g-index
274 ext. papers	19,891 ext. citations	9.2 avg, IF	5.94 L-index

#	Paper	IF	Citations
257	Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. <i>Nature Genetics</i> , 2010 , 42, 937-48	36.3	2267
256	Vascular risk factors and diabetic neuropathy. <i>New England Journal of Medicine</i> , 2005 , 352, 341-50	59.2	890
255	Meta-analysis identifies 13 new loci associated with waist-hip ratio and reveals sexual dimorphism in the genetic basis of fat distribution. <i>Nature Genetics</i> , 2010 , 42, 949-60	36.3	724
254	Fine-mapping type 2 diabetes loci to single-variant resolution using high-density imputation and islet-specific epigenome maps. <i>Nature Genetics</i> , 2018 , 50, 1505-1513	36.3	675
253	Genome-wide association analysis identifies variants associated with nonalcoholic fatty liver disease that have distinct effects on metabolic traits. <i>PLoS Genetics</i> , 2011 , 7, e1001324	6	629
252	Causal relationship between obesity and vitamin D status: bi-directional Mendelian randomization analysis of multiple cohorts. <i>PLoS Medicine</i> , 2013 , 10, e1001383	11.6	592
251	Trajectories of glycaemia, insulin sensitivity, and insulin secretion before diagnosis of type 2 diabetes: an analysis from the Whitehall II study. <i>Lancet, The</i> , 2009 , 373, 2215-21	40	555
250	Rare and low-frequency coding variants alter human adult height. <i>Nature</i> , 2017 , 542, 186-190	50.4	412
249	Genome-wide association identifies nine common variants associated with fasting proinsulin levels and provides new insights into the pathophysiology of type 2 diabetes. <i>Diabetes</i> , 2011 , 60, 2624-34	0.9	285
248	Endothelial function predicts progression of carotid intima-media thickness. <i>Circulation</i> , 2009 , 119, 1005-1012	16.7	245
247	New loci associated with birth weight identify genetic links between intrauterine growth and adult height and metabolism. <i>Nature Genetics</i> , 2013 , 45, 76-82	36.3	232
246	Refining the accuracy of validated target identification through coding variant fine-mapping in type 2 diabetes. <i>Nature Genetics</i> , 2018 , 50, 559-571	36.3	221
245	A genome-wide association study in the Japanese population identifies susceptibility loci for type 2 diabetes at UBE2E2 and C2CD4A-C2CD4B. <i>Nature Genetics</i> , 2010 , 42, 864-8	36.3	214
244	A germline variant in the TP53 polyadenylation signal confers cancer susceptibility. <i>Nature Genetics</i> , 2011 , 43, 1098-103	36.3	203
243	Dietary patterns and 15-y risks of major coronary events, diabetes, and mortality. <i>American Journal of Clinical Nutrition</i> , 2008 , 87, 1414-21	7	200
242	Protein-altering variants associated with body mass index implicate pathways that control energy intake and expenditure in obesity. <i>Nature Genetics</i> , 2018 , 50, 26-41	36.3	186
241	GLP-1 Response to Oral Glucose Is Reduced in Prediabetes, Screen-Detected Type 2 Diabetes, and Obesity and Influenced by Sex: The ADDITION-PRO Study. <i>Diabetes</i> , 2015 , 64, 2513-25	0.9	171

240	Relationship between risk factors and mortality in type 1 diabetic patients in Europe: the EURODIAB Prospective Complications Study (PCS). <i>Diabetes Care</i> , 2008 , 31, 1360-6	14.6	169
239	Genome-wide association study identifies a sequence variant within the DAB2IP gene conferring susceptibility to abdominal aortic aneurysm. <i>Nature Genetics</i> , 2010 , 42, 692-7	36.3	155
238	Diabetes and cancer (1): evaluating the temporal relationship between type 2 diabetes and cancer incidence. <i>Diabetologia</i> , 2012 , 55, 1607-18	10.3	151
237	Psychosocial stress at work doubles the risk of type 2 diabetes in middle-aged women: evidence from the Whitehall II study. <i>Diabetes Care</i> , 2009 , 32, 2230-5	14.6	151
236	Elevated levels of the anti-inflammatory interleukin-1 receptor antagonist precede the onset of type 2 diabetes: the Whitehall II study. <i>Diabetes Care</i> , 2009 , 32, 421-3	14.6	150
235	Is the association between flow-mediated dilation and cardiovascular risk limited to low-risk populations?. <i>Journal of the American College of Cardiology</i> , 2005 , 45, 1987-93	15.1	146
234	Risk factors for cardiac autonomic neuropathy in type 1 diabetes mellitus. <i>Diabetologia</i> , 2005 , 48, 164-71	10.3	138
233	Exome sequencing of 20,791 cases of type 2 diabetes and 24,440 controls. <i>Nature</i> , 2019 , 570, 71-76	50.4	129
232	Moving to an A1C-based diagnosis of diabetes has a different impact on prevalence in different ethnic groups. <i>Diabetes Care</i> , 2010 , 33, 580-2	14.6	129
231	Estimation of allele frequency and association mapping using next-generation sequencing data. <i>BMC Bioinformatics</i> , 2011 , 12, 231	3.6	126
230	Cancer occurrence in Danish diabetic patients: duration and insulin effects. <i>Diabetologia</i> , 2012 , 55, 948-58	10.3	125
229	Relation between blood glucose and coronary mortality over 33 years in the Whitehall Study. <i>Diabetes Care</i> , 2006 , 29, 26-31	14.6	120
228	Agreement Between 35 Published Frailty Scores in the General Population. <i>American Journal of Epidemiology</i> , 2017 , 186, 420-434	3.8	118
227	Apolipoprotein(a) genetic sequence variants associated with systemic atherosclerosis and coronary atherosclerotic burden but not with venous thromboembolism. <i>Journal of the American College of Cardiology</i> , 2012 , 60, 722-9	15.1	118
226	Cardiovascular mortality in Dutch men during 1996 European football championship: longitudinal population study. <i>BMJ: British Medical Journal</i> , 2000 , 321, 1552-4		118
225	Inflammation, insulin resistance, and diabetes--Mendelian randomization using CRP haplotypes points upstream. <i>PLoS Medicine</i> , 2008 , 5, e155	11.6	114
224	Human pancreatic islet three-dimensional chromatin architecture provides insights into the genetics of type 2 diabetes. <i>Nature Genetics</i> , 2019 , 51, 1137-1148	36.3	111
223	Early involvement of the spinal cord in diabetic peripheral neuropathy. <i>Diabetes Care</i> , 2006 , 29, 2664-9	14.6	108

222	Natural selection affects multiple aspects of genetic variation at putatively neutral sites across the human genome. <i>PLoS Genetics</i> , 2011 , 7, e1002326	6	107
221	Work stress, obesity and the risk of type 2 diabetes: gender-specific bidirectional effect in the Whitehall II study. <i>Obesity</i> , 2012 , 20, 428-33	8	106
220	Exome sequencing-driven discovery of coding polymorphisms associated with common metabolic phenotypes. <i>Diabetologia</i> , 2013 , 56, 298-310	10.3	102
219	Genome-wide association studies in the Japanese population identify seven novel loci for type 2 diabetes. <i>Nature Communications</i> , 2016 , 7, 10531	17.4	99
218	Accelerated increase in serum interleukin-1 receptor antagonist starts 6 years before diagnosis of type 2 diabetes: Whitehall II prospective cohort study. <i>Diabetes</i> , 2010 , 59, 1222-7	0.9	98
217	Risk Factors for Incident Diabetic Polyneuropathy in a Cohort With Screen-Detected Type 2 Diabetes Followed for 13 Years: ADDITION-Denmark. <i>Diabetes Care</i> , 2018 , 41, 1068-1075	14.6	96
216	Insulin Resistance Is Accompanied by Increased Fasting Glucagon and Delayed Glucagon Suppression in Individuals With Normal and Impaired Glucose Regulation. <i>Diabetes</i> , 2016 , 65, 3473-3481	0.9	93
215	Genome-wide population-based association study of extremely overweight young adults--the GOYA study. <i>PLoS ONE</i> , 2011 , 6, e24303	3.7	90
214	Cardiovascular and all-cause mortality in relation to various anthropometric measures of obesity in Europeans. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2015 , 25, 295-304	4.5	88
213	Sex differences in glucose levels: a consequence of physiology or methodological convenience? The Inter99 study. <i>Diabetologia</i> , 2010 , 53, 858-65	10.3	83
212	Diabetes is a risk factor for pulmonary tuberculosis: a case-control study from Mwanza, Tanzania. <i>PLoS ONE</i> , 2011 , 6, e24215	3.7	81
211	Genetic evidence of a causal effect of insulin resistance on branched-chain amino acid levels. <i>Diabetologia</i> , 2017 , 60, 873-878	10.3	79
210	Role of fasting duration and weekday in incretin and glucose regulation. <i>Endocrine Connections</i> , 2021 , 10, X2-X3	3.5	78
209	Arterial stiffness, physical function, and functional limitation: the Whitehall II Study. <i>Hypertension</i> , 2011 , 57, 1003-9	8.5	76
208	Dietary glycemic index and glycemic load are associated with high-density-lipoprotein cholesterol at baseline but not with increased risk of diabetes in the Whitehall II study. <i>American Journal of Clinical Nutrition</i> , 2007 , 86, 988-94	7	76
207	Trajectories of cardiometabolic risk factors before diagnosis of three subtypes of type 2 diabetes: a post-hoc analysis of the longitudinal Whitehall II cohort study. <i>Lancet Diabetes and Endocrinology</i> , 2013 , 1, 43-51	18.1	74
206	Prevalence of neuropathy and peripheral arterial disease and the impact of treatment in people with screen-detected type 2 diabetes: the ADDITION-Denmark study. <i>Diabetes Care</i> , 2011 , 34, 2244-9	14.6	72
205	A Genome-Wide Association Study of Diabetic Kidney Disease in Subjects With Type 2 Diabetes. <i>Diabetes</i> , 2018 , 67, 1414-1427	0.9	71

204	Diabetes is a strong predictor of mortality during tuberculosis treatment: a prospective cohort study among tuberculosis patients from Mwanza, Tanzania. <i>Tropical Medicine and International Health</i> , 2013 , 18, 822-9	2.3	71
203	Low birthweight and premature birth are both associated with type 2 diabetes in a random sample of middle-aged Danes. <i>Diabetologia</i> , 2010 , 53, 2526-30	10.3	68
202	Risk of Cardiovascular Disease and Death in Individuals With Prediabetes Defined by Different Criteria: The Whitehall II Study. <i>Diabetes Care</i> , 2018 , 41, 899-906	14.6	67
201	Serum 25(OH)D and type 2 diabetes association in a general population: a prospective study. <i>Diabetes Care</i> , 2012 , 35, 1695-700	14.6	66
200	Determinants of aortic stiffness: 16-year follow-up of the Whitehall II study. <i>PLoS ONE</i> , 2012 , 7, e37165	3.7	65
199	Associations between features of glucose exposure and A1C: the A1C-Derived Average Glucose (ADAG) study. <i>Diabetes</i> , 2010 , 59, 1585-90	0.9	65
198	Serum heat shock protein 27 and diabetes complications in the EURODIAB prospective complications study: a novel circulating marker for diabetic neuropathy. <i>Diabetes</i> , 2008 , 57, 1966-70	0.9	65
197	Severe hypoglycemia and cardiovascular disease incidence in type 1 diabetes: the EURODIAB Prospective Complications Study. <i>Diabetes Care</i> , 2012 , 35, 1598-604	14.6	63
196	Dairy product intake in relation to glucose regulation indices and risk of type 2 diabetes. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2013 , 23, 822-8	4.5	61
195	Patterns of obesity development before the diagnosis of type 2 diabetes: the Whitehall II cohort study. <i>PLoS Medicine</i> , 2014 , 11, e1001602	11.6	60
194	HbA _{1c} and mean blood glucose show stronger associations with cardiovascular disease risk factors than do postprandial glycaemia or glucose variability in persons with diabetes: the A1C-Derived Average Glucose (ADAG) study. <i>Diabetologia</i> , 2011 , 54, 69-72	10.3	59
193	Prospective study of coffee and tea consumption in relation to risk of type 2 diabetes mellitus among men and women: the Whitehall II study. <i>British Journal of Nutrition</i> , 2008 , 100, 1046-53	3.6	57
192	First Genome-Wide Association Study of Latent Autoimmune Diabetes in Adults Reveals Novel Insights Linking Immune and Metabolic Diabetes. <i>Diabetes Care</i> , 2018 , 41, 2396-2403	14.6	57
191	Type 2 diabetes risk alleles near ADCY5, CDKAL1 and HHEX-IDE are associated with reduced birthweight. <i>Diabetologia</i> , 2010 , 53, 1908-16	10.3	56
190	Motivational counseling to reduce sitting time: a community-based randomized controlled trial in adults. <i>American Journal of Preventive Medicine</i> , 2014 , 47, 576-86	6.1	55
189	Associations of Mitochondrial and Nuclear Mitochondrial Variants and Genes with Seven Metabolic Traits. <i>American Journal of Human Genetics</i> , 2019 , 104, 112-138	11	54
188	Prevalence of micro- and macrovascular diabetes complications at time of type 2 diabetes diagnosis and associated clinical characteristics: A cross-sectional baseline study of 6958 patients in the Danish DD2 cohort. <i>Journal of Diabetes and Its Complications</i> , 2018 , 32, 34-40	3.2	53
187	Sleep duration and sleep quality are associated differently with alterations of glucose homeostasis. <i>Diabetic Medicine</i> , 2012 , 29, e354-60	3.5	53

186	Re-analysis of public genetic data reveals a rare X-chromosomal variant associated with type 2 diabetes. <i>Nature Communications</i> , 2018 , 9, 321	17.4	50
185	Measurement of flow-mediated dilatation of the brachial artery is affected by local elastic vessel wall properties in high-risk patients. <i>Atherosclerosis</i> , 2005 , 182, 323-30	3.1	50
184	Impact of early detection and treatment of diabetes on the 6-year prevalence of cardiac autonomic neuropathy in people with screen-detected diabetes: ADDITION-Denmark, a cluster-randomised study. <i>Diabetologia</i> , 2013 , 56, 101-8	10.3	47
183	Macrophage inhibitory cytokine-1 is increased in individuals before type 2 diabetes diagnosis but is not an independent predictor of type 2 diabetes: the Whitehall II study. <i>European Journal of Endocrinology</i> , 2010 , 162, 913-7	6.5	46
182	Studies of metabolic phenotypic correlates of 15 obesity associated gene variants. <i>PLoS ONE</i> , 2011 , 6, e23531	3.7	46
181	QTc interval prolongation is independently associated with severe hypoglycemic attacks in type 1 diabetes from the EURODIAB IDDM complications study. <i>Diabetes Care</i> , 2012 , 35, 125-7	14.6	43
180	Trajectories of glycaemia, insulin sensitivity and insulin secretion in South Asian and white individuals before diagnosis of type 2 diabetes: a longitudinal analysis from the Whitehall II cohort study. <i>Diabetologia</i> , 2017 , 60, 1252-1260	10.3	42
179	Antidepressant medication use and risk of hyperglycemia and diabetes mellitus: a noncausal association?. <i>Biological Psychiatry</i> , 2011 , 70, 978-84	7.9	42
178	Study of agreement between LDL size as measured by nuclear magnetic resonance and gradient gel electrophoresis. <i>Journal of Lipid Research</i> , 2004 , 45, 1069-76	6.3	42
177	Evidence of a liver-alpha cell axis in humans: hepatic insulin resistance attenuates relationship between fasting plasma glucagon and glucagonotropic amino acids. <i>Diabetologia</i> , 2018 , 61, 671-680	10.3	41
176	Adiponectin trajectories before type 2 diabetes diagnosis: Whitehall II study. <i>Diabetes Care</i> , 2012 , 35, 2540-7	14.6	41
175	Real-life glycaemic profiles in non-diabetic individuals with low fasting glucose and normal HbA1c: the A1C-Derived Average Glucose (ADAG) study. <i>Diabetologia</i> , 2010 , 53, 1608-11	10.3	41
174	A meta-analysis of excess cardiac mortality on Monday. <i>European Journal of Epidemiology</i> , 2005 , 20, 401-62.1	6.1	41
173	Studies of the association of Arg72Pro of tumor suppressor protein p53 with type 2 diabetes in a combined analysis of 55,521 Europeans. <i>PLoS ONE</i> , 2011 , 6, e15813	3.7	41
172	Pluralizing field analysis: Toward a relational understanding of the field of power. <i>Social Science Information</i> , 2017 , 56, 49-73	0.6	40
171	Biomarkers of subclinical inflammation and increases in glycaemia, insulin resistance and beta-cell function in non-diabetic individuals: the Whitehall II study. <i>European Journal of Endocrinology</i> , 2016 , 175, 367-77	6.5	40
170	Diabetes in Greenland and its relationship with urbanization. <i>Diabetic Medicine</i> , 2012 , 29, 755-60	3.5	40
169	Risk Factors for the Presence and Progression of Cardiovascular Autonomic Neuropathy in Type 2 Diabetes: ADDITION-Denmark. <i>Diabetes Care</i> , 2018 , 41, 2586-2594	14.6	40

168	Pathophysiological Characteristics Underlying Different Glucose Response Curves: A Latent Class Trajectory Analysis From the Prospective EGIR-RISC Study. <i>Diabetes Care</i> , 2018 , 41, 1740-1748	14.6	40
167	Antidepressant use before and after the diagnosis of type 2 diabetes: a longitudinal modeling study. <i>Diabetes Care</i> , 2010 , 33, 1471-6	14.6	39
166	Protocol for ADDITION-PRO: a longitudinal cohort study of the cardiovascular experience of individuals at high risk for diabetes recruited from Danish primary care. <i>BMC Public Health</i> , 2012 , 12, 1078	4.1	38
165	Soluble intercellular adhesion molecule 1 and flow-mediated dilatation are related to the estimated risk of coronary heart disease independently from each other. <i>Atherosclerosis</i> , 2003 , 170, 147-53	3.3	38
164	Comparative analysis of the association between 35 frailty scores and cardiovascular events, cancer, and total mortality in an elderly general population in England: An observational study. <i>PLoS Medicine</i> , 2018 , 15, e1002543	11.6	37
163	New diagnostic criteria for diabetes: is the change from glucose to HbA1c possible in all populations?. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2010 , 95, E333-6	5.6	37
162	Low peripheral nerve conduction velocities and amplitudes are strongly related to diabetic microvascular complications in type 1 diabetes: the EURODIAB Prospective Complications Study. <i>Diabetes Care</i> , 2010 , 33, 2648-53	14.6	36
161	The FOXO3A rs2802292 G-allele associates with improved peripheral and hepatic insulin sensitivity and increased skeletal muscle-FOXO3A mRNA expression in twins. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2011 , 96, E119-24	5.6	36
160	Implications of central obesity-related variants in LYPLAL1, NRXN3, MSRA, and TFAP2B on quantitative metabolic traits in adult Danes. <i>PLoS ONE</i> , 2011 , 6, e20640	3.7	36
159	Glucose-Dependent Insulinotropic Polypeptide Is Associated With Lower Low-Density Lipoprotein But Unhealthy Fat Distribution, Independent of Insulin: The ADDITION-PRO Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2016 , 101, 485-93	5.6	34
158	Type 2 diabetes risk allele near CENTD2 is associated with decreased glucose-stimulated insulin release. <i>Diabetologia</i> , 2011 , 54, 1052-6	10.3	34
157	Hyperglycemia, type 2 diabetes, and depressive symptoms: the British Whitehall II study. <i>Diabetes Care</i> , 2009 , 32, 1867-9	14.6	34
156	New Blood Pressure-Associated Loci Identified in Meta-Analyses of 475 000 Individuals. <i>Circulation: Cardiovascular Genetics</i> , 2017 , 10,		33
155	The association between n-3 fatty acids in erythrocyte membranes and insulin resistance: the Inuit Health in Transition Study. <i>International Journal of Circumpolar Health</i> , 2009 , 68, 327-36	1.7	33
154	Relationship between insulin resistance and β -cell dysfunction in subphenotypes of prediabetes and type 2 diabetes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015 , 100, 707-16	5.6	32
153	Genetic determinants of circulating interleukin-1 receptor antagonist levels and their association with glycemic traits. <i>Diabetes</i> , 2014 , 63, 4343-59	0.9	32
152	Socioeconomic status and reduced kidney function in the Whitehall II Study: role of obesity and metabolic syndrome. <i>American Journal of Kidney Diseases</i> , 2011 , 58, 389-97	7.4	32
151	Association studies of novel obesity-related gene variants with quantitative metabolic phenotypes in a population-based sample of 6,039 Danish individuals. <i>Diabetologia</i> , 2012 , 55, 105-13	10.3	31

150	The frequent UCP2 -866G>A polymorphism protects against insulin resistance and is associated with obesity: a study of obesity and related metabolic traits among 17 636 Danes. <i>International Journal of Obesity</i> , 2013 , 37, 175-81	5.5	31
149	and Loci Identified through Large-Scale Exome Chip Analysis Regulate Kidney Development and Function. <i>Journal of the American Society of Nephrology: JASN</i> , 2017 , 28, 981-994	12.7	30
148	National study of the prevalence of gestational diabetes mellitus among Danish women from 2004 to 2012. <i>Scandinavian Journal of Public Health</i> , 2017 , 45, 811-817	3	30
147	The role of serum methylglyoxal on diabetic peripheral and cardiovascular autonomic neuropathy: the ADDITION Denmark study. <i>Diabetic Medicine</i> , 2015 , 32, 778-85	3.5	30
146	Combined heart rate- and accelerometer-assessed physical activity energy expenditure and associations with glucose homeostasis markers in a population at high risk of developing diabetes: the ADDITION-PRO study. <i>Diabetes Care</i> , 2013 , 36, 3062-9	14.6	30
145	A standardized vascular disease health check in europe: a cost-effectiveness analysis. <i>PLoS ONE</i> , 2013 , 8, e66454	3.7	30
144	Glucose patterns during an oral glucose tolerance test and associations with future diabetes, cardiovascular disease and all-cause mortality rate. <i>Diabetologia</i> , 2018 , 61, 101-107	10.3	29
143	Methylglyoxal is associated with changes in kidney function among individuals with screen-detected Type 2 diabetes mellitus. <i>Diabetic Medicine</i> , 2016 , 33, 1625-1631	3.5	29
142	Low serum adiponectin predicts 10-year risk of type 2 diabetes and HbA1c independently of obesity, lipids, and inflammation: Whitehall II study. <i>Hormone and Metabolic Research</i> , 2009 , 41, 626-9	3.1	29
141	The pro-inflammatory biomarker soluble urokinase plasminogen activator receptor (suPAR) is associated with incident type 2 diabetes among overweight but not obese individuals with impaired glucose regulation: effect modification by smoking and body weight status. <i>Diabetologia</i> , 2013 , 56, 1542-6	10.3	28
140	Effect of population screening for type 2 diabetes and cardiovascular risk factors on mortality rate and cardiovascular events: a controlled trial among 1,912,392 Danish adults. <i>Diabetologia</i> , 2017 , 60, 2183-2191	10.3	28
139	Comparing risk profiles of individuals diagnosed with diabetes by OGTT and HbA1c The Danish Inter99 study. <i>Diabetic Medicine</i> , 2010 , 27, 906-10	3.5	28
138	Adiponectin, biomarkers of inflammation and changes in cardiac autonomic function: Whitehall II study. <i>Cardiovascular Diabetology</i> , 2017 , 16, 153	8.7	27
137	ANTI-HSP60 and ANTI-HSP70 antibody levels and micro/ macrovascular complications in type 1 diabetes: the EURODIAB Study. <i>Journal of Internal Medicine</i> , 2009 , 266, 527-36	10.8	27
136	The association between father's social class and adult obesity is not explained by educational attainment and an unhealthy lifestyle in adulthood. <i>European Journal of Epidemiology</i> , 2008 , 23, 573-9	12.1	27
135	Nondiabetic Glucometabolic Status and Progression of Aortic Stiffness: The Whitehall II Study. <i>Diabetes Care</i> , 2017 , 40, 599-606	14.6	26
134	Reversion from prediabetes to normoglycaemia and risk of cardiovascular disease and mortality: the Whitehall II cohort study. <i>Diabetologia</i> , 2019 , 62, 1385-1390	10.3	26
133	Psychological distress, cardiovascular complications and mortality among people with screen-detected type 2 diabetes: follow-up of the ADDITION-Denmark trial. <i>Diabetologia</i> , 2014 , 57, 710-7	10.3	26

132	The diabetogenic VPS13C/C2CD4A/C2CD4B rs7172432 variant impairs glucose-stimulated insulin response in 5,722 non-diabetic Danish individuals. <i>Diabetologia</i> , 2011 , 54, 789-94	10.3	26
131	Performance of existing risk scores in screening for undiagnosed diabetes: an external validation study. <i>Diabetic Medicine</i> , 2010 , 27, 46-53	3.5	26
130	Changes in C-reactive protein levels before type 2 diabetes and cardiovascular death: the Whitehall II study. <i>European Journal of Endocrinology</i> , 2010 , 163, 89-95	6.5	26
129	Discovery of rare variants associated with blood pressure regulation through meta-analysis of 1.3 million individuals. <i>Nature Genetics</i> , 2020 , 52, 1314-1332	36.3	26
128	The PNPLA3 rs738409 G-allele associates with reduced fasting serum triglyceride and serum cholesterol in Danes with impaired glucose regulation. <i>PLoS ONE</i> , 2012 , 7, e40376	3.7	25
127	Predictors of future fasting and 2-h post-OGTT plasma glucose levels in middle-aged men and women-the Inter99 study. <i>Diabetic Medicine</i> , 2009 , 26, 377-83	3.5	25
126	Abdominal Fat Distribution and Cardiovascular Risk in Men and Women With Different Levels of Glucose Tolerance. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015 , 100, 3340-7	5.6	24
125	Differential relationship between physical activity and progression to diabetes by glucose tolerance status: the Inter99 Study. <i>Diabetologia</i> , 2010 , 53, 70-8	10.3	24
124	Vascular risk factors and diabetic neuropathy. <i>New England Journal of Medicine</i> , 2005 , 352, 1925-7; author reply 1925-7	59.2	24
123	The role of diabetes co-morbidity for tuberculosis treatment outcomes: a prospective cohort study from Mwanza, Tanzania. <i>BMC Infectious Diseases</i> , 2012 , 12, 165	4	22
122	Associations between ultrasound measures of abdominal fat distribution and indices of glucose metabolism in a population at high risk of type 2 diabetes: the ADDITION-PRO study. <i>PLoS ONE</i> , 2015 , 10, e0123062	3.7	22
121	No interactions between previously associated 2-hour glucose gene variants and physical activity or BMI on 2-hour glucose levels. <i>Diabetes</i> , 2012 , 61, 1291-6	0.9	21
120	Risk-Factor Trajectories Preceding Diabetic Polyneuropathy: ADDITION-Denmark. <i>Diabetes Care</i> , 2018 , 41, 1955-1962	14.6	20
119	Effect of secular trends on age-related trajectories of cardiovascular risk factors: the Whitehall II longitudinal study 1985-2009. <i>International Journal of Epidemiology</i> , 2014 , 43, 866-77	7.8	20
118	The effect of PCSK1 variants on waist, waist-hip ratio and glucose metabolism is modified by sex and glucose tolerance status. <i>PLoS ONE</i> , 2011 , 6, e23907	3.7	20
117	The role of anthropometric and other predictors for diabetes among urban Tanzanians with tuberculosis. <i>International Journal of Tuberculosis and Lung Disease</i> , 2012 , 16, 1680-5	2.1	20
116	Do gene variants influencing adult adiposity affect birth weight? A population-based study of 24 loci in 4,744 Danish individuals. <i>PLoS ONE</i> , 2010 , 5, e14190	3.7	20
115	Prospective Association Among Diabetes Diagnosis, HbA _{1c} , Glycemia, and Frailty Trajectories in an Elderly Population. <i>Diabetes Care</i> , 2019 , 42, 1903-1911	14.6	19

114	Bioinformatics-driven identification and examination of candidate genes for non-alcoholic fatty liver disease. <i>PLoS ONE</i> , 2011 , 6, e16542	3.7	19
113	Effect of intensive multifactorial treatment compared with routine care on aortic stiffness and central blood pressure among individuals with screen-detected type 2 diabetes: the ADDITION-Denmark study. <i>Diabetes Care</i> , 2012 , 35, 2207-14	14.6	19
112	The role of diabetes on the clinical manifestations of pulmonary tuberculosis. <i>Tropical Medicine and International Health</i> , 2012 , 17, 877-83	2.3	18
111	Heterogeneity in glucose response curves during an oral glucose tolerance test and associated cardiometabolic risk. <i>Endocrine</i> , 2017 , 55, 427-434	4	18
110	Solute carrier family 2 member 1 is involved in the development of nonalcoholic fatty liver disease. <i>Hepatology</i> , 2013 , 57, 505-14	11.2	18
109	Long-term effects of intensive multifactorial therapy in individuals with screen-detected type 2 diabetes in primary care: 10-year follow-up of the ADDITION-Europe cluster-randomised trial. <i>Lancet Diabetes and Endocrinology</i> , 2019 , 7, 925-937	18.1	18
108	Incremental predictive value of adding past blood pressure measurements to the Framingham hypertension risk equation: the Whitehall II Study. <i>Hypertension</i> , 2010 , 55, 1058-62	8.5	17
107	Effect of duration and burden of microvascular complications on mortality rate in type 1 diabetes: an observational clinical cohort study. <i>Diabetologia</i> , 2019 , 62, 633-643	10.3	17
106	Effectiveness of Liraglutide and Lixisenatide in the Treatment of Type 2 Diabetes: Real-World Evidence from The Health Improvement Network (THIN) Database in the United Kingdom. <i>Diabetes Therapy</i> , 2017 , 8, 417-431	3.6	16
105	Sex differences in glucose and insulin trajectories prior to diabetes diagnosis: the Whitehall II study. <i>Acta Diabetologica</i> , 2014 , 51, 315-9	3.9	16
104	NH2-terminal probrain natriuretic peptide is associated with diabetes complications in the EURODIAB Prospective Complications Study: the role of tumor necrosis factor- α . <i>Diabetes Care</i> , 2012 , 35, 1931-6	14.6	16
103	HbA1c, fasting plasma glucose and the prediction of diabetes: Inter99, AusDiab and D.E.S.I.R. <i>Diabetes Research and Clinical Practice</i> , 2012 , 96, 392-9	7.4	15
102	Reproducibility of ultrasonography for assessing abdominal fat distribution in a population at high risk of diabetes. <i>Nutrition and Diabetes</i> , 2013 , 3, e82	4.7	15
101	Clustering of microvascular complications in Type 1 diabetes mellitus. <i>Journal of Diabetes and Its Complications</i> , 2018 , 32, 393-399	3.2	14
100	Soluble CD163, adiponectin, C-reactive protein and progression of dysglycaemia in individuals at high risk of type 2 diabetes mellitus: the ADDITION-PRO cohort. <i>Diabetologia</i> , 2016 , 59, 2467-2476	10.3	14
99	Homozygous carriers of the G allele of rs4664447 of the glucagon gene (GCG) are characterised by decreased fasting and stimulated levels of insulin, glucagon and glucagon-like peptide (GLP)-1. <i>Diabetologia</i> , 2011 , 54, 2820-31	10.3	14
98	Anti-heat shock protein 27 antibody levels and diabetes complications in the EURODIAB study. <i>Diabetes Care</i> , 2009 , 32, 1269-71	14.6	14
97	Associations of common genetic variants with age-related changes in fasting and postload glucose: evidence from 18 years of follow-up of the Whitehall II cohort. <i>Diabetes</i> , 2011 , 60, 1617-23	0.9	14

96	Functional and genetic epidemiological characterisation of the FFAR4 (GPR120) p.R270H variant in the Danish population. <i>Journal of Medical Genetics</i> , 2016 , 53, 616-23	5.8	14
95	Glucose patterns during the OGTT and risk of future diabetes in an urban Indian population: The CARRS study. <i>Diabetes Research and Clinical Practice</i> , 2017 , 126, 192-197	7.4	13
94	Risk scores for diabetes and impaired glycaemia in the Middle East and North Africa. <i>Diabetic Medicine</i> , 2013 , 30, 443-51	3.5	13
93	Studies of association between LPIN1 variants and common metabolic phenotypes among 17,538 Danes. <i>European Journal of Endocrinology</i> , 2010 , 163, 81-7	6.5	13
92	Excess cardiac mortality on Monday: the importance of gender, age and hospitalisation. <i>European Journal of Epidemiology</i> , 2005 , 20, 395-9	12.1	13
91	Social relations, depressive symptoms, and incident type 2 diabetes mellitus: The English Longitudinal Study of Ageing. <i>Diabetes Research and Clinical Practice</i> , 2017 , 126, 86-94	7.4	12
90	Spousal cardiometabolic risk factors and incidence of type 2 diabetes: a prospective analysis from the English Longitudinal Study of Ageing. <i>Diabetologia</i> , 2018 , 61, 1572-1580	10.3	12
89	BCG protects against tuberculosis irrespective of HIV status: a matched case-control study in Mwanza, Tanzania. <i>Thorax</i> , 2013 , 68, 288-9	7.3	12
88	The birth weight lowering C-allele of rs900400 near LEKR1 and CCNL1 associates with elevated insulin release following an oral glucose challenge. <i>PLoS ONE</i> , 2011 , 6, e27096	3.7	12
87	The role of physical activity in the development of first cardiovascular disease event: a tree-structured survival analysis of the Danish ADDITION-PRO cohort. <i>Cardiovascular Diabetology</i> , 2018 , 17, 126	8.7	12
86	Associations of objectively measured physical activity and abdominal fat distribution. <i>Medicine and Science in Sports and Exercise</i> , 2015 , 47, 983-9	1.2	11
85	Higher physical activity is associated with lower aortic stiffness but not with central blood pressure: the ADDITION-Pro Study. <i>Medicine (United States)</i> , 2015 , 94, e485	1.8	11
84	Engineering of strained III/V heterostructures for high hole mobility 2009 ,		11
83	Physical activity energy expenditure vs cardiorespiratory fitness level in impaired glucose metabolism. <i>Diabetologia</i> , 2015 , 58, 2709-17	10.3	10
82	Development of Microvascular Complications and Effect of Concurrent Risk Factors in Type 1 Diabetes: A Multistate Model From an Observational Clinical Cohort Study. <i>Diabetes Care</i> , 2018 , 41, 2297-2305	14.6	10
81	Sex-specific effects of naturally occurring variants in the dopamine receptor D2 locus on insulin secretion and type 2 diabetes susceptibility. <i>Diabetic Medicine</i> , 2014 , 31, 1001-8	3.5	10
80	Effect of time of day and fasting duration on measures of glycaemia: analysis from the Whitehall II Study. <i>Diabetologia</i> , 2013 , 56, 294-7	10.3	10
79	Physical Activity and Improvement of Glycemia in Prediabetes by Different Diagnostic Criteria. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2017 , 102, 3712-3721	5.6	10

78	Reduction of specific circulating lymphocyte populations with metabolic risk factors in patients at risk to develop type 2 diabetes. <i>PLoS ONE</i> , 2014 , 9, e107140	3.7	10
77	The effect of FOXA2 rs1209523 on glucose-related phenotypes and risk of type 2 diabetes in Danish individuals. <i>BMC Medical Genetics</i> , 2012 , 13, 10	2.1	10
76	The minor C-allele of rs2014355 in ACADS is associated with reduced insulin release following an oral glucose load. <i>BMC Medical Genetics</i> , 2011 , 12, 4	2.1	10
75	Trans-ancestry genetic study of type 2 diabetes highlights the power of diverse populations for discovery and translation		10
74	Incidence of register-based diabetes 10 years after a stepwise diabetes screening programme: the ADDITION-Denmark study. <i>Diabetologia</i> , 2016 , 59, 989-97	10.3	10
73	Invasively Measured Aortic Systolic Blood Pressure and Office Systolic Blood Pressure in Cardiovascular Risk Assessment: A Prospective Cohort Study. <i>Hypertension</i> , 2016 , 68, 768-74	8.5	10
72	Prevalence and geographical distribution of insulin pump therapy in the Central Denmark Region and its association with metabolic parameters. <i>Diabetes Research and Clinical Practice</i> , 2018 , 141, 148-155	7.4	10
71	A combined analysis of 48 type 2 diabetes genetic risk variants shows no discriminative value to predict time to first prescription of a glucose lowering drug in Danish patients with screen detected type 2 diabetes. <i>PLoS ONE</i> , 2014 , 9, e104837	3.7	9
70	Heart Rate, Autonomic Function, and Future Changes in Glucose Metabolism in Individuals Without Diabetes: The Whitehall II Cohort Study. <i>Diabetes Care</i> , 2019 , 42, 867-874	14.6	8
69	Association of self-perceived body image with body mass index and type 2 diabetes-The ADDITION-PRO study. <i>Preventive Medicine</i> , 2015 , 75, 64-9	4.3	8
68	Impact of intensive treatment on serum methylglyoxal levels among individuals with screen-detected type 2 diabetes: the ADDITION-Denmark study. <i>Acta Diabetologica</i> , 2015 , 52, 929-36	3.9	8
67	Differences Between Randomized Clinical Trial Participants and Real-World Empagliflozin Users and the Changes in Their Glycated Hemoglobin Levels. <i>JAMA Network Open</i> , 2020 , 3, e1920949	10.4	8
66	Common variants in the hERG (KCNH2) voltage-gated potassium channel are associated with altered fasting and glucose-stimulated plasma incretin and glucagon responses. <i>BMC Genetics</i> , 2018 , 19, 15	2.6	8
65	Genetic Correlation between Body Fat Percentage and Cardiorespiratory Fitness Suggests Common Genetic Etiology. <i>PLoS ONE</i> , 2016 , 11, e0166738	3.7	8
64	Prospective Study of Neuropathic Symptoms Preceding Clinically Diagnosed Diabetic Polyneuropathy: ADDITION-Denmark. <i>Diabetes Care</i> , 2019 , 42, 2282-2289	14.6	7
63	Glucose Measurements at Various Time Points During the OGTT and Their Role in Capturing Glucose Response Patterns. <i>Diabetes Care</i> , 2019 , 42, e56-e57	14.6	7
62	Glycaemic threshold for diabetes-specific retinopathy among individuals from Saudi Arabia, Algeria and Portugal. <i>Diabetes Research and Clinical Practice</i> , 2014 , 103, e44-6	7.4	7
61	Human pancreatic islet 3D chromatin architecture provides insights into the genetics of type 2 diabetes		7

60	Multi-ancestry genetic study of type 2 diabetes highlights the power of diverse populations for discovery and translation.. <i>Nature Genetics</i> , 2022 ,	36.3	7
59	Association between protein signals and type 2 diabetes incidence. <i>Acta Diabetologica</i> , 2013 , 50, 697-704.	5.9	6
58	Physical Activity Dimensions Associated with Impaired Glucose Metabolism. <i>Medicine and Science in Sports and Exercise</i> , 2017 , 49, 2176-2184	1.2	6
57	Household and familial resemblance in risk factors for type 2 diabetes and related cardiometabolic diseases in rural Uganda: a cross-sectional community sample. <i>BMJ Open</i> , 2017 , 7, e015214	3	6
56	Do the Joint British Society (JBS2) guidelines on prevention of cardiovascular disease with respect to plasma glucose improve risk stratification in the general population? Prospective cohort study. <i>Diabetic Medicine</i> , 2010 , 27, 550-5	3.5	6
55	Netzwerke als transversale Felder 2019 , 25-61		6
54	Plasma lipid metabolites associate with diabetic polyneuropathy in a cohort with type 2 diabetes. <i>Annals of Clinical and Translational Neurology</i> , 2021 , 8, 1292-1307	5.3	6
53	Does training of general practitioners for intensive treatment of people with screen-detected diabetes have a spillover effect on mortality and cardiovascular morbidity in at risk individuals with normoglycaemia? Results from the ADDITION-Denmark cluster-randomised controlled trial. <i>Diabetologia</i> , 2017 , 60, 1016-1021	10.3	5
52	Glycemic index and glycemic load in relation to glucose intolerance among Greenland Inuit population. <i>Diabetes Research and Clinical Practice</i> , 2012 , 97, 298-305	7.4	5
51	Determinants of penetrance and variable expressivity in monogenic metabolic conditions across 77,184 exomes. <i>Nature Communications</i> , 2021 , 12, 3505	17.4	5
50	Validity of Danish register diagnoses of myocardial infarction and stroke against experts in people with screen-detected diabetes. <i>BMC Public Health</i> , 2019 , 19, 228	4.1	4
49	Heterogeneous effect of gestational weight gain on birth weight: quantile regression analysis from a population-based screening. <i>Annals of Epidemiology</i> , 2015 , 25, 133-7, 137.e1	6.4	4
48	Studies of a genetic variant in HK1 in relation to quantitative metabolic traits and to the prevalence of type 2 diabetes. <i>BMC Medical Genetics</i> , 2011 , 12, 99	2.1	4
47	Das Feld der Macht 2020 ,		4
46	Identifying hotspots of cardiometabolic outcomes based on a Bayesian approach: The example of Chile. <i>PLoS ONE</i> , 2020 , 15, e0235009	3.7	3
45	Geordnete Verhältnisse? Vielfalt und Einheit relationalen Denkens in der Soziologie. <i>Berliner Journal Für Soziologie</i> , 2017 , 27, 347-376	0.4	3
44	Association of weight loss and weight loss maintenance following diabetes diagnosis by screening and incidence of cardiovascular disease and all-cause mortality: An observational analysis of the ADDITION-Europe trial. <i>Diabetes, Obesity and Metabolism</i> , 2021 , 23, 730-741	6.7	3
43	Associations between glycaemic deterioration and aortic stiffness and central blood pressure: the ADDITION-PRO Study. <i>Journal of Hypertension</i> , 2017 , 35, 1832-1840	1.9	2

42	Effect of familial diabetes status and age at diagnosis on type 2 diabetes risk: a nation-wide register-based study from Denmark. <i>Diabetologia</i> , 2020 , 63, 934-943	10.3	2
41	Studies of association of AGPAT6 variants with type 2 diabetes and related metabolic phenotypes in 12,068 Danes. <i>BMC Medical Genetics</i> , 2013 , 14, 113	2.1	2
40	Abdominal obesity has the highest impact on metabolic profile in an overweight African population. <i>Annals of Human Biology</i> , 2012 , 39, 530-3	1.7	2
39	Role of fasting duration and weekday in incretin and glucose regulation. <i>Endocrine Connections</i> , 2020 , 9, 279-288	3.5	2
38	Obesity and Kidney Function: A Two-Sample Mendelian Randomization Study.. <i>Clinical Chemistry</i> , 2021 ,	5.5	2
37	Der Staat und die gelehrigen Körper. <i>Politologische Aufklärung - Konstruktivistische Perspektiven</i> , 2019 , 211-233	0.3	2
36	Rare coding variants in 35 genes associate with circulating lipid levels in a multi-ancestry analysis of 170,000 exomes		2
35	Genetic discovery and translational decision support from exome sequencing of 20,791 type 2 diabetes cases and 24,440 controls from five ancestries		2
34	Trajectories of obesity by spousal diabetes status in the English Longitudinal Study of Ageing. <i>Diabetic Medicine</i> , 2019 , 36, 105-109	3.5	2
33	Ein ambivalentes Erbe: Von Durkheims (un-)politischer Soziologie zu Bourdieus Religionssoziologie des Staates. <i>Berliner Journal Fur Soziologie</i> , 2018 , 28, 307-337	0.4	2
32	Prospective association between late evening food consumption and risk of prediabetes and diabetes: the Whitehall II cohort study. <i>Diabetic Medicine</i> , 2019 , 36, 1256-1260	3.5	1
31	Genetic determinants of blood pressure traits are associated with carotid arterial thickening and plaque formation in patients with type 2 diabetes. <i>Diabetes and Vascular Disease Research</i> , 2019 , 16, 13-21	3.3	1
30	Response to Comment on Fitch et al. GLP-1 Response to Oral Glucose Is Reduced in Prediabetes, Screen-Detected Type 2 Diabetes, and Obesity and Influenced by Sex: The ADDITION-PRO Study. <i>Diabetes</i> 2015;64:2513-2525. <i>Diabetes</i> , 2015 , 64, e30-e31	0.9	1
29	Assessment of time to glucose peak during an oral glucose tolerance test. <i>Clinical Endocrinology</i> , 2017 , 87, 879-881	3.4	1
28	Reply to K Esposito and D Giugliano. <i>American Journal of Clinical Nutrition</i> , 2008 , 88, 1180-1181	7	1
27	Does Training and Support of General Practitioners in Intensive Treatment of People with Screen-Detected Diabetes Improve Medication, Morbidity and Mortality in People with Clinically-Diagnosed Diabetes? Investigation of a Spill-Over Effect in a Cluster RCT. <i>PLoS ONE</i> , 2017 , 12, e0170697	3.7	1
26	Habitual physical activity is associated with lower fasting and greater glucose-induced GLP-1 response in men. <i>Endocrine Connections</i> , 2019 , 8, 1607-1617	3.5	1
25	Das Feld der Macht in der Soziologie Bourdieus 2020 , 15-35		1

24	Changes in type 2 diabetes incidence and mortality associated with introduction of HbA1c as diagnostic option: A Danish 24-year population-based study.. <i>Lancet Regional Health - Europe, The</i> , 2022 , 14, 100291		1
23	The effect of training GPs in motivational interviewing on incident cardiovascular disease and mortality in people with screen-detected diabetes. Results from the ADDITION-Denmark randomised trial. <i>BJGP Open</i> , 2020 , 4,	3.1	1
22	Determinants of penetrance and variable expressivity in monogenic metabolic conditions across 77,184 exomes		1
21	Factors associated with attendance at clinical follow-up of a cohort with screen-detected type 2 diabetes: ADDITION-Denmark. <i>Primary Care Diabetes</i> , 2020 , 14, 239-245	2.4	1
20	Potentially inappropriate medications (PIMs): frequency and extent of GP-related variation in PIMs: a register-based cohort study. <i>BMJ Open</i> , 2021 , 11, e046756	3	1
19	Heart Rate and Heart Rate Variability Changes Are Not Related to Future Cardiovascular Disease and Death in People With and Without Dysglycemia: A Downfall of Risk Markers? The Whitehall II Cohort Study. <i>Diabetes Care</i> , 2021 , 44, 1012-1019	14.6	1
18	Greater glucagon-like peptide-1 responses to oral glucose are associated with lower central and peripheral blood pressures. <i>Cardiovascular Diabetology</i> , 2019 , 18, 130	8.7	0
17	Die zwei Gesichter der Autonomie 2017 , 383-423		0
16	Towards precision medicine in diabetes? A critical review of glucotypes. <i>PLoS Biology</i> , 2021 , 19, e3000890	7	0
15	Diabetic Polyneuropathy Early in Type 2 Diabetes Is Associated With Higher Incidence Rate of Cardiovascular Disease: Results From Two Danish Cohort Studies. <i>Diabetes Care</i> , 2021 , 44, 1714-1721	14.6	0
14	Objective and subjective sleep measures are associated with HbA1c and insulin sensitivity in the general population: Findings from the ORISCAV-LUX-2 study. <i>Diabetes and Metabolism</i> , 2021 , 48, 101263	5.4	0
13	Duration of diabetes-related complications and mortality in type 1 diabetes: a national cohort study. <i>International Journal of Epidemiology</i> , 2021 , 50, 1250-1259	7.8	0
12	Prospects for a Relational Sociology of Islam: Some Remarks on Differentiation Theory, Multiple Modernities, and the Pitfalls of Occidentalism 2021 , 65-88		0
11	Genome-wide association study of circulating levels of glucagon during an oral glucose tolerance test. <i>BMC Medical Genomics</i> , 2021 , 14, 3	3.7	0
10	The Arg82Cys Polymorphism of the Protein Nepmucin Implies a Role in HDL Metabolism.. <i>Journal of the Endocrine Society</i> , 2022 , 6, bvac034	0.4	0
9	Health-related quality of life for normal glycaemia, prediabetes and type 2 diabetes mellitus: Cross-sectional analysis of the ADDITION-PRO study.. <i>Diabetic Medicine</i> , 2022 , e14825	3.5	0
8	Letter to the Editor: "One-Hour Postload Hyperglycemia: Implications for Prediction and Prevention of Type 2 Diabetes". <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019 , 104, 674-675	5.6	
7	Can insulin response patterns predict metabolic disease risk in individuals with normal glucose tolerance? Reply to Crofts CAP, Brookler K, Henderson G [letter]. <i>Diabetologia</i> , 2018 , 61, 1234-1235	10.3	

6 Vom nationalen zum globalen Feld der Macht **2020**, 103-152

5 Zur Pluralisierung der Feldanalyse: Das Feld der Macht als Feld der Felder **2020**, 61-102

4 r-cubed: Guiding the overwhelmed scientist from random wrangling to Reproducible Research in R. *The Journal of Open Source Education*, **2021**, 4, 122 1.2

3 Das Feld der Macht als gesellschaftstheoretisches Schlüsselkonzept **2020**, 1-13

2 Response to the Letter: Comment on "Abdominal Fat Distribution and Cardiovascular Risk in Men and Women With Different Levels of Glucose Tolerance" by Scheuer S.H., et al. *Journal of Clinical Endocrinology and Metabolism*, **2016**, 101, L13-4 5.6

1 Response to Comment on Andersen et al. Risk-Factor Trajectories Preceding Diabetic Polyneuropathy: ADDITION-Denmark. *Diabetes Care* 2018;41:1955-1962. *Diabetes Care*, **2018**, 41, e148-e149 14.6