

Mohammad Hassan Safaralizadeh

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2775559/publications.pdf>

Version: 2024-02-01

10

papers

39

citations

1937685

4

h-index

1872680

6

g-index

10

all docs

10

docs citations

10

times ranked

52

citing authors

#	ARTICLE	IF	CITATIONS
1	Synergy between parasitoids and pathogens for biological control of <i>Helicoverpa armigera</i> in chickpea. <i>Entomologia Experimentalis Et Applicata</i> , 2020, 168, 70-75.	1.4	8
2	Behavioral and Developmental Responses of <i>Habrobracon hebetor</i> (Hymenoptera: Braconidae) to Larvae of <i>Helicoverpa armigera</i> (Lepidoptera: Noctuidae) Inoculated With Various Concentrations of <i>Bacillus thuringiensis</i> var. <i>kurstaki</i> (Bacillales: Bacillaceae). <i>Journal of Insect Science</i> , 2020, 20, .	1.5	3
3	Negative Life History Impacts for <i>Habrobracon hebetor</i> (Hymenoptera: Braconidae) that Develop in Bollworm Larvae Inoculated with <i>Helicoverpa armigera</i> Nucleopolyhedrovirus. <i>Journal of Economic Entomology</i> , 2020, 113, 1648-1655.	1.8	3
4	Effect of fungal isolates and imidacloprid on cabbage aphid <i>Brevicoryne brassicae</i> and its parasitoid <i>Diaeretiella rapae</i> . <i>Zemdirbyste</i> , 2020, 107, 255-262.	0.8	2
5	Faunistic study of Thysanoptera in West Azerbaijan Province, north-western Iran. <i>Zoology and Ecology</i> , 2018, 28, 155-162.	0.2	0
6	<p>New species of <i>Hexabdella</i> and <i>Neomolgus</i> (Acari: Prostigmata: Bdellidae)from Iran</p>. <i>Zootaxa</i> , 2016, 4072, 291.	0.5	4
7	Effects of <i>Bacillus thuringiensis</i> var. <i>kurstaki</i> and Spinosad on three larval stages 1st, 2nd and 3rd of tomato borer, <i>Tuta absoluta</i> (Meyrick) (Lepidoptera: Gelechiidae) in laboratory conditions. <i>Archives of Phytopathology and Plant Protection</i> , 2015, 48, 377-384.	1.3	4
8	Two new species of <i>Cyta</i> (Acari: Prostigmata: Bdellidae) from Western Iran. <i>Zootaxa</i> , 2014, 3847, 567-75.	0.5	5
9	Virulence of some native <i>Bacillus thuringiensis</i> isolates against <i>Ephestia kuehniella</i> (Zeller) (Lep., Pyralidae) and <i>Pieris brassicae</i> (Lep., Pieridae) larvae isolated from stored products of Urmia city. <i>Archives of Phytopathology and Plant Protection</i> , 2014, 47, 610-614.	1.3	0
10	Fumigant toxicity of <i>Laurus nobilis</i> and <i>Myrtus communis</i> essential oils on larvae and adults of the Red flour beetle, <i>Tribolium castaneum</i> Herbst (Col.: Tenebrionidae). <i>Archives of Phytopathology and Plant Protection</i> , 2014, 47, 472-476.	1.3	10