

Christopher Conrad

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2781478/publications.pdf>

Version: 2024-02-01

98
papers

3,258
citations

159358

30
h-index

161609

54
g-index

100
all docs

100
docs citations

100
times ranked

3702
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Impact of feature selection on the accuracy and spatial uncertainty of per-field crop classification using Support Vector Machines. <i>ISPRS Journal of Photogrammetry and Remote Sensing</i> , 2013, 85, 102-119. | 4.9 | 219 |
| 2 | Per-Field Irrigated Crop Classification in Arid Central Asia Using SPOT and ASTER Data. <i>Remote Sensing</i> , 2010, 2, 1035-1056. | 1.8 | 143 |
| 3 | Integration of Optical and Synthetic Aperture Radar Imagery for Improving Crop Mapping in Northwestern Benin, West Africa. <i>Remote Sensing</i> , 2014, 6, 6472-6499. | 1.8 | 139 |
| 4 | Crop type classification using a combination of optical and radar remote sensing data: a review. <i>International Journal of Remote Sensing</i> , 2019, 40, 6553-6595. | 1.3 | 126 |
| 5 | Grassland habitat mapping by intra-annual time series analysis – Comparison of RapidEye and TerraSAR-X satellite data. <i>International Journal of Applied Earth Observation and Geoinformation</i> , 2015, 34, 25-34. | 1.4 | 114 |
| 6 | Quantifying and Mapping Ecosystem Services Supplies and Demands: A Review of Remote Sensing Applications. <i>Environmental Science & Technology</i> , 2012, 46, 8529-8541. | 4.6 | 112 |
| 7 | Estimating the fractional cover of growth forms and bare surface in savannas. A multi-resolution approach based on regression tree ensembles. <i>Remote Sensing of Environment</i> , 2013, 129, 90-102. | 4.6 | 105 |
| 8 | Large-area assessment of impervious surface based on integrated analysis of single-date Landsat-7 images and geospatial vector data. <i>Remote Sensing of Environment</i> , 2009, 113, 1678-1690. | 4.6 | 102 |
| 9 | Mapping and assessing water use in a Central Asian irrigation system by utilizing MODIS remote sensing products. <i>Irrigation and Drainage Systems</i> , 2007, 21, 197-218. | 0.5 | 99 |
| 10 | Spatio-temporal analyses of cropland degradation in the irrigated lowlands of Uzbekistan using remote-sensing and logistic regression modeling. <i>Environmental Monitoring and Assessment</i> , 2013, 185, 4775-4790. | 1.3 | 95 |
| 11 | Temporal segmentation of MODIS time series for improving crop classification in Central Asian irrigation systems. <i>International Journal of Remote Sensing</i> , 2011, 32, 8763-8778. | 1.3 | 88 |
| 12 | MODIS NDVI trends and fractional land cover change for improved assessments of vegetation degradation in Burkina Faso, West Africa. <i>Journal of Arid Environments</i> , 2018, 153, 66-75. | 1.2 | 82 |
| 13 | TiSeG: A Flexible Software Tool for Time-Series Generation of MODIS Data Utilizing the Quality Assessment Science Data Set. <i>IEEE Transactions on Geoscience and Remote Sensing</i> , 2008, 46, 3296-3308. | 2.7 | 79 |
| 14 | Derivation of temporal windows for accurate crop discrimination in heterogeneous croplands of Uzbekistan using multitemporal RapidEye images. <i>Computers and Electronics in Agriculture</i> , 2014, 103, 63-74. | 3.7 | 76 |
| 15 | Mapping abandoned agricultural land in Kyzyl-Orda, Kazakhstan using satellite remote sensing. <i>Applied Geography</i> , 2015, 62, 377-390. | 1.7 | 72 |
| 16 | Measuring rural settlement expansion in Uzbekistan using remote sensing to support spatial planning. <i>Applied Geography</i> , 2015, 62, 29-43. | 1.7 | 69 |
| 17 | Decision fusion and non-parametric classifiers for land use mapping using multi-temporal RapidEye data. <i>ISPRS Journal of Photogrammetry and Remote Sensing</i> , 2015, 108, 191-204. | 4.9 | 63 |
| 18 | Crop Type Classification Using Fusion of Sentinel-1 and Sentinel-2 Data: Assessing the Impact of Feature Selection, Optical Data Availability, and Parcel Sizes on the Accuracies. <i>Remote Sensing</i> , 2020, 12, 2779. | 1.8 | 63 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Land cover classification with coarse spatial resolution data to derive continuous and discrete maps for complex regions. <i>Remote Sensing of Environment</i> , 2011, 115, 3264-3275. | 4.6 | 62 |
| 20 | Snow-cover variability in central Asia between 2000 and 2011 derived from improved MODIS daily snow-cover products. <i>International Journal of Remote Sensing</i> , 2013, 34, 3879-3902. | 1.3 | 61 |
| 21 | Analysis of Settlement Expansion and Urban Growth Modelling Using Geoinformation for Assessing Potential Impacts of Urbanization on Climate in Abuja City, Nigeria. <i>Remote Sensing</i> , 2016, 8, 220. | 1.8 | 59 |
| 22 | Analysis of uncertainty in multi-temporal object-based classification. <i>ISPRS Journal of Photogrammetry and Remote Sensing</i> , 2015, 105, 91-106. | 4.9 | 57 |
| 23 | Identifying Changing Snow Cover Characteristics in Central Asia between 1986 and 2014 from Remote Sensing Data. <i>Remote Sensing</i> , 2014, 6, 12752-12775. | 1.8 | 48 |
| 24 | Multi-Temporal Landsat Images and Ancillary Data for Land Use/Cover Change (LULCC) Detection in the Southwest of Burkina Faso, West Africa. <i>Remote Sensing</i> , 2015, 7, 12076-12102. | 1.8 | 48 |
| 25 | Remote Sensing and Hydrological Measurements for Irrigation Performance Assessments in a Water User Association in the Lower Amu Darya River Basin. <i>Water Resources Management</i> , 2011, 25, 2467-2485. | 1.9 | 45 |
| 26 | Satellite based calculation of spatially distributed crop water requirements for cotton and wheat cultivation in Fergana Valley, Uzbekistan. <i>Global and Planetary Change</i> , 2013, 110, 88-98. | 1.6 | 45 |
| 27 | Water balance estimation of a poorly gauged catchment in West Africa using dynamically downscaled meteorological fields and remote sensing information. <i>Physics and Chemistry of the Earth</i> , 2009, 34, 225-235. | 1.2 | 41 |
| 28 | Analysing irrigated crop rotation patterns in arid Uzbekistan by the means of remote sensing: A case study on post-Soviet agricultural land use. <i>Journal of Arid Environments</i> , 2016, 124, 150-159. | 1.2 | 38 |
| 29 | Optimising Phenological Metrics Extraction for Different Crop Types in Germany Using the Moderate Resolution Imaging Spectrometer (MODIS). <i>Remote Sensing</i> , 2017, 9, 254. | 1.8 | 34 |
| 30 | Evaluating the sequential masking classification approach for improving crop discrimination in the Sudanian Savanna of West Africa. <i>Computers and Electronics in Agriculture</i> , 2015, 118, 380-389. | 3.7 | 33 |
| 31 | Suitability of satellite remote sensing data for yield estimation in northeast Germany. <i>Precision Agriculture</i> , 2022, 23, 52-82. | 3.1 | 30 |
| 32 | Interdisciplinary Geo-ecological Research across Time Scales in the Northeast German Lowland Observatory (TERENO-NE). <i>Vadose Zone Journal</i> , 2018, 17, 1-25. | 1.3 | 29 |
| 33 | Land Use/Cover Response to Rainfall Variability: A Comparing Analysis between NDVI and EVI in the Southwest of Burkina Faso. <i>Climate</i> , 2015, 3, 63-77. | 1.2 | 28 |
| 34 | Standardized FAO-LCCS land cover mapping in heterogeneous tree savannas of West Africa. <i>Journal of Arid Environments</i> , 2010, 74, 1083-1091. | 1.2 | 27 |
| 35 | Remote sensing and hydrological measurement based irrigation performance assessments in the upper Amu Darya Delta, Central Asia. <i>Physics and Chemistry of the Earth</i> , 2013, 61-62, 52-62. | 1.2 | 27 |
| 36 | Spatial targeting of land rehabilitation: a relational analysis of cropland productivity decline in arid Uzbekistan. <i>Erdkunde</i> , 2013, 67, 167-181. | 0.4 | 27 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | On the relationship between vegetation and climate in tropical and northern Africa. Theoretical and Applied Climatology, 2014, 115, 341-353. | 1.3 | 26 |
| 38 | Multiscale Remote Sensing to Map the Spatial Distribution and Extent of Cropland in the Sudanian Savanna of West Africa. Remote Sensing, 2017, 9, 839. | 1.8 | 26 |
| 39 | Evaluation of the CropSyst model for simulating the potential yield of cotton. Agronomy for Sustainable Development, 2008, 28, 345-354. | 2.2 | 25 |
| 40 | Cropping Intensity in the Aral Sea Basin and Its Dependency from the Runoff Formation 2000â€“2012. Remote Sensing, 2016, 8, 630. | 1.8 | 25 |
| 41 | Derivation of leaf area index for grassland within alpine upland using multi-temporal RapidEye data. International Journal of Remote Sensing, 2013, 34, 8628-8652. | 1.3 | 24 |
| 42 | Assessing gaps in irrigated agricultural productivity through satellite earth observationsâ€”A case study of the Fergana Valley, Central Asia. International Journal of Applied Earth Observation and Geoinformation, 2017, 59, 118-134. | 1.4 | 24 |
| 43 | Modeling of Cotton Yields in the Amu Darya River Floodplains of Uzbekistan Integrating Multitemporal Remote Sensing and Minimum Field Data. Agronomy Journal, 2007, 99, 1317-1326. | 0.9 | 23 |
| 44 | Validation of the collection 5 MODIS FPAR product in a heterogeneous agricultural landscape in arid Uzbekistan using multitemporal RapidEye imagery. International Journal of Remote Sensing, 2012, 33, 6818-6837. | 1.3 | 23 |
| 45 | Object-based identification of vegetation cover decline in irrigated agro-ecosystems in Uzbekistan. Quaternary International, 2013, 311, 163-174. | 0.7 | 23 |
| 46 | Modelling the Gross Primary Productivity of West Africa with the Regional Biomass Model RBM+, using optimized 250 m MODIS FPAR and fractional vegetation cover information. International Journal of Applied Earth Observation and Geoinformation, 2015, 43, 177-194. | 1.4 | 23 |
| 47 | Remote sensing-based assessments of land use, soil and vegetation status, crop production and water use in irrigation systems of the Aral Sea Basin. A review. Water Security, 2020, 11, 100078. | 1.2 | 23 |
| 48 | A Comparison between Support Vector Machine and Water Cloud Model for Estimating Crop Leaf Area Index. Remote Sensing, 2021, 13, 1348. | 1.8 | 23 |
| 49 | Water Management in Khorezm: Current Situation and Options for Improvement (Hydrological) Tj ETQq1 1 0.784314 rgBT /Qoverlock 21 | | |
| 50 | Reconstructing the Spatio-Temporal Development of Irrigation Systems in Uzbekistan Using Landsat Time Series. Remote Sensing, 2012, 4, 3972-3994. | 1.8 | 21 |
| 51 | Classification of burn severity using Moderate Resolution Imaging Spectroradiometer (MODIS): A case study in the jarrah-marri forest of southwest Western Australia. Journal of Geophysical Research, 2007, 112, . | 3.3 | 19 |
| 52 | Numerical modelling and remote sensing based approaches for investigating groundwater dynamics under changing land-use and climate in the agricultural region of Pakistan. Journal of Hydrology, 2020, 581, 124408. | 2.3 | 18 |
| 53 | Modelling Crop Biomass from Synthetic Remote Sensing Time Series: Example for the DEMMIN Test Site, Germany. Remote Sensing, 2020, 12, 1819. | 1.8 | 18 |
| 54 | Spatial Transferability of Random Forest Models for Crop Type Classification Using Sentinel-1 and Sentinel-2. Remote Sensing, 2022, 14, 1493. | 1.8 | 18 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Spatio-temporal supply–demand of surface water for agroforestry planning in saline landscape of the lower Amudarya Basin. <i>Journal of Arid Environments</i> , 2019, 162, 53-61. | 1.2 | 15 |
| 56 | Statistical Exploration of SENTINEL-1 Data, Terrain Parameters, and in-situ Data for Estimating the Near-Surface Soil Moisture in a Mediterranean Agroecosystem. <i>Frontiers in Water</i> , 2021, 3, . | 1.0 | 15 |
| 57 | Important Variables of a RapidEye Time Series for Modelling Biophysical Parameters of Winter Wheat. <i>Photogrammetrie, Fernerkundung, Geoinformation</i> , 2016, 2016, 285-299. | 1.2 | 15 |
| 58 | Adapting to water scarcity: constraints and opportunities for improving irrigation management in Khorezm, Uzbekistan. <i>Water Science and Technology: Water Supply</i> , 2013, 13, 337-348. | 1.0 | 14 |
| 59 | Mapping and assessing crop diversity in the irrigated Fergana Valley, Uzbekistan. <i>Applied Geography</i> , 2017, 86, 102-117. | 1.7 | 14 |
| 60 | Optimal parameters for delineating agricultural parcels from satellite images based on supervised Bayesian optimization. <i>Computers and Electronics in Agriculture</i> , 2020, 178, 105696. | 3.7 | 14 |
| 61 | Statistical derivation of fPAR and LAI for irrigated cotton and rice in arid Uzbekistan by combining multi-temporal RapidEye data and ground measurements. <i>Proceedings of SPIE</i> , 2010, , . | 0.8 | 13 |
| 62 | Understanding Hydrological Repartitioning and Shifts in Drought Regimes in Central and South-West Asia Using MODIS Derived Perpendicular Drought Index and TRMM Data. <i>IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing</i> , 2014, 7, 983-993. | 2.3 | 13 |
| 63 | Evaluation of Sentinel-1 and Sentinel-2 Feature Sets for Delineating Agricultural Fields in Heterogeneous Landscapes. <i>IEEE Access</i> , 2021, 9, 116702-116719. | 2.6 | 12 |
| 64 | Quantifying water volumes of small lakes in the inner Aral Sea Basin, Central Asia, and their potential for reaching water and food security. <i>Environmental Earth Sciences</i> , 2016, 75, 1. | 1.3 | 10 |
| 65 | Inverse Parametrization of a Regional Groundwater Flow Model with the Aid of Modelling and GIS: Test and Application of Different Approaches. <i>ISPRS International Journal of Geo-Information</i> , 2018, 7, 22. | 1.4 | 10 |
| 66 | Cabbage whiteflies colonise <i>Brassica</i> vegetables primarily from distant, upwind source habitats. <i>Entomologia Experimentalis Et Applicata</i> , 2019, 167, 713-721. | 0.7 | 10 |
| 67 | Agro-Meteorological Trends of Recent Climate Development in Khorezm and Implications for Crop Production. , 2012, , 25-36. | | 10 |
| 68 | Consequences of chemical pretreatments in particle size analysis for modelling wind erosion. <i>Geoderma</i> , 2021, 396, 115073. | 2.3 | 9 |
| 69 | A crop type dataset for consistent land cover classification in Central Asia. <i>Scientific Data</i> , 2020, 7, 250. | 2.4 | 8 |
| 70 | Unsupervised Parameterization for Optimal Segmentation of Agricultural Parcels from Satellite Images in Different Agricultural Landscapes. <i>Remote Sensing</i> , 2020, 12, 3096. | 1.8 | 8 |
| 71 | Per-field crop classification in irrigated agricultural regions in middle Asia using random forest and support vector machine ensemble. , 2012, , . | | 7 |
| 72 | Potentials of RapidEye time series for improved classification of crop rotations in heterogeneous agricultural landscapes: experiences from irrigation systems in Central Asia. , 2011, , . | | 6 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | Stepwise Automated Pixel-Based Generation of Time Series Using Ranked Data Quality Indicators. IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing, 2011, 4, 272-280. | 2.3 | 5 |
| 74 | Biomass Assessment of Agricultural Crops Using Multi-temporal Dual-Polarimetric TerraSAR-X Data. PFG - Journal of Photogrammetry, Remote Sensing and Geoinformation Science, 2019, 87, 159-175. | 0.7 | 5 |
| 75 | Remote Sensing and Modelling Based Framework for Valuing Irrigation System Efficiency and Steering Indicators of Consumptive Water Use in an Irrigated Region. Sustainability, 2020, 12, 9535. | 1.6 | 5 |
| 76 | The Impact of Phenological Developments on Interferometric and Polarimetric Crop Signatures Derived from Sentinel-1: Examples from the DEMMIN Study Site (Germany). Remote Sensing, 2021, 13, 2951. | 1.8 | 5 |
| 77 | Digital In Situ Data Collection in Earth Observation, Monitoring and Agriculture – Progress towards Digital Agriculture. Remote Sensing, 2022, 14, 393. | 1.8 | 5 |
| 78 | Dynamics of MODIS Time Series for Ecological Applications in Southern Africa. , 2007, , . | | 4 |
| 79 | Einfluss der thematischen und räumlichen Auflösung auf die überwachte, fernerkundungsbasierte Feldfrucht-Klassifizierung.. Photogrammetrie, Fernerkundung, Geoinformation, 2015, 2015, 7-20. | 1.2 | 4 |
| 80 | Vergleich zweier statistischer Methoden zur Ableitung des Anteils absorbierter Photosynthese wirksamer Strahlung (FAPAR) für Baumwolle. Photogrammetrie, Fernerkundung, Geoinformation, 2015, 2015, 55-67. | 1.2 | 4 |
| 81 | Modelling End-of-Season Soil Salinity in Irrigated Agriculture Through Multi-temporal Optical Remote Sensing, Environmental Parameters, and In Situ Information. PFG - Journal of Photogrammetry, Remote Sensing and Geoinformation Science, 2018, 86, 221-233. | 0.7 | 4 |
| 82 | Agricultural land use mapping in the sudanian savanna of West Africa: Current status and future possibilities. , 2012, , . | | 3 |
| 83 | Object-based cropland degradation identification: a case study in Uzbekistan. , 2012, , . | | 3 |
| 84 | Unveiling Undercover Cropland Inside Forests Using Landscape Variables: A Supplement to Remote Sensing Image Classification. PLoS ONE, 2015, 10, e0130079. | 1.1 | 3 |
| 85 | Integrating ground-based and remote sensing-based monitoring of near-surface soil moisture in a Mediterranean environment. , 2019, , . | | 3 |
| 86 | Modeling seasonal actual evapotranspiration with remote sensing and GIS in Khorezm region, Uzbekistan. , 2004, , . | | 2 |
| 87 | Development of a satellite-based multi-scale land use classification system for land and water management in Uzbekistan and Kazakhstan. Proceedings of SPIE, 2011, , . | 0.8 | 2 |
| 88 | Assessing irrigated cropland dynamics in central Asia between 2000 and 2011 based on MODIS time series. Proceedings of SPIE, 2012, , . | 0.8 | 2 |
| 89 | Crop classification at subfield level using RapidEye time series and graph theory algorithms. , 2012, , . | | 2 |
| 90 | Land Surface Phenology from MODIS data in Germany. , 2013, , . | | 1 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 91 | 3-D numerical modelling of groundwater flow for scenario-based analysis and management. Water S A, 2018, 44, . | 0.2 | 1 |
| 92 | 3.6 Mapping marginal land in Khorezm using GIS and remote sensing techniques. , 2015, , 167-178. | | 1 |
| 93 | MODELLING BIOPHYSICAL PARAMETERS OF MAIZE USING LANDSAT 8 TIME SERIES. International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences - ISPRS Archives, 0, XLI-B2, 171-175. | 0.2 | 1 |
| 94 | Land Cover Analysis on Sub-Continental Scale: FAO LCCS Standard with 250 Meter MODIS Satellite Observations in West Africa. , 2008, , . | | 0 |
| 95 | A Multi-Scale Approach for Retrieving Proportional Cover of Life Forms. , 2008, , . | | 0 |
| 96 | Mapping crop distribution in administrative districts of southwest Germany using multi-sensor remote sensing data. , 2010, , . | | 0 |
| 97 | Relationships between high resolution RapidEye based fPAR and MODIS vegetation indices in a heterogeneous agricultural region. , 2011, , . | | 0 |
| 98 | A re-examination of perpendicular drought indices over Central and Southwest Asia. Proceedings of SPIE, 2012, , . | 0.8 | 0 |