

Ryuji Furihata

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2782234/publications.pdf>

Version: 2024-02-01

32
papers

699
citations

516710

16
h-index

580821

25
g-index

35
all docs

35
docs citations

35
times ranked

786
citing authors

#	ARTICLE	IF	CITATIONS
1	An Aggregate Measure of Sleep Health Is Associated With Prevalent and Incident Clinically Significant Depression Symptoms Among Community-Dwelling Older Women. <i>Sleep</i> , 2017, 40, .	1.1	78
2	The association between sleep problems and perceived health status: A Japanese nationwide general population survey. <i>Sleep Medicine</i> , 2012, 13, 831-837.	1.6	72
3	Effect of <i>Enterococcus faecalis</i> 2001 on colitis and depressive-like behavior in dextran sulfate sodium-treated mice: involvement of the brain-gut axis. <i>Journal of Neuroinflammation</i> , 2019, 16, 201.	7.2	59
4	Association of short sleep duration and short time in bed with depression: A Japanese general population survey. <i>Sleep and Biological Rhythms</i> , 2015, 13, 136-145.	1.0	38
5	Napping and associated factors: a Japanese nationwide general population survey. <i>Sleep Medicine</i> , 2016, 20, 72-79.	1.6	38
6	Seasonal changes in sleep duration and sleep problems: A prospective study in Japanese community residents. <i>PLoS ONE</i> , 2019, 14, e0215345.	2.5	37
7	Improvement of psychiatrists' clinical knowledge of the treatment guidelines for schizophrenia and major depressive disorders using the "Effectiveness of Guidelines for Dissemination and Education in Psychiatric Treatment (EGUIDE)" project: A nationwide dissemination, education, and evaluation study. <i>Psychiatry and Clinical Neurosciences</i> , 2019, 73, 642-648.	1.8	35
8	Stressful events and coping strategies associated with symptoms of depression: A Japanese general population survey. <i>Journal of Affective Disorders</i> , 2018, 238, 482-488.	4.1	33
9	Prescription patterns in patients with schizophrenia in Japan: First-quality indicator data from the survey of "Effectiveness of Guidelines for Dissemination and Education in psychiatric treatment (EGUIDE)" project. <i>Neuropsychopharmacology Reports</i> , 2020, 40, 281-286.	2.3	32
10	Characteristics of discharge prescriptions for patients with schizophrenia or major depressive disorder: Real-world evidence from the Effectiveness of Guidelines for Dissemination and Education (EGUIDE) psychiatric treatment project. <i>Asian Journal of Psychiatry</i> , 2021, 63, 102744.	2.0	28
11	A composite measure of sleep health is associated with symptoms of depression among Japanese female hospital nurses. <i>Comprehensive Psychiatry</i> , 2020, 97, 152151.	3.1	27
12	Self-help behaviors for sleep and depression: A Japanese nationwide general population survey. <i>Journal of Affective Disorders</i> , 2011, 130, 75-82.	4.1	26
13	Unhealthy lifestyle factors and depressive symptoms: A Japanese general adult population survey. <i>Journal of Affective Disorders</i> , 2018, 234, 156-161.	4.1	26
14	Association between reduced serum BDNF levels and insomnia with short sleep duration among female hospital nurses. <i>Sleep Medicine</i> , 2020, 68, 167-172.	1.6	26
15	Unmet needs of patients with major depressive disorder: Findings from the "Effectiveness of Guidelines for Dissemination and Education in Psychiatric Treatment (EGUIDE)" project: A nationwide dissemination, education, and evaluation study. <i>Psychiatry and Clinical Neurosciences</i> , 2020, 74, 667-669.	1.8	20
16	Improvements in the degree of understanding the treatment guidelines for schizophrenia and major depressive disorder in a nationwide dissemination and implementation study. <i>Neuropsychopharmacology Reports</i> , 2021, 41, 199-206.	2.3	17
17	Hypnotic medication use among inpatients with schizophrenia and major depressive disorder: results of a nationwide study. <i>Sleep Medicine</i> , 2022, 89, 23-30.	1.6	16
18	Association between the examination rate of treatment-resistant schizophrenia and the clozapine prescription rate in a nationwide dissemination and implementation study. <i>Neuropsychopharmacology Reports</i> , 2022, 42, 3-9.	2.3	14

#	ARTICLE	IF	CITATIONS
19	The characteristics of patients receiving psychotropic pro re nata medication at discharge for the treatment of schizophrenia and major depressive disorder: A nationwide survey from the EGUIDE project. <i>Asian Journal of Psychiatry</i> , 2022, 69, 103007.	2.0	12
20	Subjective assessment of participants in education programs on clinical practice guidelines in the field of psychiatry. <i>Neuropsychopharmacology Reports</i> , 2022, 42, 221-225.	2.3	12
21	A dissemination and education programme to improve the clinical behaviours of psychiatrists in accordance with treatment guidelines for schizophrenia and major depressive disorders: the Effectiveness of Guidelines for Dissemination and Education in Psychiatric Treatment (EGUIDE) project. <i>BIPsych Open</i> , 2022, 8, e83.	0.7	11
22	Clozapine Treatment Is Associated With Higher Prescription Rate of Antipsychotic Monotherapy and Lower Prescription Rate of Other Concomitant Psychotropics: A Real-World Nationwide Study. <i>International Journal of Neuropsychopharmacology</i> , 2022, 25, 818-826.	2.1	11
23	Association of serum BDNF levels and the BDNF Val66Met polymorphism with the sleep pattern in healthy young adults. <i>PLoS ONE</i> , 2018, 13, e0199765.	2.5	9
24	Prescription of Anticholinergic Drugs in Patients With Schizophrenia: Analysis of Antipsychotic Prescription Patterns and Hospital Characteristics. <i>Frontiers in Psychiatry</i> , 2022, 13, .	2.6	9
25	Sleep disturbance is associated with not only shorter sleep duration, but also longer time in bed: a Japanese general population survey. <i>Sleep and Biological Rhythms</i> , 2019, 17, 407-415.	1.0	6
26	High-dose zolpidem abuse in a patient with insomnia comorbid with major depressive disorder. <i>Sleep and Biological Rhythms</i> , 2020, 18, 155-157.	1.0	3
27	REM sleep behavior disorder predisposed by neuroleptic malignant syndrome: a case report. <i>Sleep and Biological Rhythms</i> , 2019, 17, 155-157.	1.0	2
28	Association between Typus melancholicus and depressive symptoms: A Japanese general population survey. <i>Psychiatry and Clinical Neurosciences</i> , 2020, 74, 672-673.	1.8	2
29	An Epidemiological Study on Psychosocial Factors Associated with Depression among Japanese General Adult Population. <i>Journal of the Nihon University Medical Association</i> , 2016, 75, 81-87.	0.0	0
30	Identification of Chochukei (Zhang Zhongjing) as the revolutionist of decoctions. <i>Traditional & Kampo Medicine</i> , 2020, 7, 17-23.	0.6	0
31	The Relationship between Exploratory Eye Movement Dysfunction and Psychiatric Symptoms in Schizophrenia. <i>Journal of the Nihon University Medical Association</i> , 2015, 74, 283-291.	0.0	0
32	Association of insomnia symptoms and non-restorative sleep with Typus melancholicus: a Japanese general population survey. <i>Sleep and Biological Rhythms</i> , 0, , 1.	1.0	0